

## Detection of *Enterobacter sakazakii* and other *Enterobacter* sp from Dairy Cow's Milk in Boyolali and Sleman

A.E.T.H. Wahyuni<sup>1)</sup> and T.Y. Budiarso<sup>2)</sup>

<sup>1)</sup>Department of Microbiology, Faculty of Veterinary Medicine, Gadjah Mada University, Indonesia, Yogyakarta  
email:wahyuni@yahoo.com\*

<sup>2)</sup> Faculty of Biology, Duta Wacana Christian University, Yogyakarta, Indonesia

### ABSTRACT

*Enterobacter sakazakii* is considered as an opportunistic pathogen that has been implicated in severe forms of necrotizing colitis and meningitis especially in neonates with a mortality rare varying from 40%-80%. The natural habitat *E.sakazakii* is not well understood and has been reported as frequency isolated from different environments including soil, rats, flies, milk powder factories, chocolate factories and households. A total of 100 samples were obtained from dairy cow's milk were studied. The presence of *E.sakazakii* and *Enterobacter* sp was detected using the Holt et al., 1994; Guillaume et al., 2005 and Turner et al., 2000 procedure on TSBA medium. *E.sakazakii* was not isolated from both district Sleman and Boyolali. However, *E.cloacae* was found in 33 of 75 isolates (44%) of samples from Sleman. Meanwhile 12.7% *E.cloacae* and 5.4% *E.gergoviae* was found of samples from Boyolali.

*Key words: Enterobacter sakazakii, Enterobacter sp, dairy cow's milk*

### INTRODUCTION

*Enterobacter sakazakii* and other *Enterobacter* species have caused foodborne illness through consumption of a variety of foods. *Enterobacter sakazakii* is a Gram negative, facultative, rod-shaped bacterium. *Enterobacter sakazakii* is a Gram negative, facultative, rod-shaped, non-spore forming bacterium. The organism was called "yellow-pigmented *Enterobacter cloacae*". It belongs to the family *Enterobacteriaceae* and genus *Enterobacter* that contains a number of species including *E.agglomerans*, *E.cloacae*, *E.aerogenes* and *E.gergoviae*.

The differentiation among these species is based on biochemical reactions, and serological and molecular techniques (Hoffman and Roggenkamp, 2003; Iversen et al., 2004). *E.sakazakii*, *E agglomerans*, and *E.cloacae* are considered the main species of this genus that are frequently isolated from clinical samples and food products (Farmer et al., 1980). *E.sakazakii* and *Enterobacter* species have been reported as frequently isolated from different environments including soil, rats, flies, milk powder factories, chocolate factories and households (Kandhai et al., 2004). *E.sakazakii* has been also isolated from a wide range of foods including ultra high-

temperature treated milk (UHT milk), cheese, meat, vegetables, grains, sorghum seeds, rice seeds, herbs, spices, fermented bread, fermented beverage, tofu, and sour tea (Iversen&Forsythe, 2004; Leclercq et al., 2002).

*E. sakazakii* is considered an opportunistic pathogen that has been implicated in severe forms of necrotizing colitis (Van Acker et al., 2001) and meningitis (Bar-Oz et al., 2001) especially in neonates with a mortality rate varying from 40% to 80% (Muytjens et al., 1988).

The US Food and Drug Administration (FDA, 2002) has issued an alert to health care professionals about the risk associated with *E.sakazakii* infections among neonates fed with milk-based infant formula. The alert stated that a major contribution to the avoidance of *E.sakazakii* infection in premature babies and neonates is the prevention of contamination of infant milk formula during production and bottle preparation. However, knowledge of etiological and ecological characteristic of *E.sakazakii* is sparse and its occurrence in factories that produce infant formulas and in hospital kitchens has not been studied in depth.

The natural habitat *E.sakazakii* is not well understood and has been reported as frequently isolated from different environments including