

**The Study of Genetic Diversity and Relationships on *Carica* sp.
by Means of Random Amplified Polymorphic DNA (RAPD) Analysis**

Ahmad Satori¹⁾, Sriani Sujiprihati^{1,2}, Sobir^{1,2,*} ✕

ABSTRACT

The genus of *Carica* comprises 21 species, among them three species have been domesticated and cultivated. They are *Carica papaya*, *Carica candamarcensis* and *Carica monoica*. In Indonesia, two species have been cultivated viz. *Carica papaya* and *Carica candamarcensis*. In this study, four accessions of *Carica papaya* species and one accession of *Carica candamarcensis* species were analyzed using RAPD technique. We successfully amplified a total of 40 fragments from these DNA genome by using 6 random primers with GC bases content $\geq 60\%$ of each primer. The number of fragments of each primer ranged from 5 to 8 averaged 6.7 fragments per primer. Out of total fragments, 90% showed as polymorphic ranged from 5 to 8 on average 6 fragments per primer. A dendrogram based on the UPGMA-link method using Nei and Li similarity and Principal Component Analysis (PCA) plot separated the accessions into two main groups, the *Carica papaya* species on the one side and the *Carica candamarcensis* species on the other side. The result demonstrated that RAPD analysis was able to reveal genetic difference between *Carica papaya* and *Carica candamarcensis* species, as well as genetic diversity in *Carica papaya* species.

Key words: *Carica papaya*, *Carica candamarcensis*, Random Amplified Polymorphic DNA (RAPD), Genetic diversity

INTRODUCTION

The genus *Carica* comprise 21 species, and among them, three species have been domesticated and cultivated. Those are viz. *Carica papaya* L., *Carica candamarcensis* Hook and *Carica monoica* (Nakasone and Paull, 1999). Among the three species, only the common papaya, *Carica papaya* L. has high economic value.

Indonesia is very rich in *Carica* germplasm, spreading to almost all over Indonesian archipelago and cultivated as landraces (Pusat Kajian Buah-buahan Tropika, 2000), however, information on the characterization of *Carica* is lack behind. The characterization and quantification of genetic diversity have long been a major goal in evolutionary biology. In plant breeding programs, information on genetic diversity within and among closely related crops species is essential for rational use of genetic resources and particularly useful in characterizing individual accessions and cultivars, for detecting duplications of genetic materials in germplasm collections, and as a general guide in parent's assessment for crop

improvement programs as well as developing informative mapping populations for genome mapping.

Molecular markers provide a quick and reliable method for estimating genetic relationships among genotypes of any organism (Thormann *et al.*, 1994). Random amplified polymorphic DNA (RAPD) analysis (Williams *et al.*, 1990) has been used for diversity analysis in a vast array of crops, widely used for the determination of genotypes (Hasizume *et al.*, 1993), gene mapping (Ohmori *et al.*, 1996) and QTL analysis (Grandillo and Tanksley, 1996). This approach is based on the polymerase chain reaction (PCR) (Saiki *et al.*, 1988) amplification of template DNA genome using short, synthetic deoxyribonucleotides of random sequence as primers. Each primer can direct the amplification of several unrelated regions of the genome (Sondur *et al.*, 1996). The resolving power of RAPD technique is several folds higher than visual and protein markers and is much simpler and technically less demanding than RFLP and other similar techniques (Williams *et al.*, 1990).

In this study, we employed RAPD analysis to analyse genetic diversity and relationships between

- 1) Department of Agronomy, Faculty of Agriculture, Bogor Agricultural University. Jl. Meranti, Kampus IPB Darmaga, Bogor 16680. Tel/Fax. +62-0251-629348;
- 2) Center for Tropical Fruit Studies. Kampus IPB Baranangsiang, Jl. Pajajaran, Bogor 16143. Tel +62-0251-326881, Fax. +62-0251-382201; (author for correspondence)



Carica papaya species and *Carica candamarcensis* species, and among genotypes of *Carica papaya* species.

MATERIALS AND METHODS

Plant Materials

Four accessions of *Carica papaya* species (GM, NM203, KD and DTM) and one accession of *Carica candamarcensis* species (Dieng) were used in this study. The accessions are part of Center for Tropical Fruit Studies germplasm collection.

DNA Isolation

Total DNA from each accession was isolated using CTAB method (Doyle and Doyle, 1987) with slight modification. Three hundred mg of leave tissue was frozen in liquid nitrogen and ground into a fine powder using a mortar and pestle. The pulverized materials were transferred to a microtube and 600 µl of extraction buffer [100 mM Tris-HCl pH 8.0, 20 mM EDTA, 1400 mM NaCl, 2% (w/v) cetyl trimethylammonium bromide (CTAB), and 0.2% (v/v) mercaptoethanol] solution was added. The tubes were vortexed for a few seconds and incubated at 60 °C for 20 minutes. Following incubation, 570 µl of chloroform: isoamylalcohol (24:1) (v/v) was added and mixture was shaken vigorously. The extracts were centrifuged for 15 min at 12,000 rpm. The supernatant transferred to a fresh microtube and then 600 µl of cold isopropanol was added. DNA fibers became visible upon gentle swirling. The DNA was transferred to a fresh tube, rinsed with 70% ethanol and dissolved in 50-100 µl of sterilized deionized water. DNA concentration was measured with UV-spectrophotometer (Shimadzu Corporation, Japan) at wavelengths of 260 and 280 nm.

RAPD Analysis

Six random primers of 10 bases in length with GC base content $\geq 60\%$ of each primer was selected: 3 from SB series, 2 from OPH (Operon Technologies, Inc.) series and 1 from OPM (Operon Technologies, Inc.) series (Table 1). PCR reactions were carried out in a 25 µl reaction mix containing approximately 25-50 ng template DNA, 10X PCR buffer (100 mM Tris-HCl, pH 9.0 at 25 °C, 500 mM KCl, 1.0 Triton X-100), 2 mM MgCl₂, 0.2 mM of each dATP, dCTP, dGTP and dTTP, 0.4 pmol of a single 10-base primer, and 1 unit of Taq

DNA polymerase (Gene Amp). Amplification was performed in Perkin Elmer thermal cycler programmed for 45 cycles of each of the following: 95 °C for 1 min, 35 °C for 1 min, and 72 °C for 2 min. A final elongation step of 5 min at 72 °C was included. Reaction products were mixed with 2.5 µl of loading dye (0.25% bromphenol blue, 0.25% xylene cyanol and 40% sucrose, w/v) and spun briefly in a microfuge before loading (Sambrook *et al.*, 1989), and then separated on 6% polyacrylamide gel by adding 7 M urea (17.9% (v/v) acrylamid 40% (38% acrylamid, 2% N, N, methylenediacylamid, w/v), 7.5% (v/v) 10X TBE, 1% (v/v) ammonium persulphate 10%, 0.1% (v/v) TEMED (N, N, N', N'-tetramethylethylenediamine) and 42% (w/v) urea) at 60 V for 5 h in TBE buffer (1 M Tris-HCl, 0.83 M boric acid, 10 mM EDTA, pH 8.3) and stained with ethidium bromide, and the gels were illuminated under UV light.

Data Analysis

Each accession product was considered to be a unit character, and scored for the presence (1) or absence (0) of a product. Genetic similarity (F) between all pairs of accessions were calculated according to Nei and Li (1979): $F = 2N_{ab} / (N_a + N_b)$, where N_{ab} = number of bands common in a and b, N_a = number of bands present in a, N_b = number of bands present in b. The similarity indices were converted into dissimilarity ($D = 1 - F$). The resulting dissimilarity matrix was subjected to hierarchical cluster analysis using the Unweighted Pair Group Method and Arithmetic Average (UPGMA) and was done by using computer program NTSYS-pc, version 1.80 (Exeter software, New York). A principal component analysis (PCA) subjected to separate among accessions and to determine the relationships based on the diversity and was conducted using computer program MINITAB release 11.12.

RESULTS AND DISCUSSIONS

Results

A total of 40 fragments were successfully generated using 6 random primers with GC bases content $\geq 60\%$ of each primer. The fragment number of each primer ranging from 5 to 8, on average 6.7 fragments per primer (Table 1). Out of total fragments, 90% showed as polymorphic with ranged from 5 to 8, on average was 6 fragments per primer (Table 1).

Table 1. Sequence of RAPD primers used and number of fragment amplification products

Primer	Oligonucleotide Sequence (5' to 3')	No. of total fragments	No. of polymorphic fragments
SB-05	GGGTAACGCC	7	5
SB-06	GTCCGAACCC	7	7
SB-18	TCGGCGATAG	6	6
OPM-06	CTGGGCAACT	8	8
OPH-18	GAATCGGCCA	7	5
OPH-20	GGGAGACATC	5	5

A dendrogram based on the UPGMA-link method using Nei and Li similarity (1979) and principal component analysis (PCA) plot were performed to separate and examine the relationships among the accessions (Figure 1 and Figure 2). The accessions separated into two main groups. *Carica papaya* (GM, NM203, KD and DTM) species were in one cluster and *Carica candamarcensis* (Dieng) in other group. These results indicate that *Carica papaya* and *Carica candamarcensis* have different genetic characters. KD and DTM accessions were the closest, sharing as many as 14 identical banding values. The longest distance was observed between NM203 and Dieng accessions that differed at seven loci. Among *Carica papaya* accessions, KD and DTM accessions were in one group, NM203 and GM accessions in other group.

Discussions

The results demonstrate that RAPD analysis was able to differentiate between species and could be employed to reveal genetic diversity among individual cultivars in one species. Stiles et al. (1993), indicated that RAPD could be used to establish the relationships among closely related cultivars and even individuals in at least some cultivars in papaya. Both results show that RAPD analysis was a powerful tool technique to elucidate genetic diversity between *Carica papaya* and *Carica candamarcensis* as well as among genotypes of *Carica papaya*.

A dendrogram based on UPGMA-link method using Nei and Li similarity (1979) and principal component analysis plot based on data produced from 6 primers separated the accessions into two main groups (Figure 1 and Figure 2). GM, NM203, KD and DTM accessions on the one side and Dieng accession on the other side. These results indicate that the accessions from one species have closer relationships if compared to other species.

Dieng accession is a *Carica candamarcensis* species which has different characters from *Carica papaya* species. The Dieng ripe fruits have a firm flesh, low acidity, fragrant and the cavity contains numerous seeds covered with a whitish juicy sarcotesta. This study revealed different DNA genome between *Carica papaya* species and *Carica candamarcensis* species,

nevertheless, also revealed the same banding pattern presence on both species which indicated presence of genetic relationship.

Among the *Carica papaya* species group, KD and DTM accessions show closest genetic distance. Both accessions were collected from same region (Bogor), however, they show several differences in morphological characters. Compared with KD and DTM accessions, the NM203 accession has distant genetic distance. This result could be proved based on origin and the morphological characters that differentiate among the accessions. NM203 accession come from Banten and has a purple stem, petiole and peduncle. The relationship between NM203 and DTM accessions was closer than NM203 and KD accessions, since more monomorphic banding pattern were observed between NM203 and DTM accessions (Figure 3).

This study also revealed that GM accession has the farthest genetic distance from other *Carica papaya* accessions. This difference could be distinctly on the morphological characters, and predictedly caused by different genetic background. GM accession is a S6 inbred line variety with green stem, petiole and peduncle and has a broadly toothed leaf. As a S6 inbred line variety, GM accession already has characters with high purity genetic structure and uncontaminated by another alleles from other varieties. Whereas, the other three accessions are open pollinated genotype that have an alleles contaminated probability by other genotypes. The genetic diversity on *Carica papaya* is important genetic resources, that offer possibility to be used in the future *Carica* improvement and breeding program.

ACKNOWLEDGMENTS

This study was funded by special grant from Centre for Tropical Fruit Studies of IPB through National Strategic Research (RUSNAS) for tropical fruit development to S and SS. We are also express grateful thanks to Centre for Crop Improvement Studies of IPB and Ornamental Plant Research Station of Ministry of Agriculture for providing the facilities.

REFERENCES

- Doyle, J. J., J. L. Doyle. 1987. a rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull.* 12: 13-15.
- Grandillo, S., S. D. Tanksley. 1996. QTL analysis of horticultural traits differentiating the cultivated tomato from the closely related species *Lycopersicon pimpinellifolium*. *Theor Appl Genet.* 92: 935-951.
- Hasizume, T., T. Sato, M. Hirai. 1993. Determination of genetic purity of hybrid seed in watermelon (*Citrullus lanatus*) and tomato (*Lycopersicon esculentum*) Using Random Amplified Polymorphic DNA (RAPD). *Japan J Breed.* 43: 367-375.
- Nakasone, H. Y., R. E. Paull. 1999. *Tropical Fruits.* CAB International. UK. 239-269.
- Nei, M., Li, W. H. 1979. Mathematical Model for Studying Genetics Variation in Terms of Restriction Endonucleases. *Proc Nat Acad Sci.* 74: 5269 –5273.
- Ohmori, T., M. Murata, F. Motoyoshi. 1996. Molecular characterization of RAPD and SCAR markers linked to the *Tm-1* locus in tomato. *Theor Appl Genet.* 92: 151-156.
- Pusat Kajian Buah-buahan Tropika. 2000. *Pengembangan Pepaya Unggulan Nasional.* hal. 65-83. *Dalam* Prosiding Lokakarya Riset Unggulan Strategis Nasional: Pengembangan Buah-buahan Unggulan Indonesia. Bogor.
- Saiki, R. K., D. H. Gelfand, S. Stoffel, S. J. Scharf, R. Higuchi, G. T. Horn, K. B. Mullis, H. A. Erlich. 1988. Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science.* 239:487-491.
- Sambrook, J., E. F. Fritsch, T. Maniatis. 1989. *Molecular Cloning: a Laboratory Manual.* Second Edition. Cold Spring Harbor Laboratory Press. New York.
- Sondur, S. N., R. M. Manshardt, J. I. Stiles. 1996. A genetic linkage map of papaya based on randomly amplified polymorphic DNA markers. *Theor Appl Genet.* 93:547-553.
- Stiles, J. I., C. Lemme, S. Sondur, M. B. Morshidi, R. Manshardt. 1993. Using randomly amplified polymorphic DNA for evaluating genetic relationships among papaya cultivars. *Theor Appl Genet.* 85: 697-701.
- Thormann, C. E., M. E. Ferreira, L. E. A. Camango, J. G. Tivang T. C. Osborn. 1994. Comparison of RFLP and RAPD markers to estimating genetic relationships within and among Cruciferous species. *Theor Appl Genet.* 88: 973-980.
- Williams, J. G. K., A. R. Kubelik, K. J. Livak, J. A. Rafalski, S. V. Tingey. 1990. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acid Res.* 18 (22): 6531-6535.




Figure 1. Principal component analysis plot of separation between *Carica Papaya* species and *Carica candamarcensis* species, and among genotypes of *Carica papaya* species.




Figure 2. Dendrogram constructed using UPGMA and based on Nei and Li dissimilarity values (1-F) from pairwise comparison of RAPD between *Carica papaya* species and *Carica candamarcensis* species and among genotypes of *Carica papaya* species




Figure 3. RAPD profile for two species of *Carica*, four *Carica papaya* and one *Carica candamarcensis* accessions amplified by primer SB-05 and OPH-18.




Figure 4. Dendrogram constructed using UPGMA and based on 100 RAPD profiles for five (*C. papaya* and *C. candamarcensis*) accessions. The dendrogram shows the genetic relationships between the accessions, with GM and NM203 being the most closely related, followed by KD and DTM, and Dieng being the most distinct.