

The 14<sup>th</sup> International Peat Congress  
Peatlands in Balance



Stockholm, Sweden  
June 3-8, 2012



*Proceedings*



International Peat Society | IPS/IPS

ABSTRACT	ABSTRACT NO
Land-use changes tropical peat characteristics.....	157
Mari Könönen, Jyrki Jauhiainen, Harri Vasander, Satu Repo, Peter Spetz, Suwido Limin	
Recent history of a modified peat dome, Coastal Riau, Sumatra.....	180
John Bathgate, Reddy Rachmady	
Recognizing the Complexity of Tropical Peatland Ecosystem and Management for Sustainability Purposes.....	216
Medrilzam Medrilzam, John Herbohn, Paul Dargusch, Carl Smith	
Tropical peat-swamp forest biodiversity: assessment of the present and prognosis for the future.....	230
Susan Page, Mark Harrison, Susan Cheyne, Nicholas Marchant, Nicholas Boyd, Bernat Ripoll Capilla, Marc Dragiewicz, Eric Perlett, Simon Husson	
STEM - enhancing academic capacity and awareness of the challenges and threats on tropical peatlands.....	235
Maija Lampela, Markku Larjavaara, Jyrki Jauhiainen, Uras Tantulo, Adi Jaya, Suwido Limin, Harri Vasander	
The impact of drainage and degradation on tropical peatland hydrology and its implications for effective rehabilitation.....	418
Grahame Applegate, Aljosja Hooijer, Dedi Mulyadi, Nasrul Ichsan, Marnix vander Vat	

## IX.2 Carbon balance and GHG fluxes in tropical peatlands Joint IX & X

Assessing the empirical basis of peat CO <sub>2</sub> emissions estimates from oil palm plantations on tropical peatland.....	16
Ross Morrison, Susan Page, Chris Malins, Jack Rieley, Aljosja Hooijer, Jyrki Jauhiainen	
Field observation of the tropical peat soil respiration rate under various ground water levels.....	68
Toshihide Nagano, Kazutoshi Osawa, Tomoyasu Ishida, Pisoot Vijarnsorn, Apchart Jongskul, Saiyud Phetsuk, Mariko Norisada, Takeshi Yamanoshita, Katsumi Kojima	

ABSTRACT	ABSTRACT NO
Emission of CO <sub>2</sub> and CH <sub>4</sub> from Plantation Forest of <i>Acacia crassicaarpa</i> on Peatlands in Indonesia.....	114
Basuki Sumawinata, Suwardi Suwardi, Canecio P. Munoz	
The carbon balance of tropical peatlands - a global perspective.....	151
Susan Page, Aljosja Hooijer, Jyrki Jauhiainen, Jukka Miettinen, Ross Morrison, Outi Lahteenoja, Chris Malins	
Peat Maturity and Peat Thickness for Estimation of CO <sub>2</sub> Emission from Peat Oxidation.....	175
Fahmuddin Agus, Anny Mulyani, Ai Dariah, M Maswar, W Wahyunto	
Disturbance history and management of tropical peatlands effects on N <sub>2</sub> O fluxes.....	207
Harri Vasander, Jyrki Jauhiainen, Hanna Silvennoinen, Riitta Hamalainen, Kitso Kusin, Suwido Limin, John Raision	
Carbon dioxide balance of tropical peat ecosystems.....	267
Takashi Hirano, Hendrik Segah, Kitso Kusin, Suwido Limin, Hidenori Takahashi, Mitsuru Osaki	
Net ecosystem CO <sub>2</sub> exchange of a tropical peat swamp forest in Sarawak, Malaysia..	300
Angela Che Ing Tang, Azmi Puking, Kevin Kemudang Musin, James Daniel Peter Ah-Came, Deniel Sang, Edward Baran Aeries, Ryuichi Hirata, Lulie Melling	
Seasonal variation of CO <sub>2</sub> exchange from a logged over tropical peat swamp forest in Sarawak, Malaysia.....	315
Edward Baran Aeries, Deniel Sang, Angela Che Ing Tang, Azmi Puking, Kevin Kemudang Musin, James Daniel Peter Ah-Came, Ryuichi Hirata, Lulie Melling	
Effect of fertilization on N <sub>2</sub> O emission from tropical peat: a laboratory incubation study.....	317
Aileen Kai Fang Sim, Che Fauziah Ishak, Ahmad Husni Mohd Hanif, Lulie Melling	
Is water table the most important factor influencing soil C flux in tropical peatland?..	330
Lulie Melling, Kah Joo Goh, Angelyn Kloni, Ryusuke Hatano	
Regulatory factors of soil CH <sub>4</sub> fluxes in different ages of oil palm plantation on tropical peatland in Sarawak, Malaysia.....	331
Lulie Melling, Kah Joo Goh, Auldry Chaddy, Ryusuke Hatano	
Towards a standard for deforestation of tropical peat forest.....	414
Bambang Setiadi	



EMISSION OF CO<sub>2</sub> AND CH<sub>4</sub> FROM PLANTATION FOREST OF *ACACIA*  
*CRASSICARPA* ON PEATLANDS IN INDONESIA

Basuki Sumawinata; Department of Soil Science and Land Resources, Faculty of Agriculture, Bogor Agricultural University; Darmaga Campus, 16880, Bogor, Indonesia, +62-0251-8629357, basukis2@yahoo.com

Suwardi; Department of Soil Science and Land Resources, Faculty of Agriculture, Bogor Agricultural University

Canecio P. Munoz; Sinar Mas Forestry, Indonesia

#### SUMMARY

Measurement of CO<sub>2</sub> and CH<sub>4</sub> fluxes from land surface of *Acacia crassicarpa* plantation on peatlands in Indonesia has been carried out for a period of one year at 1-weekly interval. This research was aimed to understanding process of the emission as well as the controlling factors to seek for better plantation management. The results show that CO<sub>2</sub> flux fluctuated over the year with no consistent trend with respect to water table but clearly correlates with age of planted trees. There is no clear evidence either to show that pristine peat forest has the lowest CO<sub>2</sub> emission. There are strong indications that respiration contributed greatly to the flux. Flux of CH<sub>4</sub> emission from the peatland was very low.

KEY WORDS: *Acacia crassicarpa*, Emission, Peatlands, Plantation forest.

#### INTRODUCTION

Use of peatlands for development activities has been considered as an important source of carbon emissions. In Indonesia, hence, conversion of peatland forests to commercial plantation forests such as *A. crassicarpa* planted forest is intensively assessed. Varying opinions still exist as to what is the net benefit of the planted forest. Data are needed that will be useful for determining best management practices in plantation forestry particularly in controlling environment deterioration and enhancing productivity.

This study aims to collect detailed and comprehensive information and to assess the factors and processes controlling the fluxes of CO<sub>2</sub> and CH<sub>4</sub> fluxes from *A. crassicarpa* plantation on peatlands.

#### MATERIAL AND METHODS

This research was carried out at three locations in Sumatra Island, Indonesia. The first location is in Riau Province, within an *Acacia* plantation area directly converted 8-10 years ago from pristine peat forest on deep peat. The second location is in Jambi Province also within an *Acacia* plantation but on moderate peat depth in an area that was logged previously. The third location is in South Sumatra Province, on shallow peat on an area that was denuded

and has been subjected to repeated burning and over drained. Location of plots and characteristics of each are presented in Table 1. Two to 4 different sites were selected within each location based on age of the trees and water level, with additional measurements made in adjacent pristine peatland forests at Riau Province location, in adjacent logged-over peatland forests and mineral soil planted with *Acacia mangium* and *Eucalyptus*, sp. at the Jambi Province location.

Table 1. Location and condition of measurement plots.

No	Location	Land Characteristics	Landuse/ Type of Planted Trees	Age (year)	
1	Riau Province	Deep peat (> 9m)	<i>Acacia crassicarpa</i> (Ac)	1	3
2	Riau Province	Deep peat (> 9m)	<i>Acacia crassicarpa</i> (Ac)		3-R-L
3	Riau Province	Deep peat (> 9m)	Pristine forest		
4	Jambi Province	Moderate peat (3-4m)	<i>Acacia crassicarpa</i> (Ac)	3	
5	Jambi Province	Moderate peat (3-4m)	Secondary forest		
6	Jambi Province	Mineral soil	<i>Eucalyptus pelita</i> (Ep)	2	
6	Jambi Province	Mineral soil	<i>Acacia Mangium</i> (Am)	1	
7	South Sumatera	Shallow peat (1-2m)	<i>Acacia crassicarpa</i> (Ac)	1	3
8	South Sumatera	Shallow peat (1-2m)	<i>Acacia crassicarpa</i> (Ac)		3-R-L
9	South Sumatera	Mineral soil	Abandoned paddy fields		

Note: -R-L: without fine root and litter

Measurements have been made for a one year period at 1-weekly to 2-weekly intervals, at some plots by collecting CO<sub>2</sub> and CH<sub>4</sub> gases via an acrylic cubic closed chamber having dimension of 30 x 30 x 30 cm<sup>3</sup> placed on the peat surface at 3-minute interval for CO<sub>2</sub> and 10-minute interval for CH<sub>4</sub>. Three chambers were placed in three adjacent plots at every site as replication. Concentration of the gas samples is determined using an IRGA instrument for CO<sub>2</sub> and using a GC of Shimadzu 14B with FID detector. Soil moisture content of the upper 5 cm peat was measured at every gas sampling. In all plots in peatlands water table was monitored at the same interval using perforated PVC tube inserted vertically through the peat and anchored in the underlying mineral sediment.

## RESULTS AND DISCUSSION

Results of CO<sub>2</sub> flux measurements are presented in Fig. 1. The figure shows differences between land-uses and soil types. CO<sub>2</sub> fluxes from mineral soil in one year old planted *A. mangium* and from paddy field (shallow peat have completely loss) are lower than those from the peatlands vegetated by different age of *A. crassicarpa*. The 1-weekly interval measurements show that for all the land characteristics including age of the trees, CO<sub>2</sub> fluxes over a year significantly fluctuated with maximum different between the lowest and the highest is more than 100 percent. This fluctuation indicating that the controlling factors of CO<sub>2</sub> flux need to be carefully interpreted.






Figure 1. Weekly fluxes of CO<sub>2</sub> from various conditions of *Acacia crassiorpa* planted peatlands and from adjacent pristine forest and mineral soils.

Positive correlation of CO<sub>2</sub> flux with water table is not detected at all in the Fig. 1, thus this research resulted in a contradictive finding compared to what Hatano *et al.* (2011) and Hooijer *et al.* (2011) published. Fig. 1 shows that the highest fluxes of CO<sub>2</sub> from plots with 1 year and 3 years *A. crassicarpa* occurred at week 7 but the lowest water table occurred at week 30. Similar condition was found for shallow peat (see Fig. 1 c) with 1 year *A. crassicarpa* in that the deepest water table was at week 39 until week 41 but during these weeks the CO<sub>2</sub> fluxes were the lowest. On the other hand, Fig. 1 clearly shows that the CO<sub>2</sub> flux increased with age of *A. crassicarpa* trees indicating that high proportion of CO<sub>2</sub> is released from plant respiration compared to that from peat decomposition (see comparison between Fig. 1 a and b). Another factor that is considered as a control to the flux is soil moisture. Fig. 2 shows that the highest CO<sub>2</sub> flux coincides with soil moisture between the field capacity of 250% (w/w) and the wilting point of 180% (w/w). At week 19 to week 30 when soil moisture decreased to lower than the wilting point, the CO<sub>2</sub> flux dropped to the lowest. During these weeks the water table was the lowest. Fig. 2 also shows that there is no clear correlation of water table with CO<sub>2</sub> flux, but the CO<sub>2</sub> seem to correspond with the soil moisture. Difference in CO<sub>2</sub> fluxes between that from the 3 years *A. crassicarpa* plantation and that from the similar plantation but with no fine root and litter (-R -L) reflects differences in rate of the respiration and root exudation.




Figure 2. Comparison of weekly fluxes of CO<sub>2</sub> from measurement plots with and without fine root and litter within *Acacia crassicarpa* plantation on peatland

Fluxes of CH<sub>4</sub> are presented in Fig. 3. The figure shows that the fluxes were much lower than those from paddy field. Suprihati *et al.* (2006) published that CH<sub>4</sub> flux from paddy field of mineral soils was around 7.5 mg C-CH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup>, whereas Inubushi *et al.* (2003) reported that the flux from rice plantation on peatland was 1.9 mg C-CH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup>.




Figure 3. Weekly fluxes of CH<sub>4</sub> from various conditions of *Acacia crassicaarpa* planted peatlands and from adjacent pristine forest.

Some data of CH<sub>4</sub> fluxes from the *A. crassicaarpa* plantation and from the pristine forest are presented in Fig. 3. The fluxes from the pristine forest were also low and the highest was only 13.0 mg C-CH<sub>4</sub> m<sup>-2</sup> day<sup>-1</sup> or 0.54 mg C-CH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup>. For the 1 year and 3 years *A. crassicaarpa* plantation the highest fluxes range at 9.6 – 13.2 mg C-CH<sub>4</sub> m<sup>-2</sup> day<sup>-1</sup> or 0.40 – 0.55 mg C-CH<sub>4</sub> m<sup>-2</sup> h<sup>-1</sup>. Fig. 3 also shows that there is no correlation at all of the fluxes and water table.

Year's fluxes of CO<sub>2</sub> and CH<sub>4</sub> are presented in Table 2. The table shows that CO<sub>2</sub> from the peat are higher than that from the mineral soil. However it is not necessarily that those from the deep peat are higher than those from the shallower peat. Table 2 also shows that CO<sub>2</sub> released from decomposition of litter and fine root are significant. All measurements at the condition of no fine root and litter (-R -L) show almost similar results in a range of 20.31 – 26.38 ton C-CO<sub>2</sub> ha<sup>-1</sup> y<sup>-1</sup> for both the deep and shallow peat, and the pristine forest. The highest CH<sub>4</sub> flux of all plots is only 7.29 kg C-CH<sub>4</sub> ha<sup>-2</sup> y<sup>-1</sup> that is surprisingly obtained from the -R -L measurement plot of the 3 year old *A. crassicaarpa* plantation that is never inundated.



Table 2. Yearly fluxes of CO<sub>2</sub> and CH<sub>4</sub> from different plots of *Acacia crassicarpa* planted peatlands and from adjacent pristine forest and mineral soils.

Land Characteristics	Landuse (age)	CO <sub>2</sub> Fluxes (ton C-CO <sub>2</sub> ha <sup>-1</sup> y <sup>-1</sup> )	CH <sub>4</sub> Fluxes (kg C-CH <sub>4</sub> ha <sup>-1</sup> y <sup>-1</sup> )
Mineral soil	<i>Acacia mangium</i> , 2y	20.23	0.77
Mineral soil	<i>Eucalyptus</i> , sp 2y	18.10	0.32
Mineral soil	Abandoned paddy field	15.97	-0.28
Peat soil, deep	<i>Acacia crassicarpa</i> 1y	35.77	-7.52
Peat soil, deep	<i>Acacia crassicarpa</i> 3y	52.43	3.73
Peat soil, deep	<i>Acacia crassicarpa</i> 3y -R-L	26.04	7.29
Peat soil, deep	Pristine forest	33.04	4.39
Peat soil, deep	Pristine forest -R-L	20.31	4.33
Peat soil, moderate	<i>Acacia crassicarpa</i> 3y	34.31	2.20
Peat soil, moderate	Secondary (logged-over) forest	36.52	0.24
Peat soil, shallow	<i>Acacia crassicarpa</i> 4y	37.59	2.43
Peat soil, shallow	<i>Acacia crassicarpa</i> 4y- R-L	26.38	1.12

Note: -R-L: without fine root and litter

## CONCLUSION

The fluxes of CO<sub>2</sub> varied with land condition and fluctuated over a year but did not show a consistent trend with respect to peat depth, land-use type, and water table, but correlated with soil moisture and age of the *A. crassicarpa* trees. There is no clear evidence either to show that pristine peat forest has the lowest CO<sub>2</sub> emission or that the lowest water table has the highest CO<sub>2</sub> emission. There are strong indications that respiration contributed greatly to the flux.

The fluxes of CH<sub>4</sub> fluctuated over a year and varied with land condition but the range were low if compared to the fluxes from paddy fields.

## ACKNOWLEDGEMENTS

We thank PT. Bukit Batu Hutani Alam, PT. Sekato Pratama Makmur, PT. Wirakarya Sakti, PT. Sebangun Bumi Andalas, Sinar Mas Forestry Management and APP Indonesia for giving permission to conduct this research on their plantation forest concession areas and financial support.

## REFERENCES

Hatano, R., Inoue, T., Yamada, H., Sato, S., Darung, U., Limin, S., June, T., Suwardi, Sumawinata, B., Wijaya, H., Wijonarko, S. and Bathgate, J. 2011. Soil greenhouse gases

emission from various land uses in tropical peatlands in Indonesia. *Proceeding of Palangkaraya International Symposium and Workshop on Tropical Peatland*, Palangkaraya, Indonesia, June 10-11, 2010.

Hooijer, A., Page, S., Jauhiainen, J., Lee, W.A., Lu, X. X., Idris, A. and Anshari, G. 2011. Subsidence and Carbon loss in drained tropical peatlands: reducing uncertainty and implication for CO<sub>2</sub> emission reduction options. *Biogeoscience* 8, 9311-9356, doi:10.5194/bgd-8-9311-2011.

Inubushi, K., Furukawa, Y., Hadi, A., Purnomo, E. and Tsuruta, H. 2003. Seasonal changes of CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O fluxes in relation to land-use change in tropical peatlands located in Coastal area of South Kalimantan. *Chemosphere* 52:603-608.

Suprihati, Anas, I., Murdiyarso, D., Sabiham, S. and Djajakirana, G. 2006. Methane Flux and Soil Characteristic in Several Cropping Systems. *Buletin Agronomy*. (34) (3) 181-187.