




USAID
FROM THE AMERICAN PEOPLE



Vegetable-Agroforestry Systems in Indonesia



Editors: Anas D. Susila, Bambang S. Purwoko, James M. Roshetko, Manuel C. Palada, Juang G. Kartika, Lia Dahlia, Kusuma Wijaya, Arif Rahmanulloh, Mahmud Raimadoya, Tri Koesoemaningtyas, Herien Puspitawati, Tisna Prasetyo, Suseno Budidarsono, Iwan Kurniawan, Manuel Reyes, Wanraya Suthumchai, Karika Kunta and Samran Sombatpanit

7. Optimum Fertilizer Rate for Kangkung (*Ipomoea aquatica* L.) Production in Ultisols of Nanggung, Bogor

Anas D. Susila¹, Tisna Prasetyo¹ and Manuel C. Palada²

Abstract

Nitrogen, phosphorus and potassium availabilities are the most limiting factors for maximum growth and yield. Kangkung (*Ipomoea aquatica* L.) is an important traditional leafy vegetable crop cultivated in Indonesia. Kangkung was grown to evaluate the optimum rates of N, P and K fertilizers in Ultisols of Nanggung, Bogor, with low pH (5.5), low organic C (1.54%), very low total N (0.12%), low K content (0.29 me/100 g), but very high soil P₂O₅ concentration (19.2 ppm) to evaluate the best crop management practices with starter solution. The experiment was conducted in Hambaro village, Nanggung subdistrict, Bogor, Indonesia, from January to April 2008. The treatments were: N, P, K fertilizer rates of 0%, 50%, 100%, 150% and 200% from the fertilizer recommendation rates (100 kg/ha N, 135 kg/ha P₂O₅ and 135 kg/ha K₂O). This experiment used Randomized Complete Block Design with four replications (each farmer field as one replication). Total number of plots was 15 x 4 = 60, with plot size of 1.5 x 5 m. Kangkung (local variety) was planted in four rows per plot, 25 cm between rows and 15 cm within rows, at 10 seeds per planting. The application of N (200 kg/ha), P₂O₅ (270 kg/ha), and K₂O (270 kg/ha) quadratically increased total and relative yields of kangkung. Based on $Y = -0.0021x^2 + 0.572x + 56.857$ for N, $Y = -0.0013x^2 + 0.3673x + 72.102$ for P₂O₅, and $Y = -0.0001x^2 + 0.0959x + 84.102$ for K₂O, the optimum rates for each nutrient was 136-141-674 kg/ha N-P₂O₅-K₂O. The fertilizer recommendation based on K threshold (no K) was 41-40-0, on P threshold was 24-0-0 kg/ha N-P₂O₅-K₂O. However, no fertilizer was needed on N threshold. In the recommendation based on optimum yield (136-141-674), the percentage increase in cost (134.0) was higher than the expected increase in yield (19.28). Based on the yield vs. cost rule therefore, the most economical recommendation would be 41-40-0 kg/ha N-P₂O₅-K₂O (K threshold).

Keywords: Kangkung, optimum fertilizer rate, Ultisols

¹Plant Production Division, Department of Agronomy and Horticulture, Faculty of Agriculture, Bogor Agricultural University, Jl. Meranti, Komplek IPB Darmaga, Bogor 16680, Indonesia. anasdsusila@ipb.ac.id, tisnaprasetyo@yahoo.com, Tel/Fax: 0251 629353

²Crop and Ecosystems Management Unit AVRDC-World Vegetable Center. P.O. Box 42, Shanhua, Tainan 74199, Taiwan. mpalada@gmail.com

1. Introduction

Kangkung (*Ipomoea aquatica* L.) is an important traditional leafy vegetable crop cultivated in Indonesia. Practically all parts of the young plants are eaten. Since older stems become fibrous, young succulent tips are preferred. Kangkung belongs to the family Convolvulaceae which can be grown in low land and upland farms. It is preferred for cultivation because of its shorter growth cycle, it is fast growing, widely adaptable, and tolerant to diseases. Kangkung usually grows in home gardens, but is now becoming one of the significant commercial vegetables. In Indonesia, yield potential of kangkung has not yet been fully exploited or studied due to low use of inputs and lack of information on production technology.

In Indonesia, growing vegetable crops on Ultisols is not widely observed, although Ultisols occupy almost 25% of total Indonesian land surface. The major problems in Ultisols are the deficiency of plant nutrients such as phosphorous (P) and potassium (K). Ultisols have an acidic to very acidic reaction, high Al and Fe saturation which are specific properties that restrict plant growth. The presence of argillic horizon in the soil influences soil physical properties such as reduction of both macro- and micro-pores, encouraging surface run-off and finally supporting soil erosion. Most studies indicated that liming, and fertilizing by organic and inorganic fertilizers could overcome some constraints in Ultisols (Prasetyo and Suriadikarta, 2006; Kasno et al., 2006).

Fertilization is one of the management practices that can be implemented to increase vegetable yield. Nitrogen and potassium are fundamental to achieve high marketable yield while phosphorus is essential for early growth and root development. The importance of potassium in ensuring normal growth and production of quality fruits is well recognized (IPNI, 2008). Nitrogen is the nutrient needed in largest quantities by plants and the one most frequently applied as fertilizer. Application rates are critical, because too much or too little directly impacts crop growth. Correct form of nitrogen is critical where the ammonium form can restrict growth and adversely affect quality. Application rates of phosphorus and potassium depend on the potential level of nutrient availability from the soil. Excessive phosphorus fertilizer can aggravate iron and zinc deficiencies and increase soil salt content. Excessive potash fertilizer can increase soil salt content too (Whiting et al., 2007).

Rochayati et al. (1999) reported that until now, fertilization practices observed by most farmers are applied to all plants, whereas fertilization rates depend on plant species and variety, soil type, location and agricultural practices. Fertilization recommendations are crop-specific and location-specific.

Proper fertilization of a crop is a requirement to obtain maximum yield. In addition to yield obtained, quality of the commodity is an important factor of profit and shelf life for high-value crops (Hochmuth et al., 1993).

This research was conducted to evaluate optimum rates of N, P and K fertilizers on kangkung (*Ipomoea aquatica* L.) grown in Ultisols of Nanggung (Jasinga soil) with low pH (5.5), low organic C (1.54%), very low total N (0.12%), low K content (0.29 me/100 g) but very high soil P₂O₅ concentration (19.2 ppm).

2. Materials and Methods

The experiment was conducted at Hambaro village, Nanggung, Bogor, Indonesia, from January to April 2008. The soil type in the location belongs to Ultisols, which typically have pH 5.5 and high P-fixation by aluminum, soil P₂O₅ concentration of 19.2 ppm, total N of 0.12%, and K content of 0.29 me/100 g. Treatments: N, P, K fertilizer rates of 0%, 50%, 100%, 150% and 200% from fertilizer recommendation rates (100 kg/ha N, 135 kg/ha P₂O₅ and 135 kg/ha K₂O). This experiment used Completely Randomized Block Design with four replications (each farmer field as one replication). Total number of plots used was 15 x 4 = 60 plots, with plot size = 1.5 x 5 m. Local variety of kangkung was planted in four rows per plot or per bed, 25 cm between rows and 15 cm within rows, 10 seeds per planting hole.

Lime (CaCO₃) was thoroughly incorporated (1.5 ton/ha) into the bed 4 weeks before planting. Furrow irrigation was carried out at 1-week intervals and weeding done when necessary. Harvesting was carried out 4 weeks after sowing.

Fertilizer application consisted of the following:

1. N fertilizer optimization: N rate was the same as the treatments, pre-plant 100% P, 50% K, side-dress 50% K (3 weeks after planting);
2. P fertilizer optimization: P rate was the same as the treatments, pre-plant 50% N and 50% K; side-dress 50% N and 50% K (3 weeks after planting);
3. K fertilizer optimization: K rate was the same as the treatments, pre-plant 100% P, 50% N; side-dress 50% N (3 weeks after planting).

Plant height and plant diameter were measured 1, 2, 3 and 4 weeks after transplanting. Yield per plant and yield per plot were measured 5 weeks after transplanting (WAT). Analysis of variance of data was calculated using SAS 8.12 (SAS Institute, NC). Polynomial regression was used to analyze N-P-K rate effect (linear or quadratic) and to find out the optimum rate for maximum yield. Economic evaluation was done to arrive at the recommendation choices.

3. Results and Discussion

Application of N, P, K fertilizers up to 200 kg/ha N, 270 kg P₂O₅ and 200 kg K₂O significantly increased vegetative growth and plant height and stem diameter of kangkung bean from 1 week to 5 weeks after planting. This range of fertilizer rates was appropriate to build optimum rate of each fertilizer in the Ultisols with soil total N of 0.12%, P₂O₅ concentration of 19.2 ppm, and K content of 0.29 me/100 g.

3.1 Plant height

Application of N fertilizer from 0 to 200 kg/ha N increased quadratically the plant height of kangkung at 1 week to 5 weeks after planting (Table 1). A similar effect was achieved with P fertilizer application, where application of P fertilizer from 0 to 270 kg/ha P₂O₅, and K fertilizer from 0 to 200 kg/ha K₂O increased quadratically the plant height of kangkung at 1 week to 5 weeks after planting (Tables 2 and 3).

Table 1. Effect of N rates on plant height of kangkung

| N rate (kg/ha N) | Plant height (cm) | | | |
|------------------|-------------------|-------|-------|-------|
| | 1 WAT | 2 WAT | 3 WAT | 4 WAT |
| 0.00 | 3.38 | 6.30 | 10.16 | 21.84 |
| 50.00 | 4.29 | 8.01 | 12.95 | 27.78 |
| 100.00 | 5.41 | 10.13 | 16.40 | 35.15 |
| 150.00 | 5.42 | 10.12 | 16.34 | 35.06 |
| 200.00 | 4.98 | 9.30 | 15.02 | 32.22 |
| Regression | Q** | Q** | Q** | Q** |

Note: **significant at P = 0.01, Regression Q = Quadratic

Table 2. Effect of P rate on plant height of kangkung

| P rate (kg/ha P ₂ O ₅) | Plant height (cm) | | | |
|-----------------------------------------------|-------------------|-------|-------|-------|
| | 1 WAT | 2 WAT | 3 WAT | 4 WAT |
| 0.00 | 3.68 | 6.87 | 11.10 | 23.82 |
| 67.50 | 4.21 | 7.86 | 12.69 | 27.23 |
| 135.00 | 4.91 | 9.16 | 14.78 | 31.74 |
| 202.50 | 4.77 | 8.91 | 14.39 | 30.88 |
| 270.00 | 4.69 | 8.77 | 14.16 | 30.39 |
| Regression | Q** | Q** | Q** | Q** |

Note: ** significant at P = 0.01, Regression Q = Quadratic

Table 3. Effect of K rate on plant height of kangkung

| K rate (kg/ ha K ₂ O) | Plant height (cm) | | | |
|-------------------------------------|-------------------|-------|-------|-------|
| | 1 WAT | 2 WAT | 3 WAT | 4 WAT |
| 0.00 | 4.26 | 7.95 | 12.84 | 27.56 |
| 50.00 | 4.29 | 8.01 | 12.92 | 27.75 |
| 100.00 | 4.60 | 8.59 | 13.86 | 29.76 |
| 150.00 | 4.86 | 9.08 | 14.65 | 31.46 |
| 200.00 | 4.76 | 8.89 | 14.35 | 30.80 |
| Regression | Q** | Q** | Q** | Q** |

Note: ** significant at P = 0.01, Regression Q = Quadratic

3.2 Stem diameter

A similar pattern of fertilizer effect occurred in stem diameter, where N, K fertilizer resulted in quadratic response, and P fertilizer resulted in linear response. Application of N fertilizer from 0 to 200 kg/ha N increased quadratically the stem diameter of kangkung at 1 week to 5 weeks after planting (Table 4) while application of P fertilizer from 0 to 270 kg/ha P₂O₅ linearly increased stem diameter of kangkung at 1 week to 5 weeks after planting (Table 5). Application of K fertilizer from 0 to 270 kg/ha K₂O increased quadratically the stem diameter of kangkung at 1 week to 5 weeks after planting (Table 6).

Table 4. Effect of N rate on stem diameter of kangkung

| N rate (kg/ ha N) | Stem diameter (cm) | | | |
|----------------------|--------------------|-------|-------|-------|
| | 1 WAT | 2 WAT | 3 WAT | 4 WAT |
| 0.00 | 0.492 | 0.918 | 1.481 | 3.179 |
| 50.00 | 0.534 | 0.998 | 1.612 | 3.460 |
| 100.00 | 0.553 | 1.033 | 1.670 | 3.581 |
| 150.00 | 0.580 | 1.085 | 1.754 | 3.762 |
| 200.00 | 0.590 | 1.102 | 1.779 | 3.817 |
| Regression | Q** | Q** | Q** | Q** |

Note: ** significant at P = 0.01, Regression Q = Quadratic

Table 5. Effect of P rate on stem diameter of kangkung

| P rate (kg/ ha P ₂ O ₅) | Stem diameter (cm) | | | |
|---------------------------------------------------|--------------------|-------|-------|-------|
| | 1 WAT | 2 WAT | 3 WAT | 4 WAT |
| 0.00 | 0.529 | 0.987 | 1.594 | 3.421 |
| 67.50 | 0.541 | 1.011 | 1.632 | 3.504 |
| 135.00 | 0.567 | 1.059 | 1.710 | 3.669 |
| 202.50 | 0.606 | 1.132 | 1.827 | 3.922 |
| 270.00 | 0.619 | 1.157 | 1.869 | 4.010 |
| Regression | L** | L** | L** | L** |

Note: ** Significant at P = 0.01, Regression L = Linear

Table 6. Effect of K rate on stem diameter of kangkung

| K rate (kg/ ha K ₂ O) | Stem diameter (cm) | | | |
|-------------------------------------|--------------------|-------|-------|-------|
| | 1 WAT | 2 WAT | 3 WAT | 4 WAT |
| 0.00 | 0.485 | 0.906 | 1.459 | 3.135 |
| 50.00 | 0.506 | 0.945 | 1.523 | 3.273 |
| 100.00 | 0.611 | 1.142 | 1.842 | 3.955 |
| 150.00 | 0.625 | 1.167 | 1.884 | 4.043 |
| 200.00 | 0.597 | 1.116 | 1.802 | 3.367 |
| Regression | Q** | Q** | Q* | Q** |

Note: ** Significant at P = 0.01, Regression Q = Quadratic

3.3 Plant yield

Application of N, P fertilizers from 0, 50%, 100%, 150%, 200% of the recommended rate resulted in quadratic response on the yield of kangkung. However, application of K fertilizer from 0, 50%, 100%, 150%, 200% of recommendation rate resulted in linear response on the yield of kangkung (Table 7). Application of K of more than 200% of the recommended rate (>270 kg/ha K₂O) still increased yield. However, application of N and P fertilizers at more than 200% of the recommended rate (>200 kg/ha N, and >270 kg/ha K₂O) reduced yield.

3.4 Multi-nutrient response interpretation

Multi-nutrient response interpretation is one method to develop fertilizer recommendations using single-nutrient quadratic model. The recommendation choice was developed using N, P, K fertilizer response curve, where the

Table 7. Effect of fertilizer rate on the yield of kangkung

| Recommended rate (%) | Yield | | |
|----------------------|--------|-------------------------------|------------------|
| | N | P ₂ O ₅ | K ₂ O |
| | Ton/ha | | |
| 0 | 2.507 | 2.720 | 3.200 |
| 50 | 3.227 | 3.173 | 3.200 |
| 100 | 4.107 | 3.680 | 3.413 |
| 150 | 4.160 | 3.653 | 3.733 |
| 200 | 3.760 | 3.547 | 3.600 |
| Response | Q** | Q** | L** |

Note: ** Non significant or significant at P = 0.01, Regression L = Linear, Q = Quadratic

100% rate = 100 kg/ha N, 135 kg/ha P₂O₅ and 135 kg/ha K₂O

first recommendation was calculated from the optimum relative yield, and the second, third and fourth recommendations determined from N, P, K threshold (0 application), respectively.

The N curve response regression equation was:

$$Y = -0.0021x^2 + 0.572x + 56.957; R^2=0.9341$$

and the optimum N rate was 136 kg/ha N.

Phosphor response regression equation was:

$$Y = -0.0013x^2 + 0.3673x + 72.102; R^2=0.9569$$

and the optimum P rate was 191 kg/ha P₂O₅.

Potassium response regression equation was:

$$Y = -0.0001x^2 + 0.0959x + 84.102; R^2=0.7649$$

and the optimum K rate was: 647 kg/ha K₂O.

Therefore, 4 fertilizer recommendations for kangkung can be applied (kg/ha N-P₂O₅-K₂O):

- Based on optimum yield = 136-191-647
- Based on N threshold = 0-0-0
- Based on P threshold = 24-0-0 and
- Based on K threshold = 41-40-0 (Fig. 3).

3.5 Economic evaluation of fertilizer recommendations

The economic evaluation of fertilizer recommendation was calculated based on the price of urea (45% N), SP36 (36% P₂O₅) and KCl (50% K₂O) which were Rp1,700, Rp3,000, and Rp8,000, respectively. Based on the increases in relative yield, the recommendation based on optimum yield was the best choice (100%).




Figure 1. Organic vegetable at home garden




Figure 2. Organic vegetable at home garden under tree system






Figure 3. Multi-nutrient response interpretation and development of fertilizer recommendation using single-nutrient quadratic model: kangkung in Ultisols at Nanggung, Bogor. Fertilizer applied (kg/ha N-P₂O₅-K₂O) for recommendation choice: to optimum yield = 161-208-309, to N threshold = 0-4-0, to K threshold = 15-28-0, to P threshold = 0-0-0.

Table 8. Economic Evaluation of Fertilizer Recommendation for Kangkong in *Ultisols* Nanggung-Bogor.

| Fertilizer Recommendation Choice | Yield Data | | | Cost Data | | | | Relative unit Cost ³⁾ |
|----------------------------------|-------------------------------------------|----------------------------------------------|--------------------------------------------|-----------------|-----------------------|----------------------------------------------|-------------------------------------------|----------------------------------|
| | Relative Yield at each nutrient threshold | Change from next lower recommendation choice | | Fertilizer Cost | Total Production Cost | Change from next lower recommendation choice | | |
| | | Increase in Relative Yield | Percentage increase in Yield ¹⁾ | | | Increase in Cost | Percentage increase in Cost ²⁾ | |
| | | | (%) | (Rp) | (Rp) | (Rp) | (%) | |
| 0 - 0 - 0 (N-threshold) | 56.90 | - | - | 0 | 7,156,000 | - | - | 125,764 |
| 24 - 0 - 0 (P-threshold) | 72.10 | 15 | 26.71 | 92,367 | 7,248,367 | 92,367 | 1.3 | 100,532 |
| 41 - 40 - 0 (K-threshold) | 84.10 | 12 | 16.64 | 488,986 | 7,644,986 | 396,619 | 5.5 | 90,904 |
| 136 - 191 - 647 (optimum) | 100.00 | 16 | 19.28 | 10,734,776 | 17,890,776 | 10,245,790 | 134.0 | 178,344 |

1) Increase in relative yield divided by the relative yield at each nutrient threshold

2) Increase in cost divided by total production cost for previous recommendation choice

3) Total production cost divided by relative yield at each nutrient threshold

However this recommendation also caused the relative cost unit to be at 178,344. Based on the fertilizer price the most economic fertilizer recommendation was 41-40-0 (kg/ha N-P₂O₅-K₂O) which had the lowest relative unit cost (90,904). However, this choice leads to only 84.10% relative yield. The economic evaluation of fertilizer recommendations for kangkung in Ultisols (Jasinga soil) is shown in Table 8.

4. Conclusions

From this study it can be concluded that:

1. The fertilizer recommendation for kangkung in Ultisols Nanggung-Bogor with soil P₂O₅ concentration of 19.2 ppm, N-total of 0.12%, and K content of 0.29 me/100 g based on optimum yield was 136-191-647 (kg/ha N-P₂O₅-K₂O).

2. The most economic fertilizer recommendation for kangkung in Ultisols of Nanggung, Bogor was 41-40-0 (kg/ha N-P₂O₅-K₂O) However, this choice resulted in 84.10% relative yield only.

5. Acknowledgments

This publication was made possible through support provided by the United States Agency for International Development and the generous support of the American People (USAID) for the Sustainable Agriculture and Natural Resources Management Collaborative Research Support Program (SANREM CRSP) under terms of Cooperative Agreement Award No. EPP-A-00-04-00013-00 to the Office of International Research and Development (OIREL), at Virginia Polytechnic Institute and State University (Virginia Tech).

References

- International Plant Nutrition Institute (IPNI). 2008.** *Potassium Improves Yield and Quality of Tomato in the Red & Lateritic Soils of West Bengal with Low Inherent Fertility.* <http://www.ppi-ppic.org>.
- Prasetyo B.H. and D.A. Suriadikarta. 2006.** Karakteristik, potensi, dan teknologi pengelolaan tanah Ultisol untuk pengembangan pertanian lahan kering di Indonesia. *Jurnal Litbang Pertanian*, 25(2): 39-46.
- Kasno A., D. Setyorini and E. Tuberkih. 2006.** Pengaruh pemupukan fosfat terhadap produktivitas tanah Inceptisol dan Ultisol. *Jurnal Ilmu-ilmu Pertanian Indonesia*, 8 (2): 91-98.
- Rochayati R., D. Setyorini, S. Suping and L.R. Widowati. 1999.** Korelasi uji tanah hara P dan K. Laporan Bagian Proyek Penelitian Sumber daya Lahan. Puslittanak (unpublished).
- Whiting D., C. O'Meara and C. Wilson. 2007.** *Vegetable Garden: Soil Management and Fertilization.* Colorado State University Extension.

Appendix: Photos from the project area in Nanggung: (upper) Visit to an organic vegetable farm and (lower) Indonesian, Philippine and Vietnamese teams during no-tillage and drip irrigation training at Nanggung, Bogor, Indonesia

