

CURRENT ISSUES OF TROPICAL PEATLAND IN INDONESIA

Basuki Sumawinata and Darmawan

Department of Soil Science and Land Resources, Faculty of Agriculture,
Bogor Agricultural University
Email: basukis2@yahoo.com

ABSTRACT

Indonesia has peatland occupying over 20 million hectares, about 10% of its total land areas, spreading in Sumatra, Kalimantan, and Papua islands. Quite vast expanses of the peatlands have been reclaimed by government for rice cultivation in transmigration projects and by private companies for forest and oil palm plantations. This paper discusses topics of carbon stock and emission in conjunction to wise utilization of peatlands in Indonesia. Discussed topics include figures of estimate of carbon stock and emission and the approaches as well, to seek for the truth if Indonesia is really in third position of carbon emitter countries, by reviewing some related publications being widely referred. Following the discussion, reviews on peatland utilizations have been made in Indonesia are described to frankly show any successful and unsuccessful of the utilizations. All the discussions and reviews are always based on a pure scientific thinking, hindering any noise of certain interest.

Keywords: *tropical peatland, carbon stock, CO₂ emission, oil palm plantation, forest plantation.*

INTRODUCTION

Indonesia possesses a considerable expanse of peatlands that mostly spreading in Sumatra, Kalimantan, and Papua islands. The expanse counts to approximately 20 million hectares, about 10% of the total lands of the country or about 6-7% of the total peatlands of the world. The Indonesian peatlands, however, count as the largest expanse of tropical peat among other tropical countries including Malaysia that possess peatlands of about 2.5 million hectares at second position.

Tropical peat has great difference with that of temperate peat in that it composed of woody organic decay instead of sphagnum material. This difference remarkably determines other characteristics; such as bulk density. Tropical peat has a much lower bulk density than that of temperate peat. The difference is certainly important to be well considered. Some of programs of utilizing peatlands in Indonesia that had been undergone with technical supervision from peat scientists of the temperate countries had mostly failed; resulted in severe environmental problems instead, such as fire and its carbon emission as well. The failure is, among others, certainly attributable to a lack of consideration on nature and properties of the tropical peatlands.



Utilization of peatland in Indonesia was started at 18th century by people of two ethnic groups, i.e. Bugis from South Sulawesi and Banjar from South Kalimantan who had utilized some of the peatlands at eastern coast of Sumatra island for growing rice as well as various annual crops and coconut. Cities of Tembilahan, Kuala Tungkal and Kuala Enok, for example, were originally peat swamp forests where the two ethnic people had come to and started living in with the above agricultural activities. This fact had become an inspiration for Indonesian government to utilize peatlands for expansion of paddy field since other flat and water abundant areas were already limited.

Development of paddy fields on peatlands had started expansively by Indonesian government in 1969 through the so-called transmigration program. One of the main purposes of this program is to resettle poor and landless people from dense Java and Bali into much less dense areas in other islands like Sumatra, Kalimantan and Sulawesi. Unfortunately the peatland development projects have mostly ended up at failure including the most recent big one, a one million hectares Mega Rice Project (MRP) in Kalimantan in 1997. Productive rice fields intended by the project could not be eventually created; the lands were drastically changed into worse condition instead. Peat layers that have original 2-3 m thickness at the upper part of the land was after a view years land cultivation found as much more tinny layers or even are not exist anymore. Disappearance of peat layers had led former underlain mineral sediment emerging to the surface. Sulphidic material containing sediment, which has a potency to form acid sulfate soil, and quartz sand dominated sediment are dominant types of peat underlain sediments of coastal peatlands. The lost of peat was mainly due to an intensive decomposition in addition to fires which are attributable to wrong drainage system, in that the system had resulted in an exsessive drying of the peat. In addition to such expansive failure of the MRP, at about the same years there were also significant numbers of utilization of peatlands for plantation of perennial crops, where fire was intensively used in land preparations. These fires have resulted in huge haze that certainly have endangered people health and disturbed transportation systems of not only Indonesia but also the neighboring countries, i.e. Malaysia, Singapore, and Brunei.

The MRP being repeatedly used by world experts of peat and environmentalists, as a tremendous example of deteriorating effect of utilizing peatlands, in criticizing Indonesia for the expansive use of peatlands. Their critic emphasized on carbon emission. With this critic they are arguing against utilization of peatlands in Indonesia for agriculture including fast growing tree plantation for pulp industries and especially oil palm plantation. An estimation made by Page *et al.* (2001) that was reanalyzed by Hooijer *et al.* (2006) shows that the carbon emission from burned Indonesian peatlands is about 3 Gt CO₂ yearly. This figure put Indonesia at third position of the biggest carbon emitter under the US at the top and China at second position.

This paper discusses topics of carbon stock and emission and wise utilizations of peatlands for forest and oil palm plantations with respect to especially water management and peat subsidence.



Carbon Stock and Emission

Estimations of the expanse of total peatland areas in Indonesia show different results that vary between 16-26 million hectares (Table 1). This variation reflects differences of definition and method of the estimations.

Table 1. Distribution and area of peatlands in Indonesia (Wetlands International. 2006)

Author/Sources	Peatland distribution and area (million ha)				Total
	Sumatra	Kalimantan	Papua	Other	
Driessen (1987)	9.70	6.30	0.10	-	16.10
Puslittanah (1981)	8.90	6.50	10.90	0.20	26.50
Euroconsult (1984)	6.84	4.93	5.46	-	17.20
Soekardi & Hidayat (1988)	4.50	9.30	4.60	<0.10	18.40
Deptrans (1988)	8.20	6.80	4.60	0.40	20.10
Subagyo <i>et al.</i> (1990)	6.40	5.40	3.10	-	14.90
Deptrans (1990)	6.90	6.40	4.20	0.30	17.80
Nugroho <i>et al.</i> (1992)	4.80	6.10	2.50	0.10	13.50*
Radjaguguk (1993)	8.25	6.79	4.62	0.40	20.10
Dwiyono & Racman (1996)	7.16	4.34	8.40	0.10	20.00
Wetlands International (2002)	7.20	5.80	8.00	-	21.00

*not including peatland associated with saline land and floodplain (2.46 milion ha)

By delineating peat distribution based on peat depth and decomposition stage using remote sensing image, which was completed with a ground check for peat depth, decomposition stage and bulk density at some points representing each delineation unit, and a laboratory analyses of carbon content of samples representing each peat decomposition stage (fibric, sapric, hemic), Wahyunto and Suradiputra (2008) have estimated carbon stock of peat in Indonesia and showed that the stock in Sumatra and Kalimantan counts to a total of 33,558 Gt. Since the image was supposed to be used as an aid in delineating spatial distribution based on reflection of surface characteristics of the peatlands, then the delineation does not necessarily coincide with distribution of the peat depth, decomposition stage, and bulk density. This inaccuracy, suggests that the above estimation approach, as well as the results, is scientifically unacceptable. Yet the approach and the results seem to be used as reference ever since and no more scientifically acceptable approach has been developed.

The MRP has been coincided with long period of dry season of El-Nino in 1997 so that the fires were continuous and extending during the year. Carbon emission in Indonesia has become a great concern ever since this prolonged fires. Page *et al.* (2001) have made a calculation to estimate carbon emission related to the MRP fires based on field data of a block named Block C having total area of 383,800 hectares with dominating peatland area of 337.632 hectares (Table 2). The fires have damaged 48% the peatland area equivalent to 54.7% of the total areas of this block. The calculated total carbon loss of this block is 0.06-0.07 Gt. The same calculation was done for the entire MRP area (988,568 hectares) and of the whole study area (area of 2,491,619 hectares including the MRP area and surroundings). Carbon loss of each is respectively 0.15-0.18 Gt and 0.24-0.28Gt.



Table 2. Effects of the 1997 fires on peat carbon stores (Page *et al.*, 2001)

	Block C of MRP, 383,800 ha	Entire MRP, 988,568 ha	Study Area, 2,491,619 ha	Extrapolated values for whole of Indonesia, 190,400,000 ha		
Peatland area (ha)	337,632	854,809	2,153,304	20,072,825 (ref.4)		
Peat volume* (10 ⁹ m ³)	7.8-14.9	19.7-37.6	49.5-94.7	461.7-883.4		
Carbon store* (Gt)	0.44-0.85	1.12-2.14	2.8-5.40	26.32-50.34		
				Lower estimate	Intermediate estimate	Upper estimate
Area of fire-damaged peatland (ha)	184,564 (48.1%)	474,009 (48.0%)	729,500 (29.3%)	1,450,000	2,441,000	6,804,688
Percent of peatland area damaged	54.7%	55.5%	33.9%	7.2%	12.1%	33.9%
Peat volume loss (10 ⁹ m ³)	0.85-1.03	2.18-2.66	3.36-4.09	6.67-8.12	11.23-13.67	31.30- 38.30
Peat carbon loss to atmosphere (Gt)	0.05-0.06	0.12-0.15	0.19-0.23	0.38-0.46	0.64-078	1.78-2.17
Percent peat carbon loss from store (Gt)	5.9-13.6%	5.6-13.4%	3.5-8.2%	0.8-1.8%	1.3-3.0%	3.5-8.2%
Biomass carbon loss [#] (Gt)	0.01	0.03	0.05	0.10	0.17	0.40
Total carbon loss (Gt)	0.06-0.07	0.15-0.18	0.24-0.28	0.48-0.56	0.81-0.95	2.18-2.57

In the same table is also written the extrapolated values for whole of Indonesia with total area of 190,400,000 hectares including 20,072,825 hectares peatlands. The table says that carbon loss of whole of Indonesia estimated by extrapolating the 33.9% damaged peatlands obtained for the study area that was based on fires in 1997 is 0.81-2.57 Gt (a range between the lowest value of the intermediate estimate and the highest value of the upper estimate).

Calculated value of carbon loss of the Block C of MRP is might be scientifically acceptable, but the value obtained for the whole 2,491,619 hectares of study area is unacceptable based on the following reasons. It should be bear in mind that inside the study area there are parts covering hundred thousand hectares, like area of Pulau Petak, Anjir Basarang, Pulang Pisau, and transmigration area of Bahaur, which had been cultivated for agriculture since long before the MRP. This fact suggests that it is hard to say that peat of those parts just lost by the fires in 1997. Therefore, saying 33.9% peatland of study area damaged by fires is certainly incorrect.

The carbon loss estimated for the whole area of Indonesia in Table 2 is, due to different reasons, also really questionable. There is no doubt that peatlands would be burned only if they are dried. But in natural condition the lands are always waterlogged or at least always moist and, therefore, hard to be burned. Peatlands will be dried if they are drained. Since massive drainages of peatlands resulting excessive peat drying were built only in certain areas, among other is in the MRP area with a finite extent however, then the rest of Indonesian peatlands are still safe from fires.




Figure 1. Hotspot data and estimated carbon emission in Indonesia (Hooijer *et al.*, 2006).

Although the above calculation and estimation of Page *et al.* (2001) are clearly seen has roughly been made, still the publication has widely caught interests especially of environmental experts. Carbon loss as high as 2.57 Gt shown in the publication was likely considered as real loss of carbon from Indonesian peatlands in 1997, and it is really an outstanding figure placing Indonesia in world third biggest carbon emitter after the US and China.

In addition to the above publication, Page with her colleagues published another estimation of carbon emission from Indonesian peatlands as published in Hooijer *et al.* (2006). They have collected data of hotspots on Kalimantan of the years 1997 to 2006 (Figure 1a) and have correlated the hotspots of the year 1997 with the carbon emission figures of Page *et al.* (2001) as shown in Figure 1 (b).

The upper estimate of carbon emission from Indonesian peatlands made by Page *et al.* (2001) is 2.57 Gt C in one year. By expressing the estimate in CO₂ as shown in Figure 1b, then the estimate carbon loss of 0.81-2.54 Gt of only the year 1997 is equal to CO₂ emission of 2970-9313 Mt. Using hotspot data in Figure 1a to extrapolate the estimate of the 1997 value into the time series has resulted that carbon emission from Indonesian peatlands during 1997-2007 varied between 1.42 and 4.32 Gt CO₂ each year with the average of 3 Gt CO₂/year (Figure 1b). This widely published tremendous figure has ever worsened Indonesian image as carbon emitter as again the biggest emitter in the world next to the US and China.

The above estimation was really made roughly and is scientifically very weak. Our argument to the estimation is as follows. They seem to have made a wrong analogy of carbon emission from peatland fire with those the emission in industrial countries. In industrial countries carbon emissions come from continuous industrial processes in which the emissions occur in all-time along with the processes. But in case of emission from peatlands, the emission will occur only if the lands are burned and once the peat has already burned out then there will no more emissions from peat, except smaller one from burned vegetation on the lands. The other thing is that the hotspot data presented in Figure 1b are actually an accumulation of lowlands and uplands, not necessarily of only hotspot on the peatlands. Correlating such hotspot data with carbon emission estimate based on only fire events in peatlands in Central-South Kalimantan

