

## Genetic Diversity of Antifungi-Producing Rhizobacteria of *Pseudomonas* sp. Isolated from Rhizosphere of Soybean Plant

ARI SUSILOWATI<sup>1</sup>, ARIS TRI WAHYUDI<sup>1\*</sup>, YULIN LESTARI<sup>1</sup>, SURYO WIYONO<sup>2</sup>  
AND ANTONIUS SUWANTO<sup>1</sup>

<sup>1</sup>Department of Biology, Faculty of Mathematics and Natural Sciences;

<sup>2</sup>Department of Plant Protection, Faculty of Agriculture, Institut Pertanian Bogor,  
Darmaga Campus, Bogor 16680, Indonesia

Antifungi-producing rhizobacteria have been recognized playing an important role in plant disease suppression. In our laboratory, 13 indigenous soybeans' rhizobacteria *Pseudomonas* sp. that showed strong growth inhibition of root pathogenic fungi, *Rhizoctonia solani*, *Fusarium oxysporum* and *Sclerotium rolfsii*, have been isolated from rhizosphere of soybean plant. For further understanding, the genetic diversity of the antifungi-producing *Pseudomonas* sp. was investigated using Amplified 16S rDNA Restriction Analysis (ARDRA) and 16S rRNA gene sequences analysis. 16S rDNA were amplified by PCR technique and digested with restriction endonuclease *Hae*III, *Rsa*I and *Alu*I. Sequences of 16S rRNA gene were analyzed using the BLAST program for similarity searches on sequence databases. ARDRA based dendrogram analysis was carried out by neighbor-joining of TREECON 1.3b software package. ARDRA indicated the variability of *Pseudomonas* sp. based on the digestion sites. Dendrogram clustering analysis based on the restriction enzymes profile of the amplified rDNA distinguished *Pseudomonas* sp. into 7 ribotype groups. The sequences of 16S rRNA gene confirmed that the isolates belonging to *Pseudomonas* sp. and the phylogenetic tree formed 4 clusters. There was a quite overlap among ARDRA groups and 16S rRNA sequence clusters. This finding suggested that antifungal producing *Pseudomonas* sp. were present in the rhizosphere of soybean plant and the level of genetic diversity exist within these species. Sequence analysis of the 16S rRNA gene of the *Pseudomonas* sp. with an identical ARDRA pattern confirmed that members of an ARDRA group were closely related to each other.

Key words: antifungi producing rhizobacteria, ARDRA, genetic diversity, *Pseudomonas* sp., 16S rRNA

Disease suppressive soil suppresses the growth of phytopathogenic fungi or the induction of severe disease on susceptible plants (Schroth and Hancock 1982). This phenomenon, despite rarely happens, has been extensively known and strong evidences show that the disease is suppressed by specific rhizobacteria that have the ability to produce antifungal compounds. Several experiments have demonstrated that a number of *Pseudomonas* sp. strains with the ability to produce antifungal metabolite 2,4-diacetylphloroglucinol (DAPG) can be isolated with high frequency from black root rot disease suppressive soil applied to tobacco (Keel *et al.* 1996). These strains are able to suppress diseases in wheat (Raaijmakers *et al.* 1997). They are also involved in many plant disease suppression that can be related to the presence of phytopathogen-antagonistic functions in the soil microbiota (Garbeva *et al.* 2004; Mazzola 2004; Garbeva *et al.* 2006).

There is a great interest in plant-associated bacteria, particularly in the genus of *Pseudomonas*. Researches aiming to improve crop responses emphasized on antibiotic producing bacteria indigenous to certain rhizosphere. Moreover, the components of suppressiveness have been described for multiple pathosystems, especially for those involving a specific pathogen and microbial antagonist (Weller *et al.* 2002; Borneman and Becker 2007). Recently, antibiotic-producing bacteria have been recognized playing an important role in disease suppression. In our laboratory, we have screened the indigenous soybeans' rhizobacteria *Pseudomonas* sp. CRB (Cirebon) that are potential as a biocontrol for rot root disease caused by pathogenic fungi.

Several *Pseudomonas* sp. CRB isolates strongly inhibited growth of root pathogenic fungi, i.e. *Rhizoctonia solani*, *Sclerotium rolfsii* or *Fusarium oxysporum*, *in vitro*.

Genetic diversity means the total number of genetic characteristics. Molecular tools such as ribotyping, *in-situ* hybridization, DNA sequence analysis and restriction fragment length polymorphism (RFLP) are now in common use to provide accurate genetic diversity information of microbes by using 16S rRNA gene. RFLP analysis on 16S rRNA gene is called amplified rDNA restriction analysis (ARDRA). This method is useful for genotype identification and can be used to infer genetic variability and similarity of microorganisms (Yang *et al.* 2007; Kidd *et al.* 2009). ARDRA can also be applied for characterizing a number of species including *Clostridium botulinum* toxinotype A strains (Pourshafie *et al.* 2005) and mycobacteria (De Baere *et al.* 2002). In this study, we described diversity of antifungi-producing *Pseudomonas* sp. CRB from rhizosphere of soybean plant employing ARDRA and 16S rRNA gene sequences analysis.

### MATERIALS AND METHODS

**Microorganism and Culture Condition.** *Pseudomonas* spp. were isolated from soybean's rhizosphere from a soybean plantation in Cirebon area, West Java, Indonesia. The rhizosphere's soil was diluted in 10 mL 0.85% NaCl solution and then serial dilution was made to obtain the appropriate bacterial number so that various individual bacteria would grow separately on agar surface. Diluted bacteria (100 µL) were then spread onto King's agar (King *et al.* 1954) and incubated for 24 hours at room temperature (27-28°C). Each visible colony with different appearance was picked and streaked on fresh medium to obtain pure

\*Corresponding author: Phone/Fax: +62-251-8622833,  
Email: aristri2003@yahoo.com

culture. Identification of *Pseudomonas* sp. was conducted based on morphological and physiological characters. Microgen™ system (GNA and GNB) that employs 24 standardized biochemical substrates in microwell was also used to complete the test. Gram negative, rods, motile, aerobic, catalase positive, and oxidize positive were the characters that lead to *Pseudomonas* identification (Holt et al. 1994). Eighty one isolates named *Pseudomonas* sp. CRB were collected. Thirteen isolates that showed strong inhibition of fungal pathogen *in vitro* were used in this study (Table 1). *Pseudomonas* sp. CRB were routinely cultivated on agar plate of King's medium B (20 g L<sup>-1</sup> bacto peptone, 1.5 g L<sup>-1</sup> K<sub>2</sub>HPO<sub>4</sub>, 1.5 g L<sup>-1</sup> MgSO<sub>4</sub>, 15 g L<sup>-1</sup> bacto agar (Difco, France), 15 mL L<sup>-1</sup> glycerol) in room temperature (27-28°C).

**DNA Extraction and PCR Amplification.** Genomic DNA was isolated from overnight cultures of *Pseudomonas* sp. CRB by using a cetyl trimethyl ammonium bromide (CTAB)-based protocol (Sambrook and Russell 2001). DNA coding for 16S rRNA of each isolate was amplified with primer 63f (5'-CAG GCC TAA CAC ATG CAA GTC-3') and 1387r (5'-GGG CGG WGT GTA CAA GGC-3') (Invitrogen, Japan). These primers amplify approximately 1300 bp of the 16S rRNA gene, specific to consensus regions that are considered as universal bacterial domains (Marchesi et al. 1998). PCR reactions (Takara, Japan) were done in a total reaction volume of 50 µL containing 25 µL GC buffer II, 8 µL dNTP mix (2.5 mM each), 20 pmol of each primer, 2.5 unit LA Taq polymerase, 100 ng DNA template and ddH<sub>2</sub>O. The thermal cycling (GeneAmp PCR System 2400, Perkin Elmer, USA) was as the following: denaturation at 94°C for 4 min, followed by 30 cycles of denaturation at 94°C for 2 min, annealing at 55°C for 1 min and extension at 72°C for 1 min 30 sec; with the final extension at 72°C for 5 min. PCR-amplified DNAs were visualized in 1% agarose gel electrophoresed in 1X TAE buffer at 100 V for 1 h. The gel then was stained with the ethidium bromide (5 µg mL<sup>-1</sup>) for 30 min, and visualized with UV transilluminator.

**ARDRA.** Amplified 16S rDNA was purified from the gel using Wizard SV Gel and PCR Clean-up System

Table 1 *Pseudomonas* sp. producing antifungal compounds used in this study with percentage of inhibition radial growth of phytopathogenic fungi

Isolates of <i>Pseudomonas</i>	Growth inhibition of fungi by <i>Pseudomonas</i> sp. CRB (%)		
	<i>Sclerotium rolfii</i>	<i>Fusarium oxysporum</i>	<i>Rhizoctonia solani</i>
CRB-3	-	-	56.7
CRB-16	-	24.6	-
CRB-17	-	14.3	-
CRB-31	-	18.7	50.0
CRB-44	-	39.2	-
CRB-75	-	11.1	37.7
CRB-80	20.0	-	52.3
CRB-82	-	11.1	38.9
CRB-86	-	30.3	36.9
CRB-102	25.0	-	60.0
CRB-109	-	-	36.9
CRB-111	-	-	30.0
CRB-112	-	-	48.1

(Promega, USA). The purified amplification products (1.5 µg) were single-digested with 5 unit *Hae*III, *Rsa*I and *Alu*I (Fermentas Life Science, EU). Enzyme digestions were carried out at 37°C for overnight. The restriction fragments were analyzed by electrophoresis at 2% of agarose gel in 1X TAE buffer at 100V for 50 minutes. The gel then was stained with ethidium bromide (5 µg mL<sup>-1</sup>) for 30 min, and visualized with UV transilluminator. DNA 1 kb ladder (Fermentas Life Science, EU) was used as a DNA molecular size marker. To determine series of bands that appeared in the gel as a result of successful restriction, estimation of the DNA fragment size was done using a mathematical equation that links migration rate to molecular weight. The relevant formula  $D = a - b (\log M)$  (Brown 1986); where  $D$  is the distance moved,  $M$  is the molecular weight,  $a$  and  $b$  are constants that depend on the electrophoresis condition; was applied for analysis. For each restriction band, binary data matrix was constructed on the basis of the presence or absence of each band, coded as 1 or 0, respectively. The band patterns obtained with each enzyme restriction were combined to obtain a single pattern for each isolate. The patterns were used to construct dendrogram by using the neighbor joining methods as a part of the TREECON 1.3b software package (Van de Peer and De Wachter 1997).

**16S rRNA Gene Sequence Analysis.** Purified PCR products were subjected to cycle sequencing. Single direction (forward primer) sequencing of PCR product was performed by DNA sequencer (ABI Prism 3100, Applied Biosystems, USA) at PT Charoen Pokphand, Indonesia. Databases search was carried out for similar nucleotide sequences (600 nucleotides) with the BLASTN network server at the NCBI (www.ncbi.nlm.gov/BLAST). For describe their phylogenetic relationship, the partial 16S rRNA gene sequences were aligned using ClustalW, and then analyzed by means of neighbor-joining method according to Jukes and Cantor model using MEGA version 4 (Kimura et al. 2007). The robustness of the inferred trees was evaluated by 100 bootstrap resamplings.

## RESULTS

**Amplified 16S rRNA Gene and ARDRA Pattern.** The 16S rDNAs of 13 *Pseudomonas* sp. CRB were amplified, resulting in 1300 bp bands (data not shown). Different patterns obtained from each of the three endonuclease enzymes, *Hae*III, *Rsa*I and *Alu*I, distinguished the isolates one from to another. In this study, digestion of 16S rDNA of *Pseudomonas* sp. CRB by the three enzymes produced 3 to 5 bands with different sizes for each treatment. Restriction fragment lengths for each of the possible patterns based on distance relative are presented in Table 2. Unique digestion profiles shown by each isolates analyzed could be used as references or as an ARDRA profile library. All of the 13 isolates of *Pseudomonas* sp. CRB have already been identified as biocontrol bacteria. ARDRA profile revealed in this study can be widely used to investigate or identify several *Pseudomonas* isolates that inhabit the rhizosphere and possessing the characters of biocontrol.

Table 2 16S rDNA restriction fragments of antifungi-producing *Pseudomonas* sp. digested with *Hae*III, *Rsa*I and *Alu*I

Strains of <i>Pseudomonas</i>	Size of rDNA fragment (bp)		
	<i>Hae</i> III	<i>Rsa</i> I	<i>Alu</i> I
CRB-3	600, 500, 270	540, 480, 380	630, 200, 140
CRB-16	740, 690, 210, 130	720, 440, 230	700, 200, 110, 80
CRB-17	720, 330, 210, 130	720, 440, 230	650, 430, 200, 110, 80
CRB-31	620, 500, 280, 160	590, 520, 480	670, 200, 110, 80
CRB-44	570, 210, 200, 180	520, 480, 400	380, 290, 200, 110, 80
CRB-75	740, 510, 280	590, 520, 480	470, 270, 230, 150
CRB-80	890, 210, 140	720, 440, 230	430, 200, 200, 110, 80
CRB-82	740, 220, 140	720, 440, 230	430, 200, 200, 110, 80
CRB-86	640, 530, 280	480, 440, 380	670, 200, 110, 80
CRB-102	890, 220, 180	720, 440, 230	430, 200, 200, 110, 80
CRB-109	650, 510, 190	520, 440, 410	700, 200, 110, 80
CRB-111	650, 510, 190	520, 440, 410	700, 200, 110, 80
CRB-112	600, 500, 270	480, 440, 340	590, 220, 150, 80

**Cluster Analysis Based-ARDRA.** The results of ARDRA using *Hae*III, *Rsa*I, and *Alu*I were subjected to cluster analysis. Neighbor-joining method was used to construct the dendrogram of bands resulting from

endonuclease digestion. In this method, similar variables were grouped. The ARDRA clustering revealed a considerable level of genetic diversity between the isolates, since 7 clusters, designated as ribotype 1-7, were identified (Fig 1). *Pseudomonas* sp. CRB-16, CRB-17, CRB-80, CRB-82 and CRB-102 are clustered in one group, i.e. ribotype 7. The isolates that occupy the same cluster are considered to share the same attributes. Therefore, isolates belonging to the same cluster are similar in some sense. Other isolates forming other groups differ from ribotype 7. Cluster dendrogram showed that ARDRA pattern of ribotype 1 was similar to ribotype 2 and differ from ribotype 7 (Fig 1). In this result, *Pseudomonas* sp. CRB-3 is the most unique isolate among others.

**16S rRNA Gene Sequences and Phylogenetic Relationship.** BLASTN results of the partial sequence of 16S rRNA gene (about 600 nucleotides) showed high similarity with *Pseudomonas* spp. (83-100%) (Table 3). The phylogenetic relationships among 13 isolates were represented as a dendrogram using neighbor joining method (Fig 2). Four clusters were generated when *Pseudomonas* sp. CRB isolates were grouped according to their 16S rRNA sequences. Isolates found in the same cluster seem to have close phylogenetic relationship. Many isolates with the same




Fig 1 Cluster dendrogram and electrophoregram of antifungi-producing *Pseudomonas* sp. CRB, based on restriction profiles obtained with *Hae*III, *Rsa*I and *Alu*I. Each ribotype is designated by a right brace. The dendrogram was constructed with TREECON and grouped by neighbor-joining method with bootstrap analysis of 100 resamples.



- Holt JG, Krieg NR, Sheath HA, Staley JT, Williams ST. 1994. Bergey's manual of determinative bacteriology 9<sup>th</sup> edition. Maryland: Williams and Wilkins. p 171-4.
- Keel C, Weller DM, Natsch A, Défago G, Cook RJ, Thomashow LS. 1996. Conservation of the 2,4-diacetylphloroglucinol biosynthesis locus among fluorescent *Pseudomonas* strain from diverse geographic location. Appl Environ Microbiol 62:552-63.
- Kidd TJ, Ramsay KA, Hu H, Bye PTP, Elkins MR, Grimwood K, Harbour C, Marks GB, Nissen MD, Robinson PJ, Rose BR, Sloots TP, Wainwright CE, Bell SC. 2009. Low rates of *Pseudomonas aeruginosa* misidentification in isolates from cystic fibrosis patients. J Clin Microbiol 47:1503-9.
- Kimura K, Dudley J, Nei M, Kumar S. 2007. MEGA 4: Molecular evolutionary genetics analysis (MEGA) Software version 4.0. Molecular biology and evolution 10.1093/molbev/msm092.
- King EO, Ward MK, Raney DE. 1954. Two simple media for the demonstration of pyocyanin and fluorescin. J Lab Clin Med 44:301-7.
- Marchesi JR, Sato T, Weightman AJ, Martin TA, Fry JC, Hiom SJ, Wade WG. 1998. Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA. Appl Environ Microbiol 64:795-9.
- Massana R, Murray AE, Preston CM, Delong EF. 1997. Vertical distribution and phylogenetic characterization of marine planktonic Archaea in the Santa Barbara channel. Appl Environ Microbiol 63: 50-6.
- Mazzola M. 2004. The science and art of plant-disease diagnosis. Annu Rev Phytopathol 42:35-59.
- Peer Y van de, Wachter R de. 1994. TREECON for Windows: a software package for the construction and drawing of evolutionary tree for the Microsoft Windows environment. Comput Appl Biosci 10:569-70.
- Pourshafie M, Vahdani P, Popoff M. 2005. Genotyping *Clostridium botulinum* toxinotype A isolated from patients using amplified rDNA restriction analysis. J Med Microbiol 54:933-6.
- Raaijmakers JM, Weller DM. 2001. Exploiting genotypic of 2,4-diacetylphloroglucinol-producing *Pseudomonas* spp.: characterization of superior root-colonizing *P. fluorescens* strain Q8r1-96. Appl Environ Microbiol 67:2545-54.
- Raaijmakers JM, Weller DM, Thomashow LS. 1997. Frequency of antibiotic-producing *Pseudomonas* spp. in natural environments. Appl Environ Microbiol 63: 881-7.
- Ramette A, Moënne-Loccoz Y, Défago G. 2001. Polymorphism of the polyketide synthase gene *phlD* in biocontrol fluorescent pseudomonads producing 2,4-diacetylphloroglucinol and comparison of *phlD* with plant polyketide synthases. Mol Plant-Microbe Interact 14:639-52.
- Relman DA. 1999. The search for unrecognized pathogens. Science 284:1308-10.
- Saldana G, Martinez-Alcantara V, Vinardell JM, Bellogin R, Ruiz-Sainz JE, Balatti PA. 2003. Genetic diversity of fast-growing rhizobia that nodulate soybean (*Glycine max* L. Merr). Arch Microbiol 180:45-52.
- Sambrook J, Russell DW. 2001. Molecular Cloning: a Laboratory Manual. 3<sup>rd</sup> edition. New Jersey: Cold Spring Harbor Laboratory Pr. p 662.
- Schroth MN, Hancock JG. 1982. Disease-suppressive soil and root-colonization bacteria. Science 216:1376-81.
- Tian F, Ding Y, Zhu H, Yao L, Du B. 2009. Genetic diversity of siderophore-producing bacteria of tobacco rhizosphere. Braz J Microbiol 40:276-84.
- Weller DM, Raaijmakers JM, McSpadden Gardener BB, Thomashow LS. 2002. Microbial populations responsible for specific soil suppressiveness to plant pathogens. Annu Rev Phytopathol 40:309-48.
- Woese CR. 2006. How we do, don't and should look at bacteria and bacteriologi. In: Prokaryotes (3) Volume 1. Martin Dworkin, editor. Singapore: Springer. p. 1-23.
- Xiang S, Yao T, An L, Xu B, Wang J. 2005. 16S rRNA sequences and differences in bacteria isolated from the Muztag Ata Glacier at increasing depths. Appl Environ Microbiol 71: 4619-27.
- Yang JH, Liu XH, Zhu GM, Pan YL, Xu LP, Guo JH. 2007. Diversity analysis of antagonists from rice-associated bacteria and their application in biocontrol of rice diseases. J Appl Microbiol 104:91-104.