

JURNAL

2, 9, 40

ISSN : 1978-1059

GIZI

Volume 2, Nomor 2, Juli 2007

dan PANGAN

Journal of Nutrition and Food



Publikasi resmi :

Perhimpunan Peminat Gizi dan Pangan
(PERGIZI PANGAN) Indonesia

Bekerjasama dengan

Departemen Gizi Masyarakat
Fakultas Ekologi Manusia (FEMA) IPB

FOOD CONSUMPTION AND NUTRITIONAL STATUS OF CHILDREN PARTICIPATING AT POSYANDU PROGRAM IN CIANJUR REGENCY

(Konsumsi Pangan dan Status Gizi Anak Peserta Posyandu di Kabupaten Cianjur)

Hadi Riyadi¹ and Faisal Anwar¹

ABSTRAK

Tujuan dari penelitian ini adalah untuk mengidentifikasi konsumsi pangan anak balita serta menganalisis status gizi balita. Penelitian ini menggunakan cross-sectional desain dengan sampel rumah tangga yang memiliki anak balita. Sampel berjumlah 300 balita. Pemilihan sampel dilakukan secara acak berlapis (*stratified random sampling*). Penelitian ini dilakukan di dua kecamatan Kabupaten Cianjur pada tahun 2006-2007. Data yang dikumpulkan meliputi konsumsi dan frekuensi konsumsi pangan balita serta data antropometri balita yang digunakan untuk menentukan status gizi balita. Data hasil wawancara dan pengukuran dientri menggunakan software excel. Data dianalisis dengan SAS (*Statistical Analysis System*). Hasil penelitian menunjukkan bahwa konsumsi energi balita secara rata-rata hanya memenuhi 80% angka kecukupan gizi yang dianjurkan (RDA). Konsumsi protein sudah memenuhi angka kecukupan protein yang dianjurkan. Prevalensi *underweight*, *stunted* dan *wasted* pada balita berturut-turut yaitu 30.0%, 43.7% dan 12.3%.

Keywords : *konsumsi, frekuensi konsumsi pangan, dan status gizi*

INTRODUCTION

Background

The economic crisis striking in Indonesia in 1997 is considered to have a big impact on the decreased quality of public health particularly for the poor people. From the results of evaluation on Posyandu (Health and Nutrition Integrated Service Center) carried out in November and December 1998, it has been found that the economic crisis has led to an increased number of undernourished children under five year old. The results of National Census show that the prevalence of undernourishment for under five year old children in 1989 was 37.5% (7 986 279 children under five year old) and in 2003 27.3% (5 117 409 children). The absolute value of undernourist indicated that so far the nutritional programs have not yet been adequately effective and efficient in reducing nutritional problems.

The nutritional status of a community is influenced by a lot of interrelated and complex factors. At the level of households, the nutritional status is affected by the household ability to provide adequate food in both quantity and quality, mother's pattern of upbringing children, knowledge of nutrition, and other socio-cultural factors. This indicates a close relation between food security along with its

nutritional status and public health (Azwar, 2004).

An improved status of community nutrition has a quite significant effect on the improvement of human resource quality. A nutritional status is a determinant factor of living quality and work performance. A high infant mortality rate, children under five years old and pregnant women, and a decreased work performance as well as retarded mental development and intelligence are as a result directly or indirectly of malnutrition.

Riyadi (1995) defines nutritional status as a body health condition of a person or a group of people resulting from the consumption, absorption, and utilization of nutrients in food. The physical size of a person is closely related to the nutritional status; for this reason anthropometry is accepted as a good indicator and reliable to determine a nutritional status.

Goals and Objectives

The objectives of this research were :

1. To identify food consumption of children under five years old.
2. To analyze nutritional status of children under five years old.

¹ Lecturer on Department of Community Nutrition, Faculty of Human Ecology, Bogor Agricultural University (IPB).

METHODS

Design

This research used a cross-sectional design with samples of households who have children under five. The study was conducted in the District of Cianjur, the Province of West Java and two subdistricts were selected. The research has been conducted in 2006-2007. This article is part of study of Nutrition Program Implementation: Its Utilization by Households, Coverage, Effectiveness, and Impact on Nutritional Status in Poor Areas.

Sampling

The populations in this research are children under five years old. The populations were selected in two subdistricts (Karang Tengah and Ciranjang) of Cianjur District, West Java. Sampling was done using stratified random sampling with proportional allocation where high and low frequency visit stand as stratum. Sample size was also proportional with respect to *Posyandu* size. Sample sizes of children under five years old are 186 in Karang Tengah, while 114 in Ciranjang. Total sample sizes are 300 under-five-years children.

Cianjur District was selected since there were variations in nutrition program implemented and variations in achieving the program objectives (Personal Communication with Staff of Health Services, Cianjur, February 10, 2006). The two subdistricts were chosen to increase heterogeneity of samples, so that the performance of nutrition program was truly represented by the selected samples.

Data collection

The data collected consists of :

1. Data on food consumption and food frequency.
2. Nutritional status data (age, sex, weight and height).

The data were collected through questionnaires. The data on food consumption was collected with the food frequency and 1 x 24 hour food recall. Data of nutritional status was obtained by measuring height and weighing the weight of Children.

Data Analysis and Management

Data processing involves editing questionnaires, coding, and arrangement of file structure, data entry, editing files, raising variables, combining and separating files. Data

processing and analysis used Excel and Statistical Analysis System.

The estimations of means and standard deviation made for all quantitative variables. The estimation results were presented in the forms of tables and diagrams.

RESULTS

Consumption and Nutrition Adequacy of Children

Energy Consumption. Food has been the primary need for human to live healthily (Harper, Deaton & Driskel, 1986). The more varied the food stuffs consumed, the more varied the nutrients would be obtained, resulting in the improvement of nutritional quality. Consumption and energy/nutrients adequacy are presented at Table 1 and Table 2. The average energy consumption is 1094 kcal/day as seen at Table 1.

Table 1. Statistics of Nutrients Consumption of Children under Five Years Old

Nutrients	Frequency of Posyandu Visiting		
	Low (n=180) ($\bar{x} \pm sd$)	High (n=120) ($\bar{x} \pm sd$)	Total (n=300) ($\bar{x} \pm sd$)
Energy (kcal)	1142±564	1021±576	1094±571
Protein (g)	33.7±18.6	30.7±18.6	32.5±18.6
Calcium (mg)	523±413	477±414	504±413
Phosfor (mg)	551±316	523±335	540±324
Iron (mg)	14.0±17.4	11.0±12.6	12.8±15.7
Vitamin A (µgRE)	689±920	612±643	378±48
Vitamin C (mg)	37.6±33.6	33.1±27.8	35.8±31.5
Vitamin B1 (mg)	0.34±0.19	0.32±0.18	0.33±0.18

On the average, this energy consumption only fulfills 80% of the nutrition adequacy recommended by RDA. It can be seen from the two participation groups (Table 1) that the energy consumption of the children of low participation group (1142 kcal/day) is a little higher than that of high participation group (1021 kcal/day).

According to Hardinsyah and Martianto (1998), nutrition deficiency especially in energy and protein at the initial stage only causes the feeling of hunger; however, in a certain period, this will cause the reduction (loss) of body weight. With this low level of energy consumption, children will only obtain

two-third of the recommended energy adequacy level (RDA) (Table 2).

Tabel 2. Statistics of Nutritional Adequacy of Children Under Five Years Old

Nutrients	Frequency of Posyandu Visiting		
	Low (n=180) ($\bar{x} \pm sd$)	High (n=120) ($\bar{x} \pm sd$)	Total (n=300) ($\bar{x} \pm sd$)
Energy (%)	82.9±40.3	76.9±44.9	80.5±44.2
Protein (%)	132.7±70.9	125.6±79.9	129.9±74.6
Calcium (%)	105.1±82.9	96.7±86.2	101.8±84.2
Phosfor (%)	198.2±109.0	192.7±122.1	196.0±114.2
Iron (%)	170.5±209.3	137.8±158.7	157.6±191.2
Vitamin A (%)	181.9±235.3	164.6±167.2	175.0±210.8
Vitamin C (%)	90.7±80.5	81.0±68.7	86.9±76.1
Vitamin B1 (%)	59.9±32.3	58.2±34.9	59.2±33.3

Protein Consumption. Table 1 shows that the average protein consumption among children is 32.5 g/day. As a whole, the protein consumption is much higher than the recommended protein adequacy level (129. 0%). On the other hand, if seen from each participation group (Table 1), it can be said that the average protein consumption of low participation group is 33.7 mg which has no significant difference from that of high participation group (30.7 mg). The protein adequacy level of the children is greater than the recommended level (Table 2). This protein usually derives from salty fish.

Calcium Consumption. The calcium consumption by children is generally high, with the average consumption of 504 mg per day. The average protein consumption in low participation group is 523 mg higher than that of the children in high participation group (477 mg) (Table 1). The consumption level indicates that their need has been satisfied (Table 2). The calcium source greatly consumed by them is from anchovies.

Phosphor Consumption. The average consumption of phosphor is 540 mg/day (Table 1) in which the level has doubled the recommended phosphor adequacy level (Table 2). The consumption of both groups is not significantly different.

Iron Consumption. The tables show that the average iron consumption is approximately 12.8 mg/day. Iron consumption for the low group is rather higher (14.0 mg/day) than that of the high group (11.0 mg/day). The adequacy level of iron has reached 150%. The iron

consumption level in the low group is 170% whereas that of high group reaches 138% (Table 2).

Vitamin A Consumption. The average consumption of vitamin A is 658 ugRE or reaches 175% of the recommended level. The consumption of vitamin A for the low group is 689 ugRE and has reached 182% of the recommended level whereas that of the high group reaches 165 ugRE and has reached 165% of the recommended level.

Vitamin C Consumption. The average consumption of vitamin C is 35.8 mg or only reaches 87% of the recommendation. There is no big difference in vitamin C consumption between low and high groups. This indicates that food quality is varied among the children, which may not be related to their participation at posyandu.

Vitamin B1 Consumption. The average consumption of vitamin B1 is 0.33 mg or meets 59% of the recommended adequacy. The vitamin B1 consumption has no difference in both participation groups.

Food Frequency of Children

The Frequency of Staple Foods. Cereal food types (rice, corn, and wheat) contain high carbohydrates and sufficient protein content which will be utilized as energy sources (Winarno, 1995). The food frequency among children is presented to see their eating habits/patterns in the research area. From Table 3, the cereals consumed by the children include rice, corn, cassava and sweet potato, and more than 80 % of them consume these foods.

No difference in the frequency of cereal food consumption between the two groups. These cereal foods are mostly purchased, and they are rarely planted or given by other people. Rice is commonly purchased - this is claimed by 90% of the respondents.

Frequency of Animal Protein Consumption. Animal products contain the highest biological value of animal protein. Based on its characteristics, animal protein contains relatively high mineral and protein contents including various vitamins especially the unique vitamin B12 (Winarno, 1995). Types of animal protein sources include meat products, fish and eggs which are consumed everyday by more than 95% of the children. The foods consumed 4 times per week include salty fish and eggs (Table 4). Chicken meat and preserved

fish (*pindang*) are generally consumed once a week.

There is no difference in the frequency of animal protein consumption between the two groups. Most of these products are purchased. Only a little portion of the products are obtained from the animals raised by the respondents. There are 10% respondents stating that fresh water fish products are obtained from their own ponds or they are given by their neighbors.

Frequency of Legumes Consumption. Soybeans are the sources of vegetable proteins. They are not commonly consumed everyday by the children (Table 5). Tofu and fermented soybean (*tempeh*) are consumed by more than 98% of the children with the fre-

quency of 3 times per week. This might happen because these products are easily obtained and cheap. The research conducted by the Research Development Center of Nutrition in Bogor proves that the consumption of *tempeh* as one of the supplementary major food may increase body weight and improve the nutritional status of the children (Winarno, 1995). Other types of nuts are rarely consumed.

There is no difference in the frequency of soybean consumption between the two groups. Food of vegetable protein sources are almost 100.0 % bought by respondents. Only few respondents those stated they obtained it from others people or self-cultivated.

Table 3. Statistics of Staple Food Consumption Frequency of Children under Five Years Old

Kinds of Staple Food	Participation at Posyandu	Children under Five Years Old		Frequency of Consumption (times per)		
		n	%	Month	Week	Day
Rice (<i>Oryza sativa</i>)	Low	180	100.0	76.8	17.7	2.5
	High	120	100.0	74.3	17.2	2.5
Corn (<i>Zea mays</i>)	Low	152	83.9	2.1	0.5	0.1
	High	99	83.2	2.5	0.6	0.1
Cassava (<i>Manihot utilissima</i>)	Low	150	82.8	1.2	0.3	0.04
	High	100	84.0	1.4	0.3	0.1
Sweet potato (<i>Ipomoea batatas</i>)	Low	145	80.1	1.5	0.3	0.1
	High	95	79.8	1.8	0.4	0.1
Taro (<i>Colocasia esculenta</i>)	Low	72	39.8	0.3	0.1	0.01
	High	39	32.8	0.3	0.1	0.01
Potato (<i>Solanum tuberosum</i>)	Low	14	7.7	0.2	0.1	0.01
	High	20	16.8	1.1	0.3	0.04

Table 4. Statistics of Animal Protein Consumption Frequency of Children under Five Years Old

Kinds of Animal Protein	Participation at Posyandu	Children under Five Years Old		Frequency of Consumption (times per)		
		n	%	Month	Week	Day
Salty fish	Low	177	97.8	18.5	4.3	0.6
	High	118	99.2	18.6	4.3	0.6
Layer chicken's egg	Low	179	98.9	18.4	4.2	0.6
	High	115	96.6	18.8	4.3	0.6
Preserved fish	Low	173	95.6	8.1	1.9	0.3
	High	115	96.6	6.8	1.6	0.2
Chicken meat	Low	180	99.5	5.4	1.3	0.2
	High	118	99.2	4.6	1.1	0.2
Fresh water fish	Low	157	86.7	3.7	0.8	0.1
	High	99	83.2	3.1	0.7	0.1
Sea water fish	Low	75	41.4	1.4	0.3	0.04
	High	41	34.5	0.8	0.2	0.03
Chicken nugget	Low	58	32.0	1.3	0.3	0.04
	High	26	21.9	0.8	0.2	0.03
Kampong chicken's egg	Low	52	28.7	1.0	0.2	0.03
	High	40	33.6	1.2	0.3	0.04
Beef/goat meat	Low	169	93.4	0.9	0.2	0.03
	High	113	95.0	0.7	0.2	0.02

Table 5. Statistics of Legumes Consumption Frequency of Children under Five Years Old

Kinds of Nuts	Participation at Posyandu	Children under Five Years Old		Frequency of Consumption (times per)		
		n	%	Month	Week	Day
Tofu	Low	180	100.0	16.4	3.8	0.5
	High	120	100.0	15.1	3.5	0.5
Tempeh	Low	177	97.8	14.3	3.3	0.5
	High	119	100.0	14.3	3.3	0.5
Mung-bean (<i>Phaseolus vulgaris</i>)	Low	157	86.7	4.7	1.1	0.2
	High	108	90.8	3.5	0.8	0.1
Oncom (Traditional Indonesian Fermented Food)	Low	154	85.1	3.8	0.9	0.1
	High	98	82.4	3.6	0.8	0.1
Peanut (<i>Arachis hypogaea</i>)	Low	155	85.6	3.9	0.9	0.1
	High	98	82.4	3.0	0.7	0.1
Cowpea (<i>Vigna unguiculata L.</i>)	Low	6	3.3	0.1	0.02	0.0
	High	4	3.4	0.1	0.03	0.0

Frequency of Leaf Vegetables. Vegetables are sources of vitamin and minerals, those very useful for children under five. Generally, green vegetable is rich of important nutrients and its present on meals will improve the nutritional quality entirely.

Children under five rarely consumed leaf vegetables (Table 6). Frequency of leaf vegetables consumption only 1 time per week for mostly respondents. Kinds of those consumed leaf vegetables once a week are spinach, sawi (*Brassica rugosa*), carrot, long-bean (*Cajanus cajan*), and cabbage. There is no difference in the frequency of leaf vegetables consumption between the two groups.

Such vegetables as spinach, kangkung (*Ipomoea reptans*), carrot, sawi, cabbage, long-bean and melinjo leaves are commonly purchased by the respondents. Cassava and papaya leaves can be obtained by purchasing

them or taking them from the respondents' garden or they obtained the leaves from their neighbors.

Frequency of Fruit Vegetable. Like leaf vegetables, fruit vegetables are also rarely consumed by the children (Table 7). From these types of vegetables, tomato are the most frequently consumed by the children i.e. 6 times per week or almost everyday. Fruit vegetables those consumed once a week are cucumber and pumpkin. There is no difference in the frequency of the consumption between the two groups.

Frequency of Fruit. Fruits are also vitamin and mineral sources. The largest sources of vitamin C derive from fruits especially the fresh ones. In general, these fruits are not consumed daily by the children (Table 8). Oranges and banana are often consumed by the children i.e. 3 times a week since they are

Table 6. Statistics of Leaf Vegetables Consumption Frequency of Children under Five Years Old

Kinds of Leaf Vegetables	Participation at Posyandu	Children under Five Years Old		Frequency of Consumption (times per)		
		n	%	Month	Week	Day
Spinach	Low	172	95.0	8.1	1.9	0.3
	High	115	96.6	8.7	2.0	0.3
Kangkung (<i>Ipomoea reptans</i>)	Low	177	97.8	7.7	1.8	0.3
	High	115	96.6	9.0	2.1	0.3
Carrot	Low	178	98.3	7.3	1.7	0.2
	High	116	97.5	7.8	1.8	0.3
Cabbage	Low	168	92.8	6.5	1.5	0.2
	High	112	94.1	6.5	1.5	0.2
Sawi (<i>Brassica rugosa</i>)	Low	165	91.2	5.9	1.3	0.2
	High	109	91.6	5.8	1.3	0.2
Long-bean (<i>Cajanus cajan</i>)	Low	174	96.1	4.7	1.1	0.2
	High	115	96.6	4.6	1.1	0.2
Melinjo leaves (<i>Gnetum gnemon</i>)	Low	147	81.2	3.8	0.9	0.1
	High	97	81.5	3.7	0.9	0.1
Cassava leaves	Low	154	85.1	3.1	0.7	0.1
	High	100	84.0	3.7	0.9	0.1
Papaya leaves	Low	107	59.1	1.7	0.4	0.1
	High	71	59.7	2.5	0.6	0.1

Table 7. Statistics of Fruit Vegetables Consumption Frequency of Children under Five Years Old

Kinds of Fruit Vegetables	Participation at Posyandu	Children under Five Years Old		Frequency of Consumption (times per)		
		n	%	Month	Week	Day
Tomato	Low	180	100.0	26.8	6.2	0.9
	High	118	99.2	28.1	6.5	0.9
Cucumber	Low	175	96.7	8.2	1.9	0.3
	High	112	94.1	6.9	1.6	0.2
Pumpkin (<i>Legenaria leucantha</i>)	Low	176	97.2	5.4	1.3	0.2
	High	114	95.8	4.8	1.1	0.2
Eggplant	Low	144	79.6	3.6	0.8	0.1
	High	101	84.8	3.5	0.8	0.1
Baby jackfruit	Low	115	63.5	2.1	0.5	0.1
	High	70	58.8	1.3	0.3	0.04
Baby papaya	Low	62	34.3	1.0	0.2	0.03
	High	41	34.5	0.7	0.2	0.02

Table 8. Statistics of Fruits Consumption Frequency of Children under Five Years Old

Kinds of Fruits	Participation at Posyandu	Children under Five Years Old		Frequency of Consumption (times per)		
		n	%	Month	Week	Day
Guava	Low	158	87.3	3.0	0.7	0.1
	High	101	84.8	3.1	0.7	0.1
Papaya	Low	164	90.6	5.1	1.2	0.2
	High	110	92.4	3.5	0.8	0.1
Mango	Low	168	92.8	4.3	1.0	0.1
	High	104	87.4	3.6	0.8	0.1
Pineapple	Low	85	47.0	1.6	0.4	0.05
	High	45	37.8	0.5	0.1	0.02
Banana	Low	176	97.2	11.8	2.7	0.4
	High	116	97.5	12.3	2.8	0.4
Jackfruit	Low	140	77.4	1.1	0.3	0.04
	High	78	65.6	1.0	0.2	0.03
Hairyfruit	Low	156	86.2	0.1	0.03	0.0
	High	107	89.9	0.1	0.03	0.0
Orange	Low	179	98.9	13.8	3.2	0.5
	High	117	98.3	13.3	3.1	0.4
Apple	Low	141	77.9	3.0	0.7	0.1
	High	89	74.8	2.6	0.6	0.1
Melon	Low	1	0.6	0.01	0.0	0.0

easy to find and relatively cheap whereas papaya and mango are only consumed once a week. There is no difference in fruit consumption between the low and high participation groups.

Frequency of Snacks. Snacks containing carbohydrates are not consumed every day by the children (Table 9). Research by Widiyawati (2004) states that snacks frequently consumed by the children include chiki potato chips, candies, chocolate, and nuts. There is no difference in snacks consumption between the low and high participation groups. They are generally purchased from the small shops or traveling sellers.

Frequency of Miscellaneous Food. These miscellaneous foods such as sugar and monosodium glutamate (MSG) are almost consumed daily by the children (see Table 10).

This is because their mothers always put these foods into their meals to make them tastier. There is no difference in miscellaneous foods consumption between the low and high participation groups. Like the other foods, they are generally purchased by the children.

Nutritional Status of Children

Anthropometric Characteristics of Children

The average age of children in the research location is 2.7±1.0 years, with lowest age of 1 year and the highest age of 5 years. The average age is relatively the same in both low and high participation groups. Similarly, there is no difference in the average age for different sexes in both participation groups at posyandu.

Table 9. Statistics of Snacks Consumption Frequency of Children under Five Years Old

Kinds of Snacks	Participation at Posyandu	Children under Five Years Old		Frequency of Consumption (times per)		
		n	%	Month	Week	Day
Fried snacks	Low	179	98.9	20.5	4.7	0.7
	High	116	97.5	19.0	4.4	0.6
Fried banana	Low	167	92.3	14.3	3.3	0.5
	High	111	93.3	13.1	3.0	0.4
Meat ball	Low	174	96.1	8.9	2.1	0.3
	High	114	95.8	8.4	1.9	0.3
Bakwan (fried vegetables)	Low	116	64.1	7.5	1.7	0.3
	High	80	67.2	6.8	1.6	0.2
Siomay (kind of Chinese foods)	Low	110	60.8	3.8	0.9	0.1
	High	68	57.1	3.0	0.7	0.1
Chicken noodle	Low	111	61.3	2.2	0.5	0.1
	High	78	65.6	2.1	0.5	0.1

Table 10. Statistics of Miscellaneous Food Consumption Frequency of Children under Five Years Old

Kinds of Miscellaneous Food	Participation at Posyandu	Children under Five Years Old		Frequency of Consumption (times per)		
		n	%	Month	Week	Day
Sugar	Low	180	100.0	27.6	6.4	0.9
	High	117	98.3	26.4	6.1	0.9
Tea	Low	169	93.4	22.7	5.2	0.8
	High	99	83.2	21.0	4.8	0.7
Coffee	Low	116	64.1	13.9	3.2	0.5
	High	76	63.9	12.5	2.9	0.4
Sauce	Low	139	76.8	8.1	1.9	0.3
	High	79	66.4	6.6	1.5	0.2
Kerupuk (fried crackers)	Low	178	98.3	18.7	4.3	0.6
	High	116	97.5	18.1	4.2	0.6
Vetsin/MSG	Low	176	97.2	29.2	6.7	1.0
	High	118	99.2	30.2	6.9	1.0
Soybean ketchup	Low	172	95.0	15.3	3.5	0.5
	High	114	95.8	14.3	3.3	0.5

On the whole, the average body weight of children is 11.3±2.7 kg, with the lowest body weight of 6.3 kg and the highest of 26.1 kg. The body weights in both participation groups are not much different, but the body weights of boys tend to be greater than those of girls in both participation groups.

The average body height of all children is 85.4±10.0 cm, with the lowest body height of 59.5 cm and the highest of 112.1 cm. The body heights of children are relatively not different between the two participation groups. Boys have greater body heights than girls in both participation groups. In the low group, the body heights of boys are about 2 cm higher than the girls. In the high group, the body heights of boys are about 4 cm higher than the girls.

Weight for Age of Children (W/A)

The nutritional status discussed in this report is based on the resulted measurement of body weights and heights, so the analysis

uses the weight for age (W/A), height for age (H/A), and weight for height (W/H). The data processing uses WHO software, Antro 2005 and the new WHO anthropometric standard.

Based on the calculation of Z-score of W/A and the threshold point < - 2SD for the underweight, the prevalence of underweight among children (less than five years old) is 30.0%. The underweight prevalence is much higher at the research location than that in Indonesia (23.1 %) obtained by SUSENAS 2005 (National Survey of Social Economic). The underweight prevalence among children in the low participation group (34.3%) is much higher than that among children in the high participation group (23.5%).

There is no consistent pattern of underweight prevalence between sexes. In low participation group, the underweight prevalence is greater for boys, whereas in the high participation group it is greater for girls. The prevalence of severe underweight (Z-score < - 3 SD) in the research location is 8.3 %, the same

as the result obtained by SUSENAS in Indonesia.

The prevalence of overweight children is very low, i.e. 0.7 %. Overweight is more prevalent in the high participation group than in the low group. From Figure 1 it is clearly seen that the Z-score median of W/A for the children in the research site is much further to the left than the WHO standard of 2005. The median Z-score for the children in the research location is coming closer to the standard deviation of -1.5 or underweight.

The Z-score (-1.52) for the children of the low participation group is worse than that (-1.13) of the high participation group. Like the underweight prevalence, the Z-scores between different sexes do not have a consistent pattern. In the low participation group, the Z-score of W/A is higher for boys; whereas in the high participation group it is higher for girls. The gender-based analysis shows that the curve of Z-scores is not much different between boys and girls (see Figure 2). From Figure 5.3 it can be seen that in terms of age groups, the Z-score of W/A is the worst in its deviation for the age group of 24-47 months.




Figure 1. Distribution of Children based on the Z-score of W/A




Figure 2. Distribution of Children based on the Z-score of W/A and Sex




Figure 3. Z-score of W/A Value of Children based on Age Group

Height for Age of Children (H/A)

A body height at a certain time is the resulted cumulative growth since birth, therefore it can be used as a previous picture of nutritional status in the past. Body height is also the most sensitive index for detecting the socio-economic changes. The prevalence of stunted growth as a whole is 43.7%. This prevalence is closer to the prevalence in Indonesia 5-6 years ago.

The comparison of the prevalence of stunted growth between participation groups shows that the prevalence is greater for children in the low participation (46.4%) than in the high participation group (39.5%). The stunted prevalence between different genders does not take a consistent pattern. In the low participation group, the stunted prevalence is

higher among boys; whereas in the high participation group, it is greater among girls. The *severe stunted* prevalence (based the criteria of < -3 SD) in this research is 20.3%. Eating habit with a lack of animal protein and inadequate daily intake of energy can be the cause of *severe stunted* growth among children.

It can be seen from Figure 4 that the median Z-score of H/A for the children in the research location is farther to the left than the standard of WHO, i.e. getting closer to -2 standard deviation or *stunting*.

The gender-based analysis shows that the curve for the Z-score of H/A is slightly worse in the girls than in the boys (see Figure 4). The Z-score of H/A (-1.9) for children in the low participation group is worse than that of the high participation groups (-1.6).




Figure 4. Distribution of Children based on the Z-score of H/A




Figure 5. Distribution of Children according to the Z-score of H/A and Sex




Figure 6. Z-score of H/A Value of Children based on Age Group

Like the underweight prevalence, Z-score of H/A between different genders does not take a consistent pattern. In the low participation group, the boys have a higher Z-score of H/A, whereas in the high group the girls have a higher Z-score of H/A. Like in Figure 3, in Figure 6 in terms of age groups it can be seen that the Z-score of H/A is the worst in the age group of 24-35 months.

Weight for Height of Children (W/H)

Based on the calculation of Z-score of W/H with the *wasted* criteria of < -2 SD, the wasted prevalence is found to be 12.3 %. The prevalence of wasted children in the low participation growth (14.9%) is much higher than that of the high participation group (8.4%). The prevalence among the boys is higher than among the girls in both participation groups.

The prevalence of *severe wasted* (Z-score of < -3 SD) in this research is very low, i.e. 4.7 %. From Figure 7 it is clear that the median Z-score of W/H for the children in the research location is rather away to the left compared to the WHO standard of 2005. The median Z-score of W/H for the children in the area is approaching the standard deviation of -0.5 or wasting.

The gender-based analysis shows that the curve for the Z-score of W/H among the boys is relatively the same as that among the girls (see Figure 8). The prevalence of wasted among the boys (15.4 %) is higher than that among the girls (9.3 %). Like in Figure 3, in Figure 9 in terms of the age groups, it can be seen that the Z-score of W/H is the worst in the age group of 36-47 months.




Figure 7. Distribution of Children according to the Z-score of W/H




Figure 8. Distribution of Children based on the Z-score of W/H and Sex



Figure 9. Z-score of W/H Value of Children based on Age Group

CONCLUSION AND RECOMENDATION

Conclusion

1. The average consumption of energy for children under five years old is still below the recommended dietary allowance (80 % of RDA), whereas the protein consumption is already above the RDA.
2. The prevalence of underweight, stunted, and wasted among children are respectively 30.0 %, 43.7 %, and 12.3 %
3. The activities at posyandus have a positive impact on the nutritional status of children under five year's olds. The more frequent the visit to posyandus, the better the three nutritional statuses would be

Recommendation

The consumption of children under five years old have impact on the nutritional status of them, so that mothers having children under five years old should improve the consumption of children. And then the nutritional knowledge of mothers is necessary to make effort in improving the consumption of children.

ACKNOWLEDGEMENT

The researchers would like to extend its gratitude to the Neys-van Hoogstraten Foundation (NHF), the Netherlands for funding this research project. Thank you and appreciation to Ali Khomsan, Dadang Sukandar, and Eddy S. Mudjajanto from Department of Community Nutrition, Faculty of Human Ecology, Bogor Agricultural University as other researchers who also involved in this project.

BIBLIOGRAPHY

- Azwar A. 2004. Aspects of Health and Nutrition in Food Security. Paper of National Workshop on Food and Nutrition 2004. LIPI (Indonesian Institute of Science). Jakarta.
- Hardinsyah & D Martianto. 1988. Evaluation of Energy and Protein Adequacy Level and Nutritional Quality of Food Consumption. Wirasari, Jakarta.
- Harper IJ, BJ Deaton, & JA Driskel. 1986. Food, Nutrition, and Agriculture (Suhardjo, translator) (2nd Ed.). UI Press, Jakarta.
- Riyadi H. 1995. Evaluation and Measurement of Nutritional Status. Lecture Notes of Departement of Community Nutrition and Family Resources, Faculty of Agriculture, IPB, Bogor.
- Widiyawati R. 2004. Relation of Mother's Nutritional Knowledge and behaviour with Children under Five's Parenting Style in East Bogor Sub-district, Bogor City. Undergraduate script (unpublished). Department of Community Nutrition and Family Resources, Faculty of Agriculture, Bogor Agricultural University (IPB), Bogor.
- Winarno FG. 1997. Food Chemistry and Nutrition. Gramedia Pustaka Utama Ltd., Jakarta.
- _____. 1995. Nutrition and Food for baby and Post Breastfed baby. Pustaka Sinar Harapan, Jakarta.