
ian atau seluruh karya tulis ini tanpa mencantumkan dan menyebutkan sumber:

utip sebag



(Institut Pertanian Bogor)



EEDINGS OF THE INTERNATIONAL WORKSHOP

Tropical Bio-resources for Sustainable Development

"The cole of Innovation to Enhance German Alumni in Scientific and Professional Capacities"

Editors:

Syarifah Iis Aisyah Nandi Kosmaryandi Anuraga Jayanegara Ronald F. Kuehne

IPB International Convention Center August 13rd-15th, 2014 Bogor, Indonesia Dilarang mengutip sebagian atau seluruh karya tulis ini tanpa mencantumkan dan menyebutkan sumber Hak Cipta Dilindungi Undang-Undang

PROCEEDINGS OF THE INTERNATIONAL WORKSHOP TROPICAL BIO-RESOURCES FOR SUSTAINABLE DEVELOPMENT

The Role of Innovation to Enhance German Alumni in milik IPB (Institut Pertanian Bogor) Scientific and Professional Capacities"

13-15 August 2014 Bogor, Indonesia

Editors

Syarifah Iis Aisyah (Bogor Agricultural University, Indonesia) Nandi Kosmaryandi (Bogor Agricultural University, Indonesia) Anuraga Jayanegara (Bogor Agricultural University, Indonesia) Ronald F. Kuehne (Georg-August-Universitaet Goettingen, Germany)

Bogor Agricultural Universi







Dilarang mengutip sebagian atau seluruh karya tulis ini tanpa mencantumkan dan menyebutkan sumber

PREFACE

it is really honoured and very pleased to have this 6thSEAG International workshop, which is organized by SEAG (South East Asia-Germany) Alumni Network-Indonesia,in collaboration with CDA (Career Development and Alumni Affairs), Bogor Agricultural University. SEAG is the German Alumni-networking group, which was established in year 2000, among countries in South-East Asia.

Since 1999, the Federal Ministry for Economic Cooperation and Development (BMZ) and the German Academic Exchange Service (DAAD) have been systematically supporting alumni networks of graduates from German Universities. The University of Goettingen, Kassel and Marburg established an alumni consortium to support and maintain efficiently local and regional alumni networks in Egypt-Arab-Region (GEAR), in Latin America (ReCALL), in Iran (GIAN) and in South East Asia (SEAG).

The objectives of the alumni networks are to establish an alumni database to enable the exchange of scientific experiences among the alumni and their host universities in Germany, and finally to create and maintain local and region network. In order to achieve these goals, the consortium uses many tools, e.g. organizing symposium, mini workshop, international workshop, summerschool, etc.

For regional Indonesia, some Mini Workshops had been done several times which were taken placed in many universities in difference provinces. The first SEAG mini workshop had been done in Brawijaya University, Malang, on April 2003 for those alumni who work in Agriculture economy. The second one was executed in Soedirman University, Central Java on May 2004, for Agriculturist, and the third SEAG mini workshop was conducted in Taman Safari Bogor, May 2005 for Animal scientist. The fourth was in Sam Rajulangi University – North Sulawesi, for the society of forester, with the theme of Developing Public Awareness through Sustainable Forest Management. The fifth was conducted in USU (North Sumatera University) for area of Agricultural Technology, in November 2006. The Sixth was in IPB Bogor for horticulturist, on May 2007.

As academicians or researchers who gained education, training or part of it in Germany, we should play a role as key person in our scientific society. Our partners from Germanyalso believe that their support can only be



Dilarang mengutip sebagian atau seluruh karya tulis ini tanpa mencantumkan dan menyebutkan sumber Hak Cipta Dilindungi Undang-Undang

effectively provided if it is based on cooperation with key local players. Therefore development cooperation is very essentially dependent on identifying and integrating such key persons. Indeed, as German alumni, we have to show an effort to support for economic, technological and social transformation processes in our countries.

Some of us hold important positions in government, in the administration, in business and industry and in academia. We may act as multipliers and disseminators in and within our societies. We should also introduce the requisite specialist knowledge, provide motivation for innovation and guarantee cooperative capabilities in dealings with local and foreign partners. That is why we explore the theme of Tropical Bio-resources for Sustainable Development: The Role of Innovation to Enhance German Alumni in Scientific and Professional Capacities, for this workshop.

High appreciation is conveyed to the organizing committee from SEAG-Indonesia and CDA IPB for the effort to conduct this workshop. The very sincered thank is delivered to the German Academic Exchange Service (DAAD) for continues support financially and many other aspect give us invaluable opportunities to learn from each other, to improve individual and institution competences, and to experience a lot of things across universities.

Syarifah Iis Aisyah

SEAG INDONESIA CDA IPB

Bogor Agricultural University

1. Dilarang mengutip sebagian atau seluruh karya tulis ini tanpa mencantumkan dan menyebutkan sumber: Hak Cipta Dilindungi Undang-Undang

TABLE OF CONTENTS

INT	RODUCTION1
WO	RKING GROUP 1: FUNDAMENTAL ASSESSMENT FOR
BIO	RESOURCES
1. I	Distribution pattern of Alstonia scholaris - a species containing
	medicinal substances - in different forest ecosystems
(Bambang Irawan, Iskandar Z. Siregar, Reiner Finkeldey)4
2. I	Radical scavenging activity of leafy amaranths as potential
8	an Foxidant sources (Muhammad Ikhsan Sulaiman, Rita Andini) 11
	Yield evaluation of 17 chili pepper (<i>Capsicum annuum</i> L.) lines
	n Bogor, West Java (Faradila Danasworo Putri, Muhamad
	Syakur, Syarifah Iis Aisyah)17
	The research development of in vitro embryo production on
	farm animal in Indonesia (Mohamad Agus Setiadi)23
	Carrageenan prototype food product development of seaweed at
	Slester Salabangka Islands of Central Sulawesi Province
	Marhawati Mappatoba, Asriani Hasanuddin)28
	number of Dipterocarpaceae at Soraya Research Station,
	Leuser Ecosystem Area (<i>Iqbar, Essy Harnelly</i>)
	influence of different supplemental niacin levels on intake, ligestibility and rumen fermentation of dairy cows: a meta-
	analysis (Rossy E. A. Anggreini, Erika B. Laconi, Anuraga
	[ayanegara]
	nfluence of tannin concentration in ration on fermentation
	parameters of Rumen Simulation Technique (RUSITEC): a
	neta-analysis (Anuraga Jayanegara, M. Ridla, Erika B. Laconi,
	<i>Vahrowi</i>)
	bservation on the scales of three species of <i>Varanus</i> using
	aning electron microscopy (Evy Arida)
	The potency of tropical endophytic bacteria as plant growth and
	iocontrol agents (Abdul Muñif)54
11. E	Effect of oxygen concentration on storage of sapodilla fruit
(1	Achras zapota 1.) (Bambang Susilo, Rini Yulianingsih, Dyah Ayu
A	gustiningrum)61
	Heavy metals and other elements concentration in Emilia
	onchifolia grown in topand overburden of Serpentine soil from
S	orowako, Indonesia (A. Tjoa, H. Barus)67
	Jn

. Dilarang mengutip sebagian atau seluruh karya tulis ini tanpa mencantumkan dan menyebutkan sumber:

Hak Cipta Dilindungi Undang-Undang



WORKING GROUP 2: APPLIED RESEARCH AND SCALING-UP OF BIO-RESOURCE INNOVATION

UI	OF BIO-RESOURCE INNOVATION
1.	Thermal hydrolyzed rice husk as bioherbicide to control sedge rice weed <i>Fimbristylis miliacea</i> (L.) Vahl (<i>H. Agusta, M. Syakir,</i>
	D. Guntoro, M.B. Yunindanova, B. Arifin)76
2.	Potential sustainable maize-peanut production using appropriate
	biofertilizer technology in ultisols of Moramo district, South
	Konawe regency (L. Karimuna, A. Maruf, Rahman, L.A. Sani) 81
3.	Evaluation of estrus synchronization with sponge vaginal and
	artificial insemination technologies on sheep and goat (case
	study at Juhut village, Karang Tanjung sub district, Pandeglang
	district, Banten) (Siti Darodjah Rasad, Rangga Setiawan, Toha, Kikir Winangun)92
4.	Concentrate protein albumin (probumin) from snakehead fish
	(Channa striata); local product of food supplement as cheap
	alberninprotein source for community (Abu Bakar Tawali, Meta
	Mattendradatta, Veni Hadju)96
5.	Howsehold scale environmentally friendly measures to reduce
	resource consumption (Arief Sabdo Yuwono)
5.	Use of white rot fungi- and bacterial rot in decomposition of
	cocoa pod waste and in growth reduction of <i>Phytopthora</i>
	palaivora and Lasiodiplodia theobromae (Tutik Kuswinanti, Ade
7	Rosmana, Vien Sartika Dewi, Baharuddin, Jamila)
7.	Repellence test of spices (garlic, chili, and pepper) to rat (<i>Rattus rattus diardii</i> 1.) (<i>Swastiko Priyambodo, Dewi Safitri</i>)
3.	Growth response of dragon fruit (Hylocereus costaricensis) on
).	MS medium with Gandasil and Growmore in vitro (Faridatul
	Mukminah, Busroni Asnawi, Tetra Tri Novi)
	Effect of enriched phospho-compost application on the growth
	and phosphorous content of Setaria splendida Stapf (R. Dianita,
	A. Rahman Sy, Ubaidillah)
	Implementation of life cycle assessment (LCA) on tempeh
	production in Bantul district - Yogyakarta special province -
	Indonesia (Wahyu Supartono, Lina Widyasari, Didik Purwadi) 133
	Study of characteristics floral and morphological hybrid rice
	parental lines on different seeding date (P.N. Susilawati, M.
	Surahman, B.S. Purwoko, T.K. Suharsi, Satoto)
	Natural grass and plant residue qualities and values to support
	lactating cows requirement on forage at Indonesian small scale enterprise and traditional dairy farming (<i>Despal</i> , <i>Jazmi Malyadi</i> ,
	Yessy Destianingsih, Ayu Lestari, Hari Hartono, Luki Abdullah) 145
	Desiratingsin, Tya Desirit, Hart Hartono, Bakt Hodattan) 1

Hak Cipta Dilindungi Undang-Undang

1. Dilarang mengutip sebagian atau seluruh karya tulis ini tanpa mencantumkan dan menyebutkan sumber: a. Pengutipan hanya untuk kepentingan pendidikan, penelitian, penulisan karya ilmiah, penyusunan laporan, penulisan kritik atau tinjauan suatu masalah. b. Pengutipan tidak merugikan kepentingan yang wajar IPB.

13	Effect of seed density and nutrient source on production and quality of green house fodder as dairy cattle feed (<i>Idat Galih Permana</i> , <i>Despal</i> , <i>Dara Melisa</i>)
	ORKING GROUP 3: SCIENCE-POLICY INTERFACING ON O-RESOURCE CONSERVATION AND UTILIZATION
1.	Tropical rain forest in Kalimantan as source of medicinal plants:
	a case study at Dayak Meratus ethnic in South Kalimantan (<i>Yudi Funanul Arifin, Siti Hamidah</i>)
2.	Development of Gambir (<i>Uncaria gambir</i>) for rural economy:
3.	between policy and people aspiration (<i>Andy Ahmad Zaelany</i>)166 Conservation of wildlife bio-resource management for livelihood
٥.	(Retno Iswarin Pujaningsih)
4.	Market integration analysis of sweetened condensed milk in
	In mesia: do sweetened condensed milk prices follow the prices
	of nported milk and sugar? (Venty Fitriany Nurunisa, Bonar M. Sir aga, Ratna Winandi A., Bernhard Brümmer)
5.	Trend analyses of forest and land fires towards climate change in
	Indonesia (Lailan Syaufina)
6.	Agroforestry based medicinal plants and marketing partnership
	for community empowerment: cases in Bogor district and Surabumi district, West Java province (Leti Sundawati, Ninuk
	Pubnaningsih, Edy Djauhari Purwakusumah)
7.	A survey on the community socio-economic of the district of
	coral reef rehabilitation and management program (COREMAP)
0	of Sikka, Flores Flores (Vincentius Repu)
8.	Influence of leadership style, organizational culture, and work motivationon employee performance in public company pawn
	shop branch office in Kupang City, East Nusa Tenggara,
	Indonesia (Fred Marthinus Dethan)
9.	Management of natural resources in tropical peat swamp forest
10	of Indonesia (<i>Ujang Suwarna</i>)
10.	Kota Gajah Sub-district, Lampung Tengah District, Lampung
	Province (Angga Yudhistira, Harianto, Bernhard Brümmer,
	Stephan Wessels, Nunung Kusnadi)219
11.	The sustainability of coffee plantation in West Lampung,
12	Lampung province, Indonesia (<i>Yeti Lis Purnamadewi</i>)
	customary community approaching in national park management
	in Indonesia (Nandi Kosmaryandi, Sambas Basuni, Lilik B
	Prasetyo, Soeryo Adiwibowo)233



1. Dilarang mengutip sebagian atau seluruh karya tulis ini tanpa mencantumkan dan menyebutkan sumber: Hak Cipta Dilindungi Undang-Undang

13.	Competitiveness analysis of Indonesian shrimp farming, case	
	study: PT. Surya Windu Kencana (SWK), East Java (Siti	
	Maryam, Bernhard Brümmer, Gabriele Hörstgen-Schwark,	
	Rachmat Pambudy)	242
14.	Competitiveness and policy impact analysis of feedlots in	
	Lampung (Labudda Paramecwari, Bernhard Brummer, Stefan	
	Schwarze, Rachmat Pambudy)	248
15.	The Contribution of Agricultural Sector Towards Culinary	
	Business Development at Kupang Municipality East Nusa	
	Tenggara Province (Markus Bunga)	254
16.	The Morphological Character of the "Bendi" Horse as Short	
	Distance Urban Transport Modes that are Environmentally	
	Friendly (Sri Adiani, Dordia A. Rotinsulu, Ben J Takaendengan)	261
17.	The diversity of fungi on polluted mangrove ecosystem at	
	Belawan and Jaring Halus, North Sumatra province (Yunasfi,	
	Pindi Patana)	266
18.	Natural products exploration in frame of tropical plant bio-	
	resource conservation and utilization (Enih Rosamah, Harlinda	
	Kuspradini, Rita Khairani)	273
19.	The impact of trade policy on international palm oil trade flows	
	(Biska Pujiati, M Firdaus, Andriyono K Adhi)	278
LIC	STOF PARTICIPANTS	
	orr	204
CO	MAITTEE	287

Bogor Agricultural University

Dilarang mengutip sebagian atau seluruh karya tulis ini tanpa mencantumkan dan menyebutkan sumber Hak Cipta Dilindungi Undang-Undang

Influence of tannin concentration in ration on fermentation parameters of Rumen Simulation Technique (RUSITEC): a meta-analysis

Anuraga Jayanegara^{1,*}, M. Ridla¹, Erika B. Laconi¹, Nahrowi¹

Department of Nutrition and Feed Technology, Bogor Agricultural University, Bogor, 16680, Indonesia

* Corresponding author: anuragaja@apps.ipb.ac.id

Alstract The objective of this study was to summarize and to quantify the effects of tannin on fermentation parameters of rumen simulation technique (RESITEC) through a meta-analysis approach. Experiments reporting tamin concentration and rumen fermentation using the RUSITEC system were integrated into a database, and comprised of 6 studies and 25 treatments. Parameters recorded were nutrient digestibility, gas production, methane (CH₄) emission, short-chain fatty acid (SCFA) profiles, pH, ammonia (NH₃) and microbial population (bacteria and protozoa). The analysis of the data assembled in the database was based on mixed model methodology in which different studies were treated as random effects whereas tannin concentration was treated as fixed effects. Results revealed that in vitro crude protein digestibility (CPD), neutral detergent fiber digestibility (NDFD) and acid detergent fiber digestibility (ADFD) decreased linearly as the tannin concentration increased with the P-value of 0.047, 0.005 and 0.004, respectively. Comparing the magnitude of reduction in CP and fiber digestibility by the influence of tannin, the compound appeared to cause higher negative effect on CP digestibility than that of fiber as indicated by the slopes. However, the overall OMD did not significantly decrease although the slope remained negative. Methane emission tended to decrease at higher tannin concentration when expressed per unit of substrate (P=0.066) and significantly decreased when expressed per unit of total gas produced (P=0.005). It can be concluded that tannin is a potential compound for mitigating ruminal methane emission but its use on the other hand also decrease nutrient digestibility.

Keywords tannin, rumen, fermentation, rusitec, meta-analysis

1. Introduction

Tamin is among plant secondary compounds produced by plants in their intermediary metabolism. It is polyphenolic compound with diverse

niversity

structure (such as between hydrolysable and condensed tannin) and molecular weight but has similar property: it binds and precipitates protein [1]. With respect to ruminant nutrition, tannin is considered to have both pheneficial and detrimental nutritional effects. Some of the beneficial effects of tannin are better utilization of dietary protein, faster growth rate, higher milk yield and improved animal health through prevention of bloat and infection. Negative effects of tannin have been associated with its roxicity or rumen microbes and the animals especially when present at high acconcentration in ration (>50 g/kg dry matter) [2].

Research on tannin in relation to ruminant nutrition has been conducted under various experimental conditions, either *in vivo* (directly to the animal), in sacco (fistulated animal) or in vitro (laboratory equipment that mimic rumen fermentation). The latter method is divided into two groups, i.e. in vitro batch and in vitro continuous culture. Research synthesis of tannin effect of rumen digestion and fermentation based on in vivo and in vitro batch studies across various ruminant species has been previously performed through a meta-analytical approach [3]. However, the studies based of in vitro continuous culture like in rumen simulation technique or RUSITEC [4] have not been summarized. In the present study, therefore, a meta-analysis was conducted to summarize and to quantify the effects of tannin concentration on fermentation parameters of RUSITEC.

2. Materials and Methods

RUSITEC experiments reporting tannin concentration and rumen fermentation were integrated into a database (comprised of 6 studies and 25 treatments). The studies were Sliwinski et al. [5], Hess et al. [6-7], Tiemann et al. [8], Bekele et al. [9] and Khiaosa-ard et al. [10] (Table 1). Parameters recorded were nutrient digestibility, gas production, methane (CH₄) emission, short-chain fatty acid (SCFA) profiles, pH, ammonia (NH₃) and microbial population (bacteria and protozoa). Tannin forms were either from non-extracted or extracted tannins of plant origins, and constituted of different annin types, i.e. hydrolysable, condensed or unspecified or mixed tannins. Such different tannin types were not a main point of interest but rather the amount or concentration of the tannin in the ration. Therefore, they were not stated as a categorical variable and not included in the statistical model. Studies reporting treatments with addition of polyethylene glycol (PEG) were excluded from the database since the substance may neutralize the effects of tannins under rumen environment.

2. Dilarang mengumumkan dan memperbanyak sebagian atau seluruh karya tulis ini dalam bentuk apapun tanpa izin IPB . Pengutipan hanya untuk kepentingan pendidikan, penelitian, penulisan karya ilmiah, penyusunan laporan, penulisan kritik atau tinjauan suatu masalah

Dilarang mengutip sebagian atau seluruh karya tulis ini tanpa mencantumkan dan menyebutkan sumber Hak Cipta Dilindungi Undang-Undang

The analysis of the data assembled in the database was made by a statistical meta-analysis approach [11]. Studies were treated as random effects whereas tannin concentration was treated as fixed effects using MIXED procedure of SAS version 9.2. The following statistical model was employed:

$$Y_{ij} = B_0 + B_1 X_{ij} + s_i + b_i X_{ij} + e_{ij}$$

where Y_{ij} = dependent variable, B_0 = overall intercept from all studies (fixed effect), B_1 = linear regression coefficient of Y on X (fixed effect), X_{ij} = value of the continuous predictor variable (dietary tannins), s_i = random effect of study i, b_i = random effect of study i on the regression coefficient of Y on X in study i, and e_{ij} = the unexplained residual errors. The study variable was declared in the CLASS statement since it does not contain any quantitative information. Data were weighted by the number of replicates each study and scaled to 1 to take into consideration of unequal variance among studies. Microbial population data were transformed into their logarithmic units to allow linear relationships with the independent variable. Moder statistics presented are P-value and coefficient of determination (R^2).

Table 1. Studies included in the meta-analysis of the effect of tannin concentration on fermentation parameters of rumen simulation technique (RUSITEC)

Stud no.	y Reference	Basal feed	Tannin source	Tannin level (g/kg DM)
1	Sliwinski et al. (2002a)	Grass hay, silage and barley	Chestnut	0 to 2.5
2	Hess et al. (2006)	Koronivia grass	Cratylia argentea , Calliandra calothyrsus	0 to 135
3	Hess et al. (2008)	Koronivia grass	Leucaena leucocephala, Flemingia macrophylla, Calliandra calothyrsus	0 to 62.2
4	Giemann et al.	Koronivia grass	Vigna unguiculata, Calliandra calothyrsus	0 and 71
5	Bekele et al. (2009)	Koronivia grass	Samanea saman, Acacia angustissima, Sesbania sesban, Cajanus cajan	0 to 45
6	Chiaosa-Ard et al.	Grass-clover hay	Onobrychis viciifolia, Acacia mearnsii	0 and 78.9



3. Results and Discussion

In vitro crude protein digestibility (CPD), neutral detergent fiber digestibility (NDFD) and acid detergent fiber digestibility (ADFD) decreased linearly as the tannin concentration increased with the *P*-value of 0.047, 0.005 and 0.004, respectively (Table 2). Comparing the magnitude of reduction in CP and fiber digestibility by the influence of tannin, the compound appeared to cause higher negative effect on CP digestibility than that of fiber as indicated by the slopes. An increase of tannin concentration by 1 gag declined CPD by 2.921 mg/g. The decrease was lower for the NDFD and ADFD, i.e. 1.231 and 1.549 mg/g, respectively. However, these figures might change if the slopes were corrected by the intercepts due to substantial different digestibility between CP and fiber at dictary tannins substantial different digestibility between CP and fiber at dietary tannins equal to 0 g/kg. All of these relationships had high R², i.e. higher than 0.4. The results support a theory that tannin may form complexes with some nutrients such as protein and carbohydrate and, therefore, may reduce their digestibility in the digestive tract of ruminants [1-2, 12]. However, the overall OMD did not significantly decrease although the slope remained negative. This was also the case for the total gas production.

Methane emission tended to decrease at higher tannin concentration when expressed per unit of substrate (P=0.066) and significantly decreased when expressed per unit of total gas produced (P=0.005). The latter had a high \mathbb{R}^2 , i.e. 0.677. Explanation of the methane decrease due to tannin appears to be because of the decrease in digestibility of nutrients, particularly fiber, which decreases H₂ production as a substrate for methanogenesis as well as direct inhibition on methanogen population; the latter occurs since tannin is toxic to some groups of rumen microbes including the methanogen [3, 13]. Tannins had almost no effects on all SCFA variables, except that the substance linearly decreased C₄ (P=0.013, R²=0.403). Dietary tannin had also no significant effects on ruminal pH, NH3, bacteria and protozoa population.

Table 2 Equations for linear regression of the effect of tannin concentration on fermentation parameters of rumen simulation technique (RUSITEC)

_(110	0111							
Response	n	n Parameter estimates						R ²
parameter		Intercept	SE	P intercept	Slope	SE slope	P slope	
			intercept			-		
OMD (mg/g)	25	441.9	36.81	< 0.001	-0.672	0.3918	TiS	0.178
CPD (mg/g)	14	644.2	76.19	0.014	-2.921	1.2914	0.047	0.407
NDFD (mg/g)	25	323.6	26.56	< 0.001	-1.231	0.3801	0.005	0.411

Response	n	Parameter estimates					R^2	
parameter		Intercept	SE	P intercept	Slope	SE slope	P slope	
			intercept					
ADFD (mg/g)	14	277.1	20.42	< 0.001	-1.549	0.4102	0.004	0.543
Gas (ml/g)	14	81.1	28.69	Ns	-0.170	0.1298	ns	0.220
CH ₄ (ml/g)	25	10.9	1.89	0.002	-0.0255	0.01301	0.066	0.231
CH4 (ml/l gas)	14	170.4	47.73	0.07	-0.582	0.1631	0.005	0.677
Total SCFA (mmol/l)	23	83.5	10.89	0.002	-0.0003	0.04986	ns	0.000
% total SCFA)	23	63.5	1.84	< 0.001	0.0002	0.00929	ns	0.000
C ₹	23	22.2	1.83	< 0.001	0.0155	0.01129	ns	0.140
C. cista mistCs	23	10.9	1.99	0.005	-0.0139	0.00501	0.013	0.403
is <mark>@</mark> -C4	19	0.72	0.229	0.052	0.0007	0.00117	ns	0.042
Ca	19	2.75	0.894	0.054	-0.0007	0.00166	ns	0.019
iso-C5	19	0.89	0.307	0.063	-0.0011	0.00101	ns	0.126
isa SCFA	19	1.61	0.349	0.019	-0.0005	0.00166	ns	0.009
C27C3	23	2.93	0.234	< 0.001	-0.0011	0.00166	ns	0.036
pI Z Iq	25	7.00	0.050	< 0.001	0.0001	0.00035	ns	0.010
N (mmol/I)	25	6.44	1.862	0.018	-0.0285	0.01822	ns	0.155
log bacteria (10 /ml)	23	9.11	0.313	< 0.001	0.0008	0.00063	ns	0.114
log protozoa	23	3.58	0.127	< 0.001	0.0004	0.00127	ns	0.008

ADFD, in vitro acid detergent fiber digestibility; C2, acetate; C3, propionate; C4, butyrate; CPD, in vitro apparent crude protein digestibility; n, number of treatment; NDFD, in vitro neotral detergent fiber digestibility; OMD, in vitro organic matter digestibility; R2, coefficient of determination; SCFA, short chain fatty acids; SE, standard error.

4. Conclusion

Tannin is a potential natural compound for mitigating ruminal methane emission but its use on the other hand also decrease nutrient digestibility. Further investigation is therefore required to determine an optimum concentration of tannin in ration in which it mitigates methane emission and simultaneously supports rumen digestion and fermentation.

59 References

- [1] H.P.S. Makkar. 2003. Effects and fate of tannins in ruminant animals, adaptation to tannins, and strategies to overcome detrimental effects of feeding tannin-rich feeds. Small Rum. Res. 49: 241-256.
- [2] I. Mueller-Harvey. 2006. Unravelling the conundrum of tannins in animal nutrition and health. J. Sci. Food Agric. 86: 2010-2037.
- [3] A. Jayanegara, F. Leiber and M. Kreuzer. 2012. Meta-analysis of the relationship between dietary tannin level and methane formation in



- ruminants from in vivo and in vitro experiments. J. Anim. Physiol. Anim. Nutr. 96: 365-375.
- J.W. Czerkawski and G. Breckenridge. 1977. Design and development of a long-term rumen simulation technique (Rusitec). Brit. J. Nutr. 38: 371-384.
- B.J. Sliwinski, C.R. Soliva, A. Machmüller and M. Kreuzer. 2002. Efficacy of plant extracts rich in secondary constituents to modify rumen fermentation. Anim. Feed Sci. Technol. 101: 101-114.
- H.D. Hess, T.T. Tiemann, F. Noto, S. Franzel, C.E. Lascano and M. Krauzer. 2006. The effects of cultivation site on forage quality of Calliandra calothyrsus var. Patulul. Agrofor. Syst. 68: 209-220.
- H.B. Hess, M.L. Mera, T.T. Tiemann, C.E. Lascano and M. Kreuzer. 20. In vitro assessment of the suitability of replacing the low-tannin legume Vigna unguiculata with the tanniniferous legumes Leucaena leucocephala or Calliandra calothyrsus in a tropical grass diet. Anim. Feet Sci. Technol. 147: 105-115.
- T.T. Tiemann, C.E. Lascano, M. Kreuzer and H.D. Hess. 2008. The ruminal degradability of fibre explains part of the low nutritional value and reduced methanogenesis in highly tanniniferous tropical legumes. J. Sci. Food Agric. 88: 1794-1803.
- A.Z. Bekele, C. Clement, M. Kreuzer and C.R. Soliva. 2009. Efficiency of Sesbania sesban and Acacia angustissima in limiting mechanogenesis and increasing ruminally available nitrogen in a tropical grass-based diet depends on accession. Anim. Prod. Sci. 49: 145-153.
- [10] R. Khiaosa-ard, S.F. Bryner, M.R.L. Scheeder, H.R. Wettstein, F. Leiber, M. Kreuzer and C.R. Soliva. 2009. Evidence for the inhibition of the terminal step of ruminal α-linolenic acid biohydrogenation by condensed tannins. J. Dairy Sci. 92: 177-188.
- [11] D. Sauvant, P. Schmidely, J.J. Daudin and N.R. St-Pierre. 2008. Metaanalyses of experimental data in animal nutrition. Animal 2: 1203-1214.
- [12] N. Silanikove, A. Perevolotsky and F.D. Provenza. 2001. Use of tankin-binding chemicals to assay for tannins and their negative postingestive effects in ruminants. Anim. Feed Sci. Technol. 91: 69-810
- [13] M.E. Tavendale, L.P. Meagher, D. Pacheco, N. Walker, G.T. Attwood and S. Sivakumaran. 2005. Methane production from in vitro rumen incubations with Lotus pedunculatus and Medicago sativa, and effects of extractable condensed tannin fractions on methanogenesis. Anim. Feed Sci. Technol. 123-124: 403-419.