

**Volume 12,
Number 1,
January-June 2016**

ISSN 0973-1768

**GLOBAL JOURNAL *of*
PURE and APPLIED
MATHEMATICS**

**Editor-in-chief:
Taekyun Kim**

**Managing Editor:
E.-B. Lin**

RIP Research India Publications
<http://www.ripublication.com>

RIP **RESEARCH**
INDIA
PUBLICATIONS

Research India Publications was established in 1998 and today we are one of the leading International Publishers, Distributors and Subscription Agents of books and research journals. Research India Publications is a privately owned publishing company dedicated to the global dissemination of information. We acquire, develop, market, and distribute knowledge by disseminating scholarly and professional materials throughout the world. All of journals published by us maintain the highest standards of quality, with Editorial Boards composed of scholars from around the world.

Contact Us:

B-2/84, Ground Floor, Rohini Sector-16, Delhi-110089 INDIA
Tel No.: 011-45587661, 011-27893171, +91-11-65394240, Fax No.: +91-11-27297815

Emails:

General query: info@ripublication.com
Subscription/renew orders: subscription@ripublication.com
Submission of new journal/book proposal: proposal@ripublication.com
Other services: services@ripublication.com

Website:

<http://www.ripublication.com>

© All rights are reserved with Research India Publications

Global Journal of Pure and Applied Mathematics
ISSN 0973-1768

Aims and Scope: Global Journal of Pure and Applied Mathematics (GJPAM) is an international journal of high quality devoted to the publication of original research papers from pure and applied mathematics with some emphasis on all areas and subareas of mathematical analysis and their broad range of applications. Areas and subareas of interest include (but are not limited to) approximation theory; statistics; probability; fluid mechanics; Fuzzy mathematics; transport theory; mathematical biology, including population dynamics; wave propagation; special functions; algebra and applications; combinatorics; coding theory; fractional analysis; solid mechanics; variational methods; financial mathematics; asymptotic methods; graph theory; fractals; moment theory; scattering theory; number theory and applications; geometry of Banach spaces; topology and applications; complex analysis; stochastic process; bifurcation theory; differential equations; difference equations; dynamical systems; functional differential equations; numerical analysis; partial differential equations; integral equations; operator theory; Fourier analysis; matrix theory; semigroups of operators; mathematical physics; convex analysis; applied harmonic analysis; optimization; wavelets; signal theory; ultrametric analysis; optimal control; fixed-point theory and applications; reaction-diffusion problems, discrete mathematics; automata theory and more. **Submission:** Authors are requested to submit their papers electronically to the email submit@ripublishation.com with mention journal title (GJPAM) in subject line. **Indexing and Abstracting:** GJPAM is abstracted and indexed in the SCOPUS, Mathematical Reviews, MathSciNet, Zentralblatt MATH and EBSCO databases. **Frequency:** Six issues per year.

Publication Time: Author will receive print journal copy within 15 days after payment of publication charges and submission of copyright form. **Publication Cost:** Authors of accepted papers are requested to arrange page charges of their papers at the rate of (Rs.9500 per paper for Indian authors and US\$275 per paper for foreign authors), which will be invoiced, through the author, to the author's company, institute, or agency. Twenty Five reprints of a paper are provided to the authors ex-gratis. Additional sets of reprints may be ordered at the time of proof correction. Page charges are mandatory and omitted only when it will be highly recommended by Editors-in-chief to us for publication.

Style: Manuscripts should be attached and sent to the Editor via e-mail. The file has to be written using the following software: PDF, TeX/ LaTeX. When submitting manuscripts via mail, please send three copies (double-spaced, single-sided, 12-point font, 1-inch margins all around). An un-numbered cover page should include the title and authors' names, addresses and email addresses. Manuscripts submitted to GJPAM should contain: Title of article, Authors' names, Authors' Institutional Affiliations, Abstract (200-300 words suitable for abstracting services), Introduction, Experimental (Materials and Methods), Results, Discussion, Conclusion, References, Tables (each starting with a caption) and Figure captions, List of symbols and Abbreviations, Numbered list of references, Biographical sketch, and Original figures. **Length:** An GJPAM Transactions paper should not exceed 25 typeset, printed pages.

Order of parts: Title of article, authors' names, authors' institutional affiliations, abstract (200-300 words suitable for abstracting services), main text, tables (each starting with a caption) and figure captions, list of symbols and abbreviations (if necessary), numbered list of references, biographical sketch, and original figures.

Title: The title of the paper should be concise and definitive (with key words appropriate for retrieval purposes).

Author names and affiliations: It is GJPAM policy that all those who have participated significantly in the technical aspects of a paper be recognized as co-authors or cited in the

acknowledgments. In the case of a paper with more than one author, correspondence concerning the paper will be sent to the first author unless staff is advised otherwise. Author name should consist of first name, middle initial, last name. The author affiliation should consist of the following, as applicable, in the order noted:

- Company or college (with department name of company division)
- Postal address
- City, state, zip code
- Country name
- Telephone, fax, and e-mail

Abstract: A short abstract (200-300 words maximum) should open the paper. The purposes of an abstract are:

1. to give a clear indication of the objective, scope, and results of the paper so that readers may determine whether the full text will be of particular interest to them;
2. to provide key words and phrases for indexing, abstracting, and retrieval purposes.

The abstract should not attempt to condense the whole subject matter into a few words for quick reading.

Figures should be sent as: fair copy on paper, whenever possible scaled to about 200%, or as EPS file (Encapsulated PostScript). In addition, all figures and tables should be numbered and the appropriate space reserved in the text, with the insertion point clearly indicated.

The final decision on publication is made by the Editor-in-Chief upon recommendation of Editorial board member. If the manuscript is accepted for publication in GJPAM, it must not be published in any periodical elsewhere without the permission of the Publisher.

Copyright: Each manuscript must be accompanied by a statement that it has been neither published nor submitted for publication, in whole or in part, either in a serial, professional journal or as a part in a book which is formally published and made available to the public. For the mutual benefit and protection of authors and publishers it is necessary that authors provide formal written consent to publish and transfer of Copyright Form before publication of the work.

Proofs: Proofs will be sent to the corresponding author and should be returned with one week of receipt. Corrections should be restricted to typesetting errors; any other corrections may be charged to the authors. Authors are advised to check their proofs very carefully before return, since the inclusion of late corrections cannot be acceptable. Corrected proofs are to be returned to the publishers.

Editorial Board

Editor-in-Chief:

Taekyun Kim, Kwangwoon University, Korea

Managing Editor:

E.-B. Lin, University of Toledo, USA

Associate Editor:

Sunil Mathur, University of Memphis, USA

Editorial Board Members:

S.D. Rajan, Al Musanna College of Technology, Oman, **M. Bohner**, University of Missouri at Rolla, USA, **S. Mathur**, University of Mississippi, USA, **G.M. N'Guerekata**, Morgan State University, USA, **R.P. Agarwal**, Florida Institute of Technology, USA, **N.U. Ahmed**, University of Ottawa, Canada, **G.A. Anastassiou**, University of Memphis, USA, **B.S. Attili**, United Arab Emirates University, UAE, **P.H. Bezandry**, Howard University, USA, **D. Bugajewska**, Adam Mickiewicz University, Poland, **E. Camouzis**, The American College of Greece, Greece, **C. Cesarano**, Università Campus Bio-Medico di Roma, Italy, **S.S. Cheng**, National Tsing Hua University, Taiwan, **A. Dermoune**, Université des Sciences et Technologies de Lille, France, **S.S. Dragomir**, Victoria University, Australia, **P. Eloe**, University of Dayton, USA, **A. Fiorenza**, Università' di Napoli Federico II, Italy, **D. Girela**, Universidad de Malaga, Spain, **J.R. Graef**, University of Tennessee at Chattanooga, USA, **S. Hamadene**, Université du Maine, France, **O. Hili**, National Polytechnic Institute of Yamoussoukro, Ivory Coast, **G.R. Hiremath**, University of North Carolina at Pembroke, USA, **M. Ito**, Kyoto Sangyo University, Japan, **H. Th. Jongen**, RWTH Aachen University, Germany, **I.B. Jung**, Kyungpook National University, Korea, **A. Kamal**, Sultan Qaboos University, Oman, **N. Kenmochi**, Chiba University, Japan, **H. Kitada**, University of Tokyo, Japan, **J. Koolen**, Pohang University of Science and Technology, Korea, **T. Kwembe**, Jackson State University, USA, **T. Kusano**, Fukuoka University, Japan, **D.R. Larson**, Texas A&M University, USA, **C.-O. Lee**, KAIST, Korea, **E.-B. Lin**, University of Toledo, USA, **W.B. Liu**, University of Kent, UK, **G. Lu**, Wayne State University, USA, **W.-X. Ma**, University of South Florida, USA, **M.F. Mahmood**, Howard University, USA, **G.M. Mahmoud**, Assiut University, Egypt, **T.M. Mills**, La Trobe University, Australia, **T. Nogura**, Ehime University, Japan, **H. Ouerdiane**, Campus Universitaire, Tunisia, **Y. Ouknine**, Université Cadi Ayyad, Morocco, **A. Peterson**, University of Nebraska-Lincoln, USA, **J.M. Rassias**, National and Capodistrian University of Athens, Greece, **S. Reich**, Israel Institute of Technology, Israel, **S. Saitoh**, Gunma University, Japan, **M.Z. Abu-Sbeih**, King Fahd University of Petroleum and Minerals, Saudi Arabia, **N.C. Sacheti**, Sultan Qaboos University, Oman, **S.M. El-Sayed**, Education College for Girls, Saudi Arabia, **M. Sifi**, Campus Universitaire, Tunisia, **C.W. Soh**, Jackson State University, USA, **A. Soufyane**, United Arab Emirates University, UAE, **F. Wagemann**, Université de Nantes, France, **S. Watanabe**, Gunma University, Japan, **H.K. Xu**, University of Kwazulu-Natal, South Africa, **A.A. Yakubu**, Howard University, USA, **K. Yoshimoto**, Nihon University, Japan, **X. Zha**, Middle Tennessee State University, USA.

Contents

Inverse Perfect Domination in Graphs <i>Daisy P. Salve and Enrico L. Enriquez</i>	1 - 10
Optimizing the Coordinates Position for Optimal Placement of ZigBee for Power Monitoring From Remote Lokation <i>Narendra B Soni, SMIEE, Kamal Bansal and Devendra Saini</i>	11 - 17
Method for Assessment of Sectoral Efficiency of Investments Based on Input-Output Models <i>Anton Kogan</i>	19 - 32
Connection of two nodes of a transportation network by two disjoint paths <i>Ilya Abramovich Golovinskii</i>	33 - 55
A mathematical model of thyroid tumor Eugeny Petrovich Kolpak, Inna Sergeevna Frantsuzova and Kabrits Sergey <i>Alexandrovich</i>	56 - 66
Trigonometrically-fitted explicit four-stage fourth-order Runge–Kutta–Nyström method for the solution of initial value problems with oscillatory behavior <i>M. A. Demba and N. Senu and F. Ismail</i>	67 - 80
Functions fixed by a weighted Berezin transform <i>Jaesung Lee</i>	81 - 86
Traveling Wave Solutions for the Nonlinear Boussinesq Water Wave Equation <i>Jabaruddin</i>	87 - 93
On strongly left McCoy rings <i>Sang Jo Yun, Sincheol Lee, Sung Ju Ryu, Yunbo Shin and Hyo Jin Sung</i>	95 - 104

On convergence theorems for new classes of multivalued hemicontractive-type mappings	105 - 116
<i>Samir Dashputre and Apurva Kumar Das</i>	
CESRO and HLDR Mean of Product Summability Methods	117 - 124
<i>Suyash Narayan Mishra, Piyush Kumar Tripathi and Manisha Gupta</i>	
Equitable, restrained and k-domination in intuitionistic fuzzy graphs	125 - 145
<i>G. Thamizhendhi and R. Parvathi</i>	

Traveling Wave Solutions for the Nonlinear Boussinesq Water Wave Equation

Jaharuddin

*Department of Mathematics,
Bogor Agricultural University,
Jl. Meranti, Kampus IPB Dramaga,
Bogor 16880, Indonesia.*

Abstract

The Boussinesq equation is one of the nonlinear evolution equations which describes the model of shallow water waves. By using the modified F-expansion method, we obtained some exact solutions. Some exact solutions expressed by hyperbolic function and exponential function are obtained. The results show that the modified F-expansion method is straightforward and powerful mathematical tool for solving nonlinear evolution equations in mathematical physics and engineering sciences.

AMS subject classification: 35C08, 35K99, 35P05.

Keywords: F-expansion method, traveling wave solutions, Boussinesq equation.

1. Introduction

The Boussinesq equation appears in studying the transverse motion and nonlinearity in acoustic waves on elastic rods with circular cross section. This equation arises in other physical applications such as nonlinear lattice waves [3], ion sound waves in plasma [7], and in vibrations in a nonlinear string [10]. One of the fundamental problems for these equation is to obtain their traveling waves solution as well as solitary wave solution. Traveling wave solutions represents an important type of solutions for nonlinear partial differential equations as many nonlinear PDE have been found to have a variety of traveling wave solutions. It is well-known that the investigation of the exact solutions of nonlinear PDE's plays an important role in the study of nonlinear physical phenomena. The sixth-order Boussinesq equation is one the nonlinear PDE's to be discussed. Many researchers studied the sixth-order Boussinesq equation to construct analytical solutions

by using different methods. For instance, H. Naher, F.A. Abdullah, and M.A. Akbar [6] implemented the exp-function method to investigate this equation for obtaining traveling wave solutions. In [4], M. Hosseini, H. Abdollahzadeh, and M. Abdollahzadeh employed the sin-cosine method to construct exact solutions of the same equation. A general approach to construct exact solution to the Boussinesq equation is given by Clarkson [2] has deduced conservation laws. Wazwaz [8] obtained the single soliton solutions of the sixth-order Boussinesq equation using the tanh method and multiple soliton solutions using Hirota bilinear method. Wang and Li [9] presented a powerful method which is called the F-expansion method. The F-expansion method is an effective and direct algebraic method for finding the exact solutions of nonlinear evolution equation, many nonlinear equations have been successfully solved. The F-expansion method gives a unified formation to construct various traveling wave solutions and provides a guideline to classify the various types of traveling wave solutions. Later, the further develop methods named the modified F-expansion. Using the new method, exact solutions of many nonlinear evolution equations are successfully obtained.

The purpose of this paper is to implement the modified F-expansion method to search traveling wave solutions for the sixth-order Boussinesq equation. In former literature, the traveling wave solutions to sixth-order the Boussinesq equation have not been studied by this method. The paper is organized as follows. In Section 2, the modified F-expansion method is introduced briefly. Section 3 is devoted to applying this method to the sixth-order nonlinear Boussinesq water wave equation. The last section is a short summary and discussion.

2. Description of the method

Based on [1], the main procedures of modified F-expansion method are as follows. Consider the nonlinear partial differential equation in the form:

$$G \left(u, \frac{\partial u}{\partial t}, \frac{\partial u}{\partial x}, \frac{\partial^2 u}{\partial x^2}, \frac{\partial^2 u}{\partial x \partial t}, \dots \right) = 0 \quad (2.1)$$

where $u(x, t)$ is a traveling wave solution of equation (2.1), G is a polynomial in $u(x, t)$ and its partial derivatives in which the highest order derivatives and the nonlinear terms are involved. We use the transformation

$$\xi = \alpha(x - ct), \quad (2.2)$$

where c and α are constants. Using (2.2) to transfer the nonlinear partial differential equation (2.1) to nonlinear ordinary differential equation for $U = U(\xi)$

$$H \left(U, \frac{dU}{d\xi}, \frac{d^2U}{d\xi^2}, \frac{d^3U}{d\xi^3}, \dots \right) = 0. \quad (2.3)$$

Suppose that the solution of equation (2.3) can be written as follows:

$$U(\xi) = A_0 + \sum_{k=1}^N (A_k (F(\xi))^k + B_k (F(\xi))^{-k}) \quad (2.4)$$

where A_0, A_k, B_k ($k = 1, 2, \dots, N$) are constants to be determined later. The positive integer N can be determined by considering the homogenous balance between the highest order derivatives and the nonlinear terms appearing in equation (2.3), and $F(\xi)$ satisfies the first-order linear ordinary differential equation:

$$\frac{dF}{d\xi} = h_0 + h_1F + h_2F^2, \tag{2.5}$$

where h_0, h_1, h_2 are the arbitrary constants to be determined later. The special condition chosen for the $h_0, h_1,$ and $h_2,$ we can obtain the solution of equation (2.5). For example, when $h_1 = 0,$ and $h_2 = -h_0,$ the solution of equation (2.5) is

$$F(\xi) = \tanh(h_0\xi).$$

Next, when $h_2 = 0,$ and $h_1 = -h_0,$ the solution of equation (2.5) is

$$F(\xi) = \frac{1}{h_1} \exp(h_1\xi) + 1.$$

Substituting (2.4) into (2.3) and using (2.5), we obtain a polynomial in $(F(\xi))^k$ ($k = 0, \pm 1, \pm 2, \dots$). Equating each coefficient of the resulting polynomial to zero yields a set of algebraic equations for $c, \alpha, h_0, h_1, h_2,$ and A_0, A_k, B_k ($k = 1, 2, \dots, N$). Assuming that the constants A_0, A_k, B_k ($k = 1, 2, \dots, N$) can be obtained by solving the algebraic equations and then substituting these constants and the solution of (2.5), depending on the special conditions chosen for the $h_0, h_1,$ and h_2 we can obtain the explicit solutions of equation (2.1) immediately. Obviously, different types of exact solutions of the equation (2.5) may result in different types of exact traveling wave solutions for equation (2.1).

3. Application of the method

In this section, we apply the modified F-expansion method to construct exact traveling wave solutions for the sixth-order Boussinesq equation. Let us consider the equation

$$\begin{aligned} \frac{\partial^2 u}{\partial t^2} - \frac{\partial^2 u}{\partial x^2} - 15u \frac{\partial^4 u}{\partial x^4} - 30 \frac{\partial u}{\partial x} \frac{\partial^3 u}{\partial x^3} - 15 \left(\frac{\partial^2 u}{\partial x^2} \right)^2 \\ - 45u^2 \frac{\partial^2 u}{\partial x^2} - 90u \left(\frac{\partial u}{\partial x} \right)^2 - \frac{\partial^6 u}{\partial x^6} = 0. \end{aligned} \tag{3.1}$$

The equation (3.1) was introduced by J.V. Boussinesq (1842–1929) to illustrate the propagation of long waves on the surface of water with small amplitude [5], where $u(x, t)$ is the elevation of the free surface of the fluid. Making a transformation $u(x, t) = U(\xi)$ with $\xi = \alpha(x - ct),$ equation (3.1) can be reduced to the following ordinary differential equation:

$$(c^2 - 1) \frac{\partial^2 U}{\partial \xi^2} - 15\alpha^2 u \frac{\partial^4 u}{\partial \xi^4} - 30\alpha^2 \frac{\partial u}{\partial \xi} \frac{\partial^3 u}{\partial \xi^3} - 15\alpha^2 \left(\frac{\partial^2 u}{\partial \xi^2} \right)^2 \tag{3.2}$$

$$-45u^2 \frac{\partial^2 u}{\partial \xi^2} - 90u \left(\frac{\partial u}{\partial \xi} \right)^2 - \alpha^4 \frac{\partial^6 u}{\partial \xi^6} = 0,$$

where c is wave velocity which moves a long the direction of horizontal axis and α is the wave number. By integrating equation (3.2) twice with respect to the variable ξ and assuming a zero constants of integration, we have

$$(c^2 - 1)U - 15\alpha^2 U \frac{\partial^2 u}{\partial \xi^2} - 15U^3 - \alpha^4 \frac{\partial^4 U}{\partial \xi^4} = 0. \quad (3.3)$$

The homogenous balance between U^3 and $\frac{\partial^4 U}{\partial \xi^4}$ in equation (3.3) implies $n = 2$. Suppose that solution of equation (3.3) is of the following form

$$U(\xi) = A_0 + A_1 F(\xi) + A_2 F^2(\xi) + \frac{B_1}{F(\xi)} + \frac{B_2}{F^2(\xi)} \quad (3.4)$$

where A_0, A_1, A_2, B_1 and B_2 are constants to be determined. Substituting equation (3.4) along with (2.5) into (3.3) and then setting all the coefficients of $F^k(\xi)$, ($k = -6, -5, \dots, 5, 6$) of the resulting systems to zero, we can obtain a system of algebraic equations for $c, \alpha, h_0, h_1, h_2, A_0, A_1, A_2, B_1$ and B_2 as follows

$$\begin{aligned} F^6 & : (-90a^2 A_2^2 h_2^2 - 15A_2^3 - 120a^4 A_2 h_2^4) = 0, \\ F^k & : f_k(c, \alpha, h_0, h_1, h_2, A_0, A_1, A_2, B_1, B_2) = 0, \\ F^{-6} & : (-90a^2 B_2^2 h_0^2 - 15B_2^3 - 120a^4 B_2 h_0^4) = 0. \end{aligned}$$

where f_k ($k = 0, \pm 1, \pm 2, \pm 3, \pm 4, \pm 5,$) are nonlinear function. Choose $h_1 = 0$ and $h_2 = -h_0$, solving the obtained algebraic equations by use of Mathematica 10.1, we obtain

$$A_0 = 2(\alpha h_0)^2, A_1 = 0, A_2 = -2(\alpha h_0)^2, B_1 = 0, B_2 = 0, \quad (3.5)$$

where $c = \sqrt{1 + 16(\alpha h_0)^4}$ and

$$\begin{aligned} A_0 & = (\alpha h_0)^2 \left(1 - \frac{1}{15} \sqrt{105} \right), A_1 = 0, A_2 = -2(\alpha h_0)^2, \\ B_1 & = 0, B_2 = 0, \end{aligned} \quad (3.6)$$

where $c = \sqrt{1 + 2\alpha^2 h_0^4 (11 + \sqrt{105})}$.

Substituting equation (3.5) and equation (3.6) into equation (3.4), we obtain two exact solutions for equation (3.1) in the form

$$u(x, t) = 2(\alpha h_0)^2 \left(1 - \tanh^2(h_0 \alpha (x - \sqrt{1 + 16(\alpha h_0)^4} t)) \right) \quad (3.7)$$

and

$$u(x, t) = (\alpha h_0)^2 \left(1 - \frac{1}{15} \sqrt{105} \right)$$

$$-2(\alpha h_0)^2 \tanh^2 \left(h_0 \alpha \left(x - \sqrt{1 + 2\alpha^2 h_0^4 (11 + \sqrt{105}) t} \right) \right). \quad (3.8)$$

Next, choose $h_2 = 0$ and $h_1 = -h_0$, solving the obtained algebraic equations by use of Mathematica 10.1, we obtain

$$A_0 = 0, A_1 = 0, A_2 = 0, B_1 = 2(\alpha h_0)^2, B_2 = -2(\alpha h_0)^2, \quad (3.9)$$

where $c = \sqrt{1 + (\alpha h_0)^4}$ and

$$A_0 = (\alpha h_0)^2 \left(-\frac{1}{4} + \frac{1}{60} \sqrt{105} \right), A_1 = 0, A_2 = 0, \quad (3.10)$$

$$B_1 = 2(\alpha h_0)^2, B_2 = -2(\alpha h_0)^2,$$

where $c = \sqrt{1 + \frac{1}{8} \alpha^4 h_0^4 (11 - \sqrt{105})}$.

Substituting equation (3.5) and equation (3.6) into equation (3.4), we obtain two exact solutions for equation (3.1) in the form

$$u(x, t) = \frac{2(\alpha h_0)^2}{1 - \frac{1}{h_0} \exp \left(-h_0 \alpha (x - \sqrt{1 + (\alpha h_0)^4 t}) \right)} - \frac{2(\alpha h_0)^2}{\left(1 - \frac{1}{h_0} \exp \left(-h_0 \alpha (x - \sqrt{1 + (\alpha h_0)^4 t}) \right) \right)^2} \quad (3.11)$$

and

$$u(x, t) = (\alpha h_0)^2 \left(-\frac{1}{4} + \frac{1}{60} \sqrt{105} \right) + \frac{2(\alpha h_0)^2}{1 - \frac{1}{h_0} \exp \left(-h_0 \alpha (x - \sqrt{1 + \frac{1}{8} \alpha^4 h_0^4 (11 - \sqrt{105}) t}) \right)} - \frac{2(\alpha h_0)^2}{\left(1 - \frac{1}{h_0} \exp \left(-h_0 \alpha (x - \sqrt{1 + \frac{1}{8} \alpha^4 h_0^4 (11 - \sqrt{105}) t}) \right) \right)^2} \quad (3.12)$$

The correctness of the solution is verified by substituting them into original equation (3.1). Our solutions in (3.7), (3.8), (3.11), and (3.12) are coincided with the published results for a special case which are gained by Hosseini et.al. [4]. For direct-viewing analysis, we provide the figures of $u(x, t)$ in equation (3.7) and (3.11), where we choose $\alpha = 1$ and $h_0 = -1$.




Figure 1: The graph of function (3.7) which was derived from equation (3.1) (right), and its profile graph for t_0 .




Figure 2: The graph of function (3.11) which was derived from equation (3.1) (right), and its profile graph for t_0 .

4. Conclusions

In this paper, modified F-expansion method has been successfully applied to find the some exact traveling wave solutions of the sixth-order Boussinesq equation. The method is more effective and simple than other methods and a number of solutions can be obtained at the same time. The obtained solutions are presented through the hyperbolic functions and the exponential functions. The obtained results are verified by putting them back into the original equation. Some of our solutions are in good agreement with already published results for a special case. This study shows that the modified F-expansion method is quite efficient and practically well suited for use in finding exact solutions for the nonlinear evolution equation in mathematical physics and engineering sciences.

References

- [1] Cai, G., Q. Wang, and J. Huang, 2006, “A modified F-expansion method for solving breaking soliton equation”, *Int. Journal of Nonlinear Science*, 2(2), 122–128.
- [2] Clarkson, P.A., 1990, “New exact solutions of the Boussinesq equation”, *European Journal of Applied Mathematics*, 1(3), 279–300. DOI: <http://dx.doi.org/10.1017/S095679250000022X>
- [3] Daripa, P., and W. Hua, 1999, “A numerical study of an ill-posed Boussinesq equation arising in water waves and nonlinear lattices: Filtering and regularization techniques”, *Applied Mathematics and Computation*, 101(2-3), 159–207.
- [4] Hosseini, M., H. Abdollahzadeh, and M. Abdollahzadeh, 2011, “Exact travelling solutions for the sixth-order Boussinesq equation”, *The Journal of Mathematics and Computer Science*, 2(2), 376–387.
- [5] Lai, S., Y.H. We, Y. Zhou, 2008, “Some physical structures for the (2+1)-dimensional Boussinesq water equation with positive and negative exponents”, *Comput. Math. Appl*, 56(2), 339–345. DOI:<http://dx.doi.org/10.1016/j.camwa.2007.12.013>
- [6] Naher, H., F.A. Abdullah, and M.A. Akbar, 2011, “The exp-function method for new exact solutions of the nonlinear partial differential equations”, *International Journal of the Physical Sciences*, 6(29), 6706–6716.
- [7] Nagasawa, T., Y. Nishida, 1992, “Mechanism of resonant interaction of plane ion acoustic solitons”, *Phys. Rev. A*, 46(3471), 3471–3476.
- [8] Wazwaz, A.M., 2008, “Multiple-soliton solutions for the ninth-order KdV equation and sixth-order Boussinesq equation”, *Applied Mathematics and Computation*, 203(1), 277–283. DOI:<http://dx.doi.org/10.1016/j.amc.2008.04.040>
- [9] Wang, M., and X.Li, 2005, “Applications of F-expansion to periodic wave solutions for a new Hamiltonian amplitude equations”, *Chaos, Solitons and Fractals*, 24(5), 1257–1268. DOI:<http://dx.doi.org/10.1016/j.chaos.2004.09.044>
- [10] Zakharov, V.E., 1974, “On stochastization of one dimensional chains of nonlinear oscillations”, *Sov. Phys. JETP*, 38(1), 108–110.