

## Nutritional Properties of Three Different Origins Of Indonesian *Jatropha (Jatropha Curcas)* Meal For Ruminant

Lydia Gandini Triastuty<sup>1)</sup>, Despal<sup>1)</sup> and Idat Galih Permana<sup>1)</sup>

### Abstract

An experiment to explore the nutrition properties of Indonesian *Jatropha curcas* meal for ruminant has been conducted. Three different origins of Indonesian *jatropha* (Lampung, Kebumen and Lombok) have been investigated for their nutritional values such as chemical compositions, amino acid profiles, fermentability and in vitro digestibility. Anti nutritional and toxic compounds are also determined. It was found that chemical compositions and amino acid profiles of the *Jatropha* meal vary according to the origin of *Jatropha*. The *jatropha* meal from Lampung had the highest CP (% DM) content (42.58%) compared with *jatropha* meal from another region (Kebumen 37.93%, Lombok 32.94%). The CP (%Solid Non Fat) content of *jatropha* meal from Lampung (58.59%) was higher than CP content of soybean meal (50.71%). The *jatropha* meal from three different regions in Indonesia showed a significant difference ( $p < 0.01$ ) in IVDMD and IVOMD, which is the *jatropha* meal from Lampung (60.23%) and Kebumen (60.73%) was higher than from Lombok (49.8%). Toxic and anti-nutritional factors that studied in this research are curcin, phorbol ester and phytate. The phytate and curcin level was highest in *jatropha* meal from Lampung. But the level of phorbol ester was not detectable in this meal. The contents of toxic and anti-nutritional factors did not affect on digestibility and ruminal fermentation products.

1). Department of Animal Nutrition and Feed Technology  
Faculty of Animal Science, Bogor Agricultural University-Indonesia  
Jl. Agatis, Kampus IPB Darmaga, Bogor, Email: permana@ipb.ac.id

### Introduction

*Jatropha curcas* L. is known as a prospective renewable energy source which to be developed in Indonesia. *Jatropha curcas* represent annual crop which hold up dryness so that able to grow and expand better in marginal land as in region of East Indonesia (Hambali, 2006).

The oil from the kernels of *Jatropha curcas* came through processing phase. *Jatropha* seed extraction can be conducted by using simple expeller machine. *Jatropha* meal is the residue from seed processing. The extraction will obtained 30-40% oil and 60% *jatropha* meal. The residue in the form of *jatropha* meal has potential as animal feed. However, due to the toxicity further research need to be conducted to have the basic information from *jatropha* meal.

The objective of this research were to get the basic information of *jatropha* meal include the information of the nutrition, anti nutrition and toxic factors which is the main constraint of *Jatropha curcas*, and to study the characteristics of *jatropha* meal as animal feed especially for ruminant.

### Materials and Methods

#### Sample Preparation

The *jatropha* seeds used were from three different regions from Indonesia (Lampung, Kebumen and Lombok). The seed were dehulled, dried and ground manually. The kernel was defatted using manual hydraulic press machine. The defatted and ground kernel is referred to as the meal and used for next analysis. Soybean meal was used for some analysis as comparison.

#### Chemical Analysis

Chemical compositions of *jatropha* meal were determined according to AOAC (1980) procedure. The amino acids analysis was conducted using HPLC, while the gross energy was measured using Bomb Calorimeter. The concentration of phorbol ester of *jatropha* meal was carried out by de procedures describes by Makkar *et al.* (1997), while lectin/cursin was carried out by heamagglutination assay (Aregheore *et al.* 1998). The concentration of phytate was determined using spectrophotometry procedure.