



# International Society for Horticultural Science Société Internationale de la Science Horticole

ACTA HORTICULTURAE® is a peer reviewed publication of ISHS. Information about Acta Horticulturae® and ISHS is given at the end of this book.

Check out www.actahort.org for details on our latest titles.

# Editorial Advisory Board of Acta Horticulturae®

Jules Janick, Purdue University, USA, Chair of the Editorial Advisory Board

Isabel Ferreira, Instituto Superior de Agronomia, Lisbon, Portugal

Kim Hummer, USDA ARS NCGR, Corvallis, USA

Hilde Nybom, Balsgard-Dept. Crop Science, Swedish University of Agricultural Sciences, Kristianstad, Sweden

Robert K. Prange, Agriculture and Agri-Food Canada, Kentville, Canada

Anthony David Webster, Malmesbury, Wiltshire, United Kingdom

**Executive Director of ISHS** Ir. J. Van Assche

Secretariat of ISHS

PO Box 500 3001 Leuven 1 Belaium

Phone:

+32.16.22 94 27

Fax: E-mail: Internet: www.ishs.org

+32.16.22 94 50 info@ishs.org

X

# PROCEEDINGS OF THE INTERNATIONAL ISHS-PROMUSA SYMPOSIUM ON GLOBAL PERSPECTIVES ON ASIAN CHALLENGES

Co-Conveners

Jiang Zongyong Yi Ganjun A. Molina I. Van den Bergh

Guangzhou, China

September 14-18, 2009

ISHS Section Banana and Plantain ISHS Working Group on Musa Crop Improvement ISHS Working Group on Musa Crop Production ISHS Working Group on Musa Crop Protection ISSN 0567-7572 ISBN 978 90 6605 138 6, Acta Horticulturae n°. 897 Price for non-members of ISHS: € 112,-Published by ISHS, May 2011

Executive Director of ISHS: J. Van Assche Technical Processing: M.L.M. Testor

ISHS Secretariat, PO Box 500, 3001 Leuven 1, Belgium

Printed by Drukkerij Geers Eeckhoutdriesstraat 67, 9041 Gent (Oostakker), Belgium

<sup>©</sup>2011 by the International Society for Horticultural Science (ISHS). All rights reserved. No part of this book may be reproduced and/or published in any form or by any means, electronic or mechanical, including photocopying, microfilm and recording, or by any information storage and retrieval system, without written permission from the publishers.

Photographs on the front cover:

1	2
3	4
5	6

- 1. Banana fields on the island of Hainan. Photograph by courtesy of Nicolas Roux.
- Visit to the Maosheng orchard, Shenwan Town, Zhongshan City. Photograph by courtesy of Inge Van den Bergh.
- 3. Banana at the South China Botanical Garden. Photograph by courtesy of Inge Van den Bergh.
- 4. Musella lasiocarpa at the South China Botanical Garden. Photograph by courtesy of Inge Van den Bergh.
- 5. Musella lasiocarpa at the South China Botanical Garden. Photograph by courtesy of Inge Van den Bergh.
- 6. Banana at the South China Botanical Garden. Photograph by courtesy of Inge Van den Bergh.

# Editors

I. Van den Bergh

M. Smith

R. Swennen

C. Hermanto

# **Editorial Board**

N. Roux

Huang Bingzhi

P. Heslop-Harrison

R. Miller

A. Molina

A. Churchill

M. Dita

# The International ISHS-ProMusa Symposium on Global Perspectives on Asian Challenges

#### Co-Conveners

Jiang Zongyong Yi Ganjun A. Molina I. Van den Bergh

### Scientific Committee

Chairs

N. Roux

Yi Ganjun

Members

Huang Bingzhi

M. Smith

R. Swennen

Wei Yuerong C. Hermanto

Li Chunyu

A. Molina

A. Churchill

Z. de Beer

Huang Xuelin

I. Van den Bergh

V. Galán Saúco

H.P. Singh

Jin Zhiqiang

# **Organising Committee**

Chairs

Jiang Zongyong

Yi Ganjun

A. Molina

I. Van den Bergh

Members

Bai Xianjin

Lin Jianliang

Zhang Zongwen

Chen Weixin

Peng Hongxiang

Zhang Shaoxing

Liu Guanghua

Wei Yuerong Huang Bingzhi

K. Lehrer

Chen Daoming

Qin Suijia

# **Sponsors and Organisers**











#### PREFACE

The papers contained in this volume of *Acta Horticulturae* report the peer reviewed Proceedings of the International ISHS-ProMusa Symposium on Global Perspectives on Asian Challenges. Keynote speakers and authors of selected contributed oral and poster presentations were given the opportunity to submit a manuscript for publication.

The manuscripts were reviewed by the Editors and members of the Editorial Board. Only those papers judged suitable for publication following the authors' consideration of

reviewer suggestions appear in this volume of Acta Horticulturae.

The ISHS acknowledges and appreciates the contribution of all editors and reviewers. They have made a significant contribution to improving the quality of this publication.

The ISHS Board of Directors

# LIST OF CONTENTS

Preface List of Contents List of Authors List of Participants	5 7 13 17
Opening Session	
Status, Challenges and Trends of Chinese Banana Industry C.Y. Li, Y.R. Wei, Y.L. Wu, B.Z. Huang and G.J. Yi	31
Novel Approaches to Understanding, Conserving and Using Banana Genetic Diversity	
Non-Conventional Breeding of Banana ( <i>Musa</i> spp.) Y.F. Chen, X.M. Dai, Q. Gong, X. Huang, W. Xiao, J.T. Zhao and X.L. Huang	39
From Fundamental Research Discoveries to Applications for Banana	47
Improvement R. Swennen, S.C. Carpentier, I.M. Henry, A. Vertommen, I. Van den houwe, G. Kovacs, L. Sagi, S. Remy and B. Panis	47
Genomics, Banana Breeding and Superdomestication J.S. Heslop-Harrison	55
Different Approaches to Discover the Repetitive Part of the Banana Nuclear Genome <i>E. Hřibová and J. Doležel</i>	63
Identification and Validation of EST-Derived Molecular Markers, TRAP and VNTRs, for Banana Research S.A.L. Garcia, R. Talebi, C.F. Ferreira, B.I. Vroh, L.V. Paiva, G.H.J. Kema and M.T. Souza Jr.	69
Wild Musaceae Species in China M. Häkkinen	81
Musa Collection and Characterisation in Central and Eastern DR Congo: a Chronological Overview B. Dheda Djailo, B.D. Nzawele, N. Roux, F. Ngezahayo, N. Vigheri, E. De Langhe, D. Karamura, C. Picq, P. Mobambo, R. Swennen and G. Blomme	87
Morphological Characterisation of East African AAB and AA Dessert Bananas ( <i>Musa</i> spp.)  M. Onyango, D. Karamura, S. Keeley, R. Manshardt and D. Haymer	95
The Complementarity of Farmers' and Botanical Descriptors of the East African Highland Banana Cultivars ( <i>Musa</i> , AAA) D. Karamura, A. Kiggundu and E. Karamura	107

Standardised Procedure for <i>Musa</i> Germplasm Characterisation S. Channelière, I. Van den houwe, E. Arnaud, J.P. Horry, M. Ruas and N. Roux	113
Diversity Analysis in Indian Cooking Bananas ( <i>Musa</i> , ABB) through Morphotaxonomic and Molecular Characterisation <i>M.S. Saraswathi, S. Uma, E. Vadivel, P. Durai, S.A. Siva, G. Rajagopal and S. Sathiamoorthy</i>	123
Application of Cytoplasmic and Nuclear DNA-Based Marker Systems for Elucidation of the Phylogenetic Relationship of <i>Musa acuminata</i> and <i>Musa balbisiana</i> and Their Hybrids <i>R. Boonruangrod, D. Desai, M. Berenyi, S. Fluch and K. Burg</i>	133
A Complex Approach for Unravelling <i>Musaceae</i> Phylogeny at Molecular Level <i>P. Němcová, E. Hřibová, M. Valárik and J. Doležel</i>	139
Molecular Analysis Reveals Multiple Domestications of Edible Bananas H. Volkaert	143
Microsatellite Markers for Classifying and Analysing Genetic Relationship between Banana Cultivars in Indonesia A. Retnoningsih, R. Megia and A. Hartana	153
Validation of Rapid (Colour-Based) Prescreening Techniques for Analysis of Fruit Provitamin A Contents in Banana ( <i>Musa</i> spp.)  L.C. Pereira, G.B. Ngoh Newilah, M.W. Davey and I. Van den Bergh	161
Vitamin A Biofortification in <i>Musa</i> : Status, Bottlenecks and Prospects <i>M.W. Davey, I. Van den Bergh and N. Roux</i>	169
Candidate Resistance Gene Discovery: Resistance Gene Analog Characterisation and Differential Gene Expression Analysis in <i>Musa-Mycosphaerella</i> Host-Pathogen Interactions <i>R.N.G. Miller, M.A.N. Passos, F.L. Emediato, C. de Camargo Teixeira and G.J. Pappas Júnior</i>	179
MA-ACS1: a Key Operator in Ethylene Biosynthesis in Banana - Its Role and Regulation during Fruit Ripening S. Roy Choudhury, S. Roy, S.K. Singh and D.N. Sengupta	187
Combination of Suppression-Subtractive Hybridisation with cDNA Microarray, a Novel Way to Identify Genes from Banana Involved in Fruit Ripening Z.Q. Jin, B.Y. Xu, J.H. Liu, J.B. Zhang and C.H. Jia	195
Developing Resistant Banana and Plantain Cultivars through Conventional Breeding Techniques R. Menon, A. Cherian, A. Suma, Maicykutty, P. Mathew, S. Nair and K.C. Aipe	207
Exploitation of Diploids in Indian Banana Breeding Programmes S. Uma, M.M. Mustaffa, M.S. Saraswathi and P. Durai	215

The Effect of Banana Breeding on the Diversity of East African Highland Banana ( <i>Musa</i> , AAA)  M. Nyine and M. Pillay	225
Use of Molecular Markers in Banana and Plantain Improvement J. Lorenzen, S. Hearne, G. Mbanjo, M. Nyine and T. Close	231
The Use of Molecular Markers in the CIRAD Banana Breeding Programme <i>J.P. Horry</i>	237
Development of Highly Regenerative Embryogenic Cell Suspensions of 'Nanjangud Rasbale' (syn. 'Rasthali', <i>Musa</i> , AAB, Silk Subgroup) and Transformants with <i>AMP</i> Gene S. Mohandas, H.D. Sowmya, R. Manjula, K.Y. Pratibha, S. Meenakshi and K.M. Ajay	245
Collecting and Conserving Wild <i>Musa</i> Germplasm in Guangxi, China <i>J.Y. Yan, X.Q. Quan, X. Long and H.X. Peng</i>	255
Preliminary Study on ISSR Analysis and Classification of Wild <i>Musa</i> Germplasm in Guangxi, China <i>X.Q. Qin, H.X. Peng, X. Long and J.Y. Yao</i>	259
Analysis of Genetic Diversity in Banana Cultivars ( <i>Musa</i> spp.) Using Sequence-Related Amplified Polymorphism Markers <i>J.Y. Wei, D.B. Liu, S.X. Wei, Z.S. Xie, G.Y. Xu and Y.Y. Chen</i>	263
Intersectional Relationship between <i>Eumusa</i> and <i>Rhodochlamys</i> of the Genus <i>Musa</i> Using Morphotaxonomy and Microsatellite Markers <i>P. Durai, S. Uma, M.S. Saraswathi, N. Jayabalan and M.M. Mustaffa</i>	267
Studies on Genetic Divergence of Seeded Banana in West Bengal M.A. Hasan, R. Ray Chowdhury, M. Manna, K.K. Mandal, D. Majumder and S. Jha	271
Generation of Segregating F <sub>1</sub> Populations for Mapping the <i>Musa</i> Genome A. Rekha, K.V. Ravishankar and D.S. Ambika	277
Gene Expression Analysis in Leaves of 'Bee Hee Kela', a Drought-Tolerant <i>Musa balbisiana</i> Genotype from Northeast India K.V. Ravishankar, A. Rekha, R.H. Laxman, G. Savitha and V. Swarupa	279
Studies on Time-Course Expression of Defence Genes in Banana against <i>Pratylenchus coffeae</i> for the Creation of a Subtractive cDNA Library <i>S. Backiyarani, S. Uma, P. Sundararaju, M. Mayilvaganan, M.S. Saraswathi and S. Jeeva</i>	281
Seed as an Alternative Source of DNA for Molecular Research of Inaccessible Wild Musa Species S. Uma, M.S. Saraswathi and D. Anto	285
Using an Improved Site-Directed Mutagenesis for the Creation of Constructs to Analyse Gene Function in <i>Fusarium oxysporum</i> f. sp. <i>cubense C.Y. Li, G.J. Yi, S. Chen, Q.M. Sun, C.W. Zuo, B.Z. Huang, Y.R. Wei, Y.H. Huang, Y.L. Wu, L.B. Xu and C.H. Hu</i>	289

Transformation of the Banana Cultivar 'Nanjangud Rasbale' (Syn. 'Rasthali', AAB, Silk Subgroup) with the <i>AMP</i> Gene and Screening for Fusarium Resistance with a Bioassay S. Mohandas, R. Manjula, A.K. Saxena, K.M. Ajay, B. Shakunthala, H.D. Sowmya and S. Meenakshi	293
Evaluation and Tissue-Culture Conservation, Multiplication and Distribution of Rare and Carotenoid-Rich Fe'i Banana Cultivars in Micronesia <i>V.M. Verma</i>	297
Micropropagation of Malaysian Banana Cultivars ( <i>Musa</i> spp.) Using a Modified Twin Flask System <i>L.K. Chan and V.H. Au</i>	299
The Effect of Ethyl Methane Sulphomate (EMS) and Sodium Azide (NaN <sub>3</sub> ) on Plant Regeneration Capacity of an Embryogenic Cell Suspension of 'Yueyoukang 1' ( <i>Musa</i> , AAA), a Banana Cultivar Resistant to Fusarium Wilt <i>C.X. Xu, J. Xiao, J.G. He, G.B. Hu and H.B. Chen</i>	301
Integrated Approaches to Managing Fusarium Wilt and Other Emerging Disease Threats	
Early Events in the <i>Fusarium oxysporum</i> f. sp. <i>cubense - Musa</i> Interaction Using Green Fluorescent Protein-Tagged Race 4 <i>C.Y. Li, G.J. Yi, B.Z. Huang and Y.R. Wei</i>	305
Incidence and Distribution of Fusarium Wilt Disease of Banana in Indonesia C. Hermanto, A. Sutanto, Jumjunidang, H.S. Edison, J.W. Daniells, W.T. O'Neill, V.G.O. Sinohin, A.B. Molina and P. Taylor	313
Status of Fusarium Wilt in India M.M. Mustaffa and R. Thangavelu	323
Raising Awareness of the Threat of Fusarium Wilt Tropical Race 4 in Latin America and the Caribbean L.E. Pocasangre, R.C. Ploetz, A.B. Molina and L. Perez Vicente	331
A Detection Method for Tropical Race 4 of the Banana Pathogen <i>Fusarium oxysporum</i> f. sp. <i>cubense</i> C. Waalwijk, M.A. Dita, I. Buddenhagen, L.V. Paiva, M.T. Souza Jr. and G.H.J. Kema	339
Vegetative Compatibility Group Analysis of Indonesian <i>Fusarium oxysporum</i> f. sp. <i>cubense</i> Isolates <i>W.T. O'Neill, A.B. Pattison, J.W. Daniells, C. Hermanto and A.B. Molina</i>	345
Cross Infection and Genetic Diversity of <i>Fusarium oxysporum</i> f. sp. <i>cubense</i> , the Causal Agent of Fusarium Wilt in Banana <i>R. Thangavelu, P. Suganya Devi, P.M. Chrismala and M.M. Mustaffa</i>	353
Gene Expression Analysis in Roots of <i>Musa acuminata</i> ssp. <i>burmannicoides</i> 'Calcutta-4', a Banana Genotype Tolerant to Fusarium Wilt	363

Characterisation of Early Events in Banana Roots Infected with Green Fluorescent Protein-Tagged Fusarium oxysporum f. sp. cubense X.M. Yin, B.Y. Xu, W. Zheng, Z. Wang, B.Z. Wang, W.H. Ma, H.T. Mao, J.Y. Li, Z.W. Sheng, D.X. Wang, Y.D. He, L.N. Han, S.W. Chen, J.B. Zhang, Y.G. Fu, J.B. Wang and Z.Q. Jin	371
A Greenhouse Bioassay for the <i>Fusarium oxysporum</i> f. sp. <i>cubense</i> x 'Grand Naine' ( <i>Musa</i> , AAA, Cavendish Subgroup) Interaction <i>M.A. Dita, C. Waalwijk, L.V. Paiva, M.T. Souza Jr. and G.H.J. Kema</i>	377
Discrimination of Banana Genotypes for Fusarium Wilt Resistance in the Greenhouse L.R. Ribeiro, E.P. Amorim, Z.J.M. Cordeiro, S. de Oliveira e Silva and M.A. Dita	381
Reaction of Diploid (AA) and Tetraploid (AAAB) Banana Hybrids to Fusarium Wilt under Field Conditions A.P. de Matos, Z.J.M. Cordeiro, S. de Oliveira e Silva, E.P. Amorim and D.M.V. Ferreira	387
Selection of a New Somaclone Cultivar 'Tai-Chiao No. 5' (AAA, Cavendish) with Resistance to Fusarium Wilt of Banana in Chinese Taipei S.Y. Lee, Y.U. Su, C.S. Chou, C.C. Liu, C.C. Chen and C.P. Chao	391
Comparison of Host Reaction to <i>Fusarium oxysporum</i> f. sp. <i>cubense</i> Tropical Race 4 and Agronomic Performance of Somaclonal Variant 'GCTCV-119' (AAA, Cavendish) and 'Grand Naine' (AAA, Cavendish) in Commercial Farms in the Philippines <i>A.B. Molina, E.G. Fabregar, R.O. Soquita and V.G.O. Sinohin</i>	399
Combating Banana Wilts – What Do Resistant Cultivars Have to Offer? J.W. Daniells	403
Fusarium Wilt Incidence, Growth, Yield and Post-Harvest Quality of Banana as Affected by Organic Farming in Chinese Taipei C.M. Chang, C.P. Chao, S.N. Huang and S.C. Chiang	413
Efficacy of Clay-Based Formulated <i>Serratia</i> in Reducing Inoculum of <i>Fusarium oxysporum</i> f. sp. <i>cubense</i> Tropical Race 4 A.S.Y. Ting, M.T. Fang and C.S. Tee	421
In Vitro Mutation Induction for the Improvement of <i>Mycosphaerella fijiensis</i> Resistance in Two Banana Cultivars Grown in Kenya <i>M. Mwangi, S.W. Waudo and E.M. Kahangi</i>	427
Screening <i>Musa</i> Genotypes for Banana Bunchy Top Disease Resistance in Burundi <i>C. Niyongere, E. Ateka, T. Losenge, G. Blomme and P. Lepoint</i>	439
Transgenic Banana Plants Resistant to <i>Banana Bunchy Top Virus</i> Infection W. Borth, E. Perez, K. Cheah, Y. Chen, W.S. Xie, D. Gaskill, S. Khalil, D. Sether, M. Melzer, M. Wang, R. Manshardt, D. Gonsalves and J.S. Hu	449

Enhancing Banana Crop Management by Use of Tissue-Culture Derived Planting Material in Kenya J. Njuguna, S. Nguthi, F. Wambugu, D. Gitau and M. Karuoya	459
Virus-Indexing Technology for Production of Quality Banana Planting Material: a Boon to the Tissue-Culture Industry and Banana Growers in India R. Selvarajan, V. Balasubramanian, M.M. Sheeba, R. Raj Mohan and M.M. Mustaffa	463
Monitoring Fusarium Wilt in Hainan Province and Preliminary Pathogen Characterisation X. Zhang, H. Zhang, Y. Xie, J. Pu, Y. Qi and Y. Lu	471
Banana Varieties and <i>Fusarium oxysporum</i> f. sp. <i>cubense</i> in Indonesia – Observations from Fusarium Wilt Disease Databases <i>J.W. Daniells, W.T. O'Neill, C. Hermanto and R.C. Ploetz</i>	475
Evaluation of Seeded Banana Types for Biotic Stress M.A. Hasan, R. Ray Chowdhury, M. Manna, K.K. Mandal, D. Majumder and S. Jha	479
Importance of Black Leaf Streak Control in 'Pisang Berangan' (AAA) in Peninsular Malaysia <i>K.H. Then and S. Palaniappan</i>	483
CR-M: an Important Nucleotide Acid Sequence to Distinguish <i>Banana Bunchy Top Virus</i> Isolates from Two Groups <i>Y.H. Huang, J.W. Zeng, Y.L. Wu, R. Xia and G.J. Yi</i>	485
Investigation into Low-Cost Medium for Hardening of In Vitro Banana Plantlets to Promote Adoption of Disease-Free Plants <i>B. Jhurree-Dussoruth and H. Kallydin</i>	489
Micropropagation of 'Pisang Awak' ( <i>Musa</i> , ABB Genome) as a Model for Improving Livelihoods of Rural Communities in Malaysia <i>L.K. Chan, V.H. Au, A.A.A. Noor, M. Adnan and P.L. Boey</i>	491
Closing Session	
Harnessing the Potential of Banana and Plantain in Asia and the Pacific for Inclusive Growth <i>H.P. Singh</i>	495
The Role of Genomics in Banana Improvement	
Opportunities for Bridging the Gap between Genomics and Genetic Improvement in <i>Musa</i> spp.  N. Roux, M. Rouard, X.L. Huang and M. Smith	509
Mitigating the Threat of Fusarium Wilt	
Fusarium Wilt: the Banana Disease that Refuses to Go Away R.C. Ploetz and A.C.L. Churchill	519

# Microsatellite Markers for Classifying and Analysing Genetic Relationship between Banana Cultivars in Indonesia

A. Retnoningsih<sup>1, 2</sup>, R. Megia<sup>1</sup> and A. Hartana<sup>1</sup>

Department of Biology, Faculty of Mathematics and Natural Sciences, Bogor Agricultural University, Indonesia

Agricultural University, Indonesia

<sup>2</sup>Current address: Department of Biology, Faculty of Mathematics and Natural Sciences,

Semarang State University, Indonesia

Keywords: Musa acuminata, Musa balbisiana, SSR

#### Abstract

In this study, microsatellite markers were used to identify Musa genomic groups and to investigate genetic relationships among banana accessions from various areas in Indonesia. One hundred sixteen banana accessions were analysed using MaCIR108 and Ma-3-90 primer pairs for identifying genomic groups. Six additional SSR primer pairs were used for genetic relationship analysis. The results showed that 73 accessions should be classified in the AA/AAA and AAA genomic groups, two accessions in the BB genomic group, 21 accessions in the AAB genomic group and 20 accessions in the ABB genomic group. Ninety-nine out of the 116 accessions were unique genotypes while the rest were synonyms. The dendrogram generated by UPGMA analysis separated the 116 banana accessions into two main clusters with a similarity of 0.13. All banana accessions belonging to the BB, ABB and AAB genomic groups clustered in the first main cluster, together with the majority of the accessions containing the pure A genome. The second main cluster was formed of 11 accessions of AA/AAA and AAA genome. Within the first main cluster, the accessions containing the B genome were clustered according to their genomic group, except four AAB accessions clustering with accessions containing the A genome alone. The ABB genomic group appeared closer to the BB than to the AAB genomic group. The AA and AAA banana accessions could not be significantly distinguished, although the majority of accessions tended to be clustered according to their ploidy level.

#### INTRODUCTION

Most edible bananas originated from two wild species, *Musa acuminata* Colla (2n) and *Musa balbisiana* Colla (2n). Scientifically, banana cultivars are often referred to by their genomic groups, which are categorised on the basis of their ploidy levels and the genomes they contain. Ploidy in bananas includes diploids, triploids and rare tetraploids, resulting in genomic groups: AA, BB, AAA, AAB, ABB, AAAB ... with the letters A and B representing the contribution of *M. acuminata* and *M. balbisiana*, respectively. The hybrid bananas that evolved from the two natural species mostly formed triploid cultivars (Heslop-Harison and Schwarzacher, 2007).

Genomic constitution plays an important role in the classification of bananas (Pillay et al., 2004). Cultivar identification based on morphological characters is not always easy due to the influence of environmental factors. Molecular approaches have been recognised to be more effective than morphological techniques to distinguish banana

cultivars and classify them in genomic groups (Ford-Lloyd et al., 1997).

Microsatellites are one of the most informative molecular markers to reveal genetic diversity of banana cultivars (Creste et al., 2003, 2004). Such markers have been used for genotype identification of many plant species (Perera et al., 2001; Chakravarthi and Naravaneni, 2006). Microsatellites are DNA regions that consist of short repeat units flanked by conserved sequences that are unique to specific loci in the genome (Semagn et al., 2006). In previous studies in banana, several microsatellite primers were shown to be useful in producing diagnostic characters for different genomic groups (Kaemmer et al., 1997; Creste et al., 2005), with a determination key based on application of these markers

constructed (Retnoningsih et al., 2010a).

The objectives of this study were to classify banana accessions from various regions in Indonesia using a microsatellite determination key, and to investigate the genetic relationships among these accessions based on microsatellite markers.

#### MATERIALS AND METHODS

#### Plant Materials and Primers

A total of 116 banana accessions were collected from various locations in Indonesia. The genetic relationships among banana germplasm was investigated using eight microsatellite primer pairs. The accessions were classified into their genomic groups using a molecular key developed by Retnoningsih et al. (2010a), presented in Figure 1.

DNA Extraction, Microsatellite Region Amplification and Silver Staining

Total DNA was extracted from fresh young leaves using a modified SDS procedure (Dixit, 1998). Amplification of microsatellite regions was conducted following Kaemmer et al. (1997), with modifications, using a Perkin Elmer 2400 thermocycler (Applied Biosystems, Foster City, CA, USA®). The PCR products were separated on 6% denaturing polyacrylamide gels containing 7 M urea (Sigma-Aldrich Chemicals Germany®) at 45-60 W for 2 to 3 h. Alleles of each microsatellite were visualised by silver staining according to the modified procedure of Creste et al. (2001). The size of each allele was estimated using a 100 bp DNA ladder (Invitrogen®).

**Data Analysis** 

Each variant fragment of microsatellite was considered an allele. Genetic relationship analysis was based on data generated with the eight microsatellite markers, with each allele scored as present (1) or absent (0). Jaccard's coefficient was used to calculate similarity values between accessions using the Similarity of Qualitative Data (SYMQUAL) procedure. This matrix was then analysed via cluster analysis to visualise genetic relationships among the accessions. Cluster analysis was performed using Sequential, Agglomerative, Hierarchical and Nested (SAHN) clustering of the Unweighted Pair-grouping with Arithmetic Average (UPGMA) using Numerical Taxonomy and Multivariate Analysis System (NTSYS) software, version 2.02 for PC (Rohlf, 1998).

#### RESULTS AND DISCUSSION

All microsatellite markers generated clearly distinguishable alleles and polymorphic patterns. A total of 93 alleles were detected with the eight markers. The number of alleles per locus varied from 5 to 18, with an average of 11.6 alleles per locus. The size of the alleles ranged from approximately 110 to 436 bp with the polymorphism degree per primer ranging from 5 to 18 alleles. The highest degree of polymorphism was observed with MaCIR108 and the lowest was detected with MaCIR327b (Table 1).

Classification of Banana Accessions Using a Microsatellite Determination Key

The molecular determination key used was able to classify the 116 banana accessions effectively into their genomic groups. The key is based on four diagnostic alleles of locus MaCIR108 and one allele of locus Ma-3-90 with a size of 152 bp. A MaCIR108 allele with a size longer than 270 bp was considered a specific allele for the B genome, while an allele equal to or less than 270 bp was considered specific for the A genome (Retnoningsih et al., 2010b).

Seventy-three banana accessions were classified in the AAA or AA/AAA genomic groups, because they possessed only alleles of MaCIR108 with size equal to or shorter than 270 bp. The size of the MaCIR108 allele observed in these accessions ranged from approximately 210 to 268 bp. A-genome accessions with only one or two alleles were grouped in the AA/AAA genomic group whereas the accessions having three alleles were

classified into AAA genomic group. Diploid accessions could not be easily distinguished from triploid accessions, because there are no specific alleles to differentiate the AA from the AAA genomic group, as also reported by Creste et al. (2004). Furthermore, the dosage effects of the microsatellite allele (simplex, duplex, triplex, etc.) cannot be differentiated (Zhang et al., 2000) because most cultivated bananas are naturally polyploid (Stover and Simmonds, 1987). The triploid accessions possessing two alleles in a certain locus could not be differentiated from the diploid also possessing the same allele numbers. Two accessions were classified in the BB genomic group, because they possessed a MaCIR108

allele with a size of 295 bp without any allele size shorter than 270 bp.

Twenty accessions were classified in the ABB genomic group and 21 accessions in the AAB genomic group. A combination of the diagnostic character for the B genome (MaCIR108 allele size of 295 bp alone, or in combination with allele size of 289 or 287 bp) and the diagnostic character for the A genome (one allele size shorter than 270 bp) was found in some accessions of the ABB group. Other accessions with alleles with diagnostic characters for the A and B genomes were also identified as the ABB genomic group due to the presence of a Ma-3-90 allele with a size of 152 bp. In the 21 accessions classified as AAB, some contained a MaCIR108 allele with a size of 287 bp and shorter than 270 bp, but they did not have a Ma-3-90 allele with a size of 152 bp. Other AAB group members possessed a MaCIR108 allele with a size of 275 bp and shorter than 270 bp. In the 73 accessions of pure A genome, the Ma-3-90 allele with a size of 152 bp was not observed. The study also detected a Ma-3-139 allele with a size of 132 bp found in all banana accessions containing the B genome, while none was observed in banana accessions containing the A genome alone. This suggested that the locus Ma-3-139 could be considered as a diagnostic character for distinguishing banana accessions with the B genome from those with the A genome alone.

The number of pure A-genome accessions was greater than the number of B-genome accessions and their hybrids. These results also supported the hypothesis that Indonesia is one of the main centres of diversity and the centre of origin of *M. acuminata* 

(Daniells et al., 2001).

Analysis of Genetic Relationships among Banana Germplasm in Indonesia

The dendrogram generated using UPGMA analysis separated the 116 banana accessions into two main clusters with a similarity of 0.13 (Fig. 2). All banana accessions belonging to the BB, AAB and ABB genomic group clustered in the first main cluster together with most banana accessions containing the A genome alone. The second main

cluster was formed from 11 banana accessions with the A genome alone.

The polymorphism of microsatellite loci analysed in this study was consistent with a previous study by Creste et al. (2004), but was greater than that observed by Creste et al. (2003). Using nine primers, Creste et al. (2004) observed an average of 12.8 alleles, ranging from 10 to 15 alleles in a group of 58 *Musa* genotypes, including 49 AA and 9 AAB cultivars. However, Creste et al. (2003) detected an average of 6.1 alleles at each of 11 microsatellite loci studied in a group of 35 banana genotypes of various genomic compositions and ploidy levels. One possible reason for this difference is that all accessions used in this study originated from natural cultivars, with a relatively wide genetic base.

Genetic relationship analysis among banana accessions of various genomic groups demonstrated that the accessions containing the B genome were clearly distinct from each other (Fig. 2). The BB genomic group appears closer to the ABB genomic group. All banana accessions containing the B genome were clustered according to their genomic group, except four AAB accessions, namely 'Kepok Amerika', 'Raja Sereh', 'Embok'

and 'Susu', clustering with accessions containing the A genome alone.

With a similarity of 0.18, the dendrogram showed that the first main cluster was divided into two subclusters. The first subcluster contained diploid and triploid accessions containing the A genome alone (AA/AAA and AAA) and four AAB accessions. Cluster analysis based on 93 alleles of microsatellites could not significantly distinguish diploid

from triploid banana accessions of the A genome alone due to the dosage effects of microsatellite alleles. However, the majority of accessions, especially AAA accessions, tended to be clustered according to their ploidy levels. Similar results were reported by Creste et al. (2004) and in a previous study by Retnoningsih et al. (2010b) investigating genetic relationships among A-genome cultivars. In addition to the dosage effect, characteristics of the AA and AAA accessions were not clearly distinct, possibly due to

both groups originating from the same parents.

Four banana accessions with an AAB genome were separated from the second cluster which contained all banana accessions with the B genome. This is because these accessions had one specific allele of the MaCIR108 microsatellite with a size of 275 bp. According to the microsatellite marker determination key, this allele was not detected in the other genomic groups. They were closely related to banana accessions containing the A genome alone. Seventeen AAB accessions had a MaCIR108 allele with a size of 287 bp that was also observed in several accessions of ABB. Thus, this AAB genomic group was closely related to ABB.

Cluster analysis showed that microsatellite markers could discriminate each accession, except accessions which were known to be phenotypically similar. Ninety-nine unique genotypes were identified from the analysis of 116 accessions. In identical accessions, some had the same local name, whereas other accessions were synonymous.

In conclusion, 116 banana accessions were successfully classified into their genomic groups using a molecular key based on two SSR primer pairs. The key is only applicable to accessions that contain A and B genomes. For other cultivated bananas of S and T genomes, identification using molecular markers could likely be performed if primers were available. Cluster analysis based on 93 microsatellite alleles could not significantly distinguish diploid from triploid banana accessions. Incorporation of several M. acuminata subspecies in the analysis could likely reveal further genetic relationship with the cultivars.

#### ACKNOWLEDGEMENTS

The authors are greatly indebted to Ir. Agus Sutanto, RIF Solok, West Sumatra, staff of Diperta Yogyakarta and Margareta Rahayuningsih, for help in collection of plant materials and provided information about accessions. In addition, the authors would like to express their thanks to Prof. Dr. Mien A. Rifai for his critical review of the manuscript.

#### Literature Cited

Chakravarthi, B.K. and Naravaneni, R. 2006. SSR marker based DNA fingerprinting and diversity study in rice (Oryza sativa L.). Afr. J. Biotechn. 5:684-688.

Creste, S., Neto, A.T. and Figueira, A. 2001. Detection of single sequence repeat polymorphisms in denaturing polyacrylamide sequencing gels by silver staining. Plant. Mol. Biol. Rep. 19:299-306.

Creste, S., Neto, A.T., Silva, S.O. and Figueira, A. 2003. Genetic characterization of banana cultivars (Musa spp.) from Brazil using microsatellite markers. Euphytica

132:259-268.

Creste, S., Neto, A.T., Vencovsky, R., Silva, S.O. and Figueira, A. 2004. Genetic diversity of Musa diploid and triploid accessions from the Brazilian banana breeding program estimated by microsatellite markers. Genet. Res. Crop. Evol. 51:723-733.

Creste, S., Bennati, T.R., Orsi, M.R., Risterucci, A.M. and Figueira, A. 2005. Isolation and characterization of microsatellite loci from a commercial cultivar of Musa acuminata. Mol. Ecol. Notes. doi: 10.1111/j.1471.8286.2005.01209.x.

Daniells, J., Jenni, C., Karamura, D. and Tomekpe, K. 2001. Musalogue: A Catalogue of Musa Germplasm, Diversity in the Genus Musa. INIBAP, Montpellier, France.

Dixit, A. 1998. A simple and rapid procedure for isolation of Amaranthus DNA suitable for fingerprint analysis. Plant. Mol. Biol. Rep. 16:1-8.

Ford-Lloyd, B.V., Kaemmer, D., Kahl, G. and Lagoda, P.J.L. 1997. Molecular marker assisted analysis of the Musa genome complex. INIBAP, Montpellier, France.

Heslop-Harison, J.S. and Schwarzacher T. 2007. Domestication, genomics and the future for banana. Ann. Bot. 100:1073-1083.

Kaemmer, D., Fischer, D., Jarret, R.L., Baurens, F.C., Grapin, A., Dambier, D., Noyer, J.L., Lanaud, C., Kahl, G. and Lagoda, P.J.L. 1997. Molecular breeding in the genus *Musa*: a strong case for STMS marker technology. Euphytica 96:49-63.

Perera, L., Russel, J.R., Provan, J. and Powell, W. 2001. Levels and distribution of genetic diversity of coconut (*Cocos nucifera* L. var. *Typica* form *typica*) from Sri

Lanka assessed by microsatellite markers. Euphytica 122:381-389.

Pillay, M., Tenkouano, A., Ude, G. and Ortiz, R. 2004. Molecular characterization on genomes in *Musa* and its applications. p.271-286. In: S.M. Jain and R. Swennen (eds.), Banana Improvement: Cellular, Molecular Biology, and Induced Mutations. Science Publishers, Inc. Enfield (NH).

Retnoningsih, A., Megia, R. and Hartana, A. 2010a. Molecular verification and diversity analysis of Indonesian BB, AAB and ABB banana cultivars. Tree and Forestry

Science and Biotechnology 4 (Special Issue 1):69-76.

Retnoningsih, A., Megia, R. and Hartana, A. 2010b. Characterization and evaluation of *Musa acuminata* cultivars in Indonesia based on microsatellite markers. Floribunda (in press).

Rohlf, F.J. 1998. NTSYSpc: Numerical taxonomy and multivariate analysis system version 2.0. Use guide. New York: Exerter SofwareApplied Biostatistics Inc.

Semagn, K., Bjornstad, A. and Ndjiondjop, M.N. 2006. An overview of molecular marker methods for plants. Afr. J. Biotechn. 5:2540-2568.

Simmonds, N.W. and Shepherd, K. 1955. Taxonomy and origins of the cultivated bananas. Bot. J. Lin. Soc. 55:302-312.

Stover, R.H. and Simmonds, N.W. 1987. Bananas. Longman, UK.

Zhang, D.P., Carbajulca, D., Ojeda, L., Rossel, G., Milla, S., Herrera, C. and Ghislain, M. 2000. Microsatellite analysis of genetic diversity in sweetpotato varieties from Latin America. CIP program Report 1999-2000.

#### Tables

Table 1. Alleles size, number of alleles and observed heterozygosity value produced from analysis of 116 banana accessions using eight microsatellite markers.

Primer	Allele size range (bp)	Number of allele	Observed heterozygosity
MaCIR327b	388-436	5	0.45
Ma-1-132	330-378	9	0.51
MaCIR332a	260-296	11	0.85
MaCIR108	220-295	18	0.84
Ma-3-139	132-177	17	0.83
Ma-3-90	132-172	14	0.86
Ma-1-27	122-142	8	0.57
Ma-1-17	110-154	11	0.91
Total		93	
Mean		11.6	0.73

Fig. 1. The molecular determination key of banana genomic groups (Retnoningsih et al., 2010a).




Fig. 2. UPGMA clustering of 116 banana accessions in Indonesia based on eight microsatellite primers.

# **ACTA HORTICULTURAE®**

Acta Horticulturae<sup>®</sup> is used for publishing reviewed research and review articles presented at ISHS co-sponsored meetings.

Research reports of an oral or poster presentation should contain a commentary on a finished research project. A review article deals with knowledge obtained about a certain subject. Abstracts or copies of posters are not published.

As decided by the Council of ISHS, the peer reviewed proceedings of ISHS-symposia are published in a volume of Acta Horticulturae<sup>®</sup>. The regulations for the publication of the proceedings are mentioned in a special agreement, to be signed by the organizer of a symposium, by the institution involved and by the Executive Director of ISHS.

The ISHS Secretariat takes care of the publishing.

All published Acta Horticulturae® articles are available from www.actahort.org.

# Reprints

Authors can order reprints of their article(s) from the ISHS Secretariat. An order-form for reprints can be downloaded from **www.ishs.org**.

ISSN: 0567-7572

ISBN: 978 90 6605 138 6

Price for non-members of ISHS: € 112,-

