


**FACULTY OF ANIMAL SCIENCE
BOGOR AGRICULTURAL UNIVERSITY**


THE SECOND INTERNATIONAL SEMINAR ON ANIMAL INDUSTRY
“Empowering Local Resources for Sustainable Animal Production in Adapting
to Climate Change”

5-6 July 2012
Jakarta Convention Center, Jakarta-Indonesia

PROCEEDING



Organized by:



Supported by:



LIST OF CONTENTS

List of Editors	ii
Foreword from Chairperson of Organizing Committee	iii
Remarks from Dean of Faculty of Animal Science	v
Seminar Program	vii
Seminar Layout	xiv
List of Contents	xv
Invited Speaker	1
GM and Non-GM Rumen Microbes in Enhancing Animal Productivity. T.S. Park, J. K. Seo, & Jong K. Ha.....	3
Consumer Preferences in Meat. Louw Hoffman & Donna Cawthorn.....	9
Improving Local Feed Resource to Increase Nutrient Availability to Support Sustainable Agriculture. E.R. Ørskov.....	23
Planning Dairy Development Programs in Tropical Asia. J. B. Moran & J. W Brouwer.....	27
Carrier Proteins in Milk: Basic and Potential Applications. Kenji Fukuda.....	35
Indonesia Farm Animal Genetic Resources in Adapting to Climate Change. Ronny Rachman Noor.....	53
Tropical Forages in Indonesia: Past experience and Future Opportunity. H.M. Shelton.....	64
BREEDING AND GENETICS	75
Improvement the Genetic Potential of Local Chicken By Combination of Crossbreeding, Selection Method, Cellular Analysis and Nutritional Adjustment to Produce the Candidate of Local Layer. M. Aman Yaman, Yurliasni, Zulfan, & Muhammad Daud.....	77
Physical Meat Characteristics of Local Thin Tail Sheep based on Calpastatin (CAST) Genotype Variation. M.I.A. Dagong, C. Sumantri, R.R. Noor, R. Herman & M. Yamin.....	85
Genetic Variation of the IGF1 and OPN Genes in Holstein-Friesian Dairy Cattle of Historical and Non-Historical Twins. Anneke Anggraeni, Hasanatun Hasinah, Santi Ananda Arta, Bess Tiesnamurti, Restu Misrianti, & Eryk Andreas.....	91
Genetic Marker Approach for Confirming the Existing Twinning Trait in PO Cattle. Endang T. Margawati, Paskah P. Agung, & Muhamad Ridwan.....	97

Physical Meat Characteristics of Local Thin Tail Sheep based on Calpastatin (CAST) Genotype variation

M.I.A. Dagong¹, C. Sumantri², R.R. Noor², R. Herman² & M. Yamin²

¹Animal Production Department, Faculty of Animal Science, Hasanuddin University, Makassar, 90245, Indonesia.
e-mail:iccangdagong@yahoo.com

²Department of Animal Production and Technology, Faculty of Animal Science, Bogor Agricultural University, Bogor, 16680, Indonesia

Abstract

Calpastatin (CAST) is an indigenous inhibitor of calpain that involved in regulation of protein turn over and growth. The objective of this research was to identify genetic polymorphisms in the part of intron 5 - entire exon 6 of CAST gene in local sheep and their association with meat quality and muscle composition. A PCR-SSCP method was carried out to identify genetic variation of CAST gene. In total 401 heads of sheep from 8 subpopulations were investigated, three groups of samples were thin tail sheep from Sukabumi, Jonggol and Kissar. The rest samples were Priangan sheep from Margawati and Wanaraja and fat tail sheep from Donggala, Sumbawa and Rote islands. Twenty one heads from Jonggol were used for meat and muscle identification. SSCP analysis revealed that three different SSCP patterns corresponded to three different alleles in the CAST locus (CAST-1, 2 and 3 allele) with six different genotypes. Genetic variation between local sheep populations were calculated based on genotypic and allelic frequencies. Most populations studied were polymorphic, with genotype frequencies of CAST-11, CAST-12, CAST-22, CAST-13, CAST-23, and CAST-33 were 29.7%, 38.2%, 24.2%, 2.5%, 4.5% and 0.7% respectively. CAST-1 and 2 alleles were most commonly found in all populations with total frequency 95.8%, while rare allele was CAST-3 (4.2%) and only found in thin tail population. Based on sequence analysis identified a non-synonymous amino acid variation in exon 6 induced Gln/Leu substitution. There was no association between CAST alleles and genotypes with meat quality.

Key words: calpastatin, local sheep, meat quality, PCR-SSCP

Introduction

Local sheep is one of the genetic resource potential to be developed. This is due to their several advantages which are prolific, good adaptability to the harsh

environments, disease resistance, shorter production cycles and relatively requires small capital. In addition, in some densely populated areas like Java, sheep are able to substitute some beef that have to be imported each year. Based on 2008 data, special needs of sheep in West Java alone reach the range of 3.343.365 heads (Ditjennak 2009). The amount of the request indicates that the prospects for sheep farming is still wide open.

In relation to sheep development efforts, some weaknesses of the local sheep is that -their slaughter weights and daily body weight gain are low varied between 54 - 174 g/head/day (Yamin *et al.* 2009), in addition, the carcass quality and meat is also highly variable and do not meet international market standards. In order to solve the problem the genetic quality improvement efforts are needed to increase productivity, carcass and meat quality so the impact on lamb production will increase the contribution of lamb to total meat production in the country that currently only around 5% (Ditjennak 2009).

Molecular biotechnology advances allow the selection can be done at the DNA level through the use of marker genes that have an association with the highly economic traits. One of marker genes associated with body weight in local sheep was calpastatin (Sumantri *et al.* 2008). Numerous other studies have also shown a relationship of calpastatin gene with carcass quality (Schenkel *et al.* 2006), and meat quality, especially tenderness (Casas *et al.* 2006; Curi *et al.* 2009).

Calpastatin (CAST) is a member of the calpain-calpastatin system involving three molecules of the enzyme μ -calpain, m-calpain and calpastatin that serves as both calpain inhibitor. This system plays a important role in diverse physiological processes such as regulation of protein turn over and growth (Goll *et al.* 1992), and myoblast migration (Dedieu *et al.* 2003), therefore CAST is believed as a good candidate gene for growth, carcass and meat quality.

Information on meat quality based on various calpastatin gene genotype in the local sheep currently is not yet available, so the information is needed in order to describe the effect of this gene on meat quality.

Materials and Methods

Sample and Genotyping

This study used 21 heads of Thin Tail Sheep (TTS) from UP3J Jonggol that reared intensively. Sheep then grouped based on their genotype variation based on Dagong *et al.* (2011) methods. A pair of PCR primer, forward: 5'-GTTATGAATT-GCTTTCTACTC-3' and reverse: 5'-ATACGATTGAGAGACTTCAC-3' was designed to amplify part of intron 5 and whole exon 6 of CAST gene, as described by Zhou *et al.* (2007). PCR amplification was carried out in 25 μ l reaction containing 50-100 ng genomic DNA, 0.25 μ M of each primer, 200 μ M dNTPs (Fermentas), 4.0 μ M Mg^{2+} , 0.5 U of Toptaq DNA polymerase (Qiagen, Hilden, Germany), and 1x

the reaction buffer. The condition of thermal cycling consisted of pradenaturation at 95 °C for 5 min, followed by 35 cycles of denaturation 95 °C for 30 s, annealing 56 °C for 45 s, and extension 72 °C for 45 s. The final extension step was at 72 °C for 5 min. Amplification was carried out in a thermal cycler (Mastercycler Personal 22331, Eppendorf, Germany). The PCR amplicon were checked on 1.5% agarose gels in 0.5 x TBE buffer containing 10% of ethidium bromide at 100 volt for 45 min and visualized by UV transiluminator. A SSCP procedure was used to identify variation in the amplicon of CAST locus. The sheep that have been known to represent the CAST genotype then slaughter each genotype to identify their physical meat characteristics.

Physical Meat Characteristics

Meat quality was measured based on the physical parameters which include: pH measurement by pH meter and measured after aging for 24 hours. Meat tenderness was shown by the enormous strength (kg/cm²) required to cut the meat cores indicated by the needle cutter Warner Bratzler Shear Force (WBSF) that moves over the scale with a measurement sensitivity of 0.1 kg/cm². Water Holding Capacity was measured with a planimeter by finding out the amount of water (mg H₂O). Cooking loss was measured by subtracting the initial weight with the weight after sample cooked at 80 °C for 1 hour.

Association of CAST gene polymorphisms with physical meat quality was analyzed by t-test with the following statistical equation:

$$t = \frac{\bar{X}_1 - \bar{X}_2}{\sigma \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \quad \sigma = \sqrt{\frac{\sum_{i=1}^{n_1} (\bar{X}_1 - \bar{X}_1)^2 + \sum_{i=1}^{n_2} (\bar{X}_2 - \bar{X}_2)^2}{n_1 + n_2 - 2}}$$

Where :

- \bar{X}_1 and \bar{X}_2 = Mean value in genotype 1 and genotype 2
- n_1 and n_2 = Number of sample in genotype 1 and genotype 2
- σ = Total varians

Physical meat quality data corrected in advance using the following statistical equation:

$$X_i \text{ correction} = \frac{\bar{X}_{\text{standard}}}{\bar{X}_{\text{observed}}} \times X_i \text{ observed}$$

Where :

X_i correction = value of physical meat characteristics after being corrected by sex and age

\bar{x} standard = mean of physical meat characteristics of standard population

\bar{x} observed = mean of physical meat characteristics of observed population

X_i observed = value of physical meat characteristics before being corrected by sex and age

Results and Discussion

Differences of physical meat characteristic

Means value of physical meat quality from different CAST genotypes on the local sheep are shown in Table 1. There was no significant difference ($P>0.05$) either in tenderness, water holding capacity, cooking loss and pH from three different CAST genotypes (CAST-11, CAST-12 and CAST-22) in local sheep. Similar results with a previous study by Zhou *et al.* (2008), who reported that all allelic variation (CAST-1, 2 dan 3) or variations of genotypes were identified in the CAST locus did not significantly affect the lamb tenderness.

Tenderness value of research results in the range 2 - 3 in a tender category, but no differences among the three genotypes. In contrast to some previous studies that identified a significant association between CAST variation with beef tenderness. In cattle, CAST gene variations have been used commercially as genetic markers. Two markers are currently available commercially were *GeneSTAR Tenderness* and *Igenity TenderGENE*. *GeneSTAR* using SNP G/A in 3'UTR region (Barendse 2002),

Tabel 1. Mean value of physical meat characteristics from various CAST genotypes of local Thin Tail Sheep

Physical meat characteristics	Genotypes					
	CAST-11 (n=4)	CV(%)	CAST-12 (n=10)	CV(%)	CAST-22 (n=7)	CV(%)
Tenderness (Kg/cm ²)	3.16±0.72	22.78	2.98±0.79	26.51	2.49±0.71	28.81
Cooking loss (%)	49.54±3.24	6.54	43.76±3.61	8.25	46.32±6.53	14.10
Water Holding Capacity (WHC) (MgH ₂ O)	117.31±13.70	11.68	99.46±19.83	19.94	103.73±19.27	18.58
Percentage of WHC (% MgH ₂ O)	39.10±4.56	11.66	33.15±6.60	19.91	34.57±6.42	18.57
pH _{ult} (24 h)	5.57±0.07	1.26	5.75±0.31	5.39	5.80±0.32	5.54

Note: CV = Coefficient of variation (standard deviation/mean x 100%)

while *Igenity TenderGENE* using SNP G/C in intron 5 region (Van Eenennaam *et al.* 2007).

Differences in CAST gene effect sheep meat tenderness and beef probably caused by the fact that sheep meat is more tender due to the rate of myofibrils proteolysis of sheep meat is faster than beef (Koochmaraie *et al.* 1991), therefore, the differences in sheep meat tenderness has smaller effect on meat tenderness.

Conclusion

There was no difference between physical meat quality with CAST genotype variation in local sheep.

Acknowledgements

This research was funded by DP2M Dikti through *Hibah Kompetensi* project with contract No: 224/SP2H/PP/DP2M/III/2010 and DIPA IPB through *Hibah Disertasi Doktor* with contract No : 10/I.3.24.4/SPK/PDD/2011.

References

- Barendse, WJ, 2002. DNA markers for meat tenderness in cattle. Commonwealth Scientific Industrial and Research Organisation. International patent publication WO 02/064820 A1.
- Casas, E., S. N. White, T. L. Wheeler, S. D. Shackelford, M. Koochmaraie, D. G. Riley, C. C. Chase, D. D. Johnson, & T. P. L Smith. 2006. Effects of calpastatin and mikro calpain markers in beef cattle on tenderness traits. *J. Anim. Sci.* 84: 520-525.
- Curi, R. A., L. A. L. Chardulo, M. C. Mason, M. D. B. Arrigoni, A. C. Silveira, & H. N. de Oliveira. 2009. Effect of single nucleotide polymorphism of CAPN1 and CAST genes on meat traits in Nellore beef cattle (*Bos indicus*) and their crosses with *Bos taurus*. *Anim. Genet.* 40: 456-462.
- Dedieu S, Mazeret G, Poussard S, Brustis JJ, Cottin P. 2003. Myoblast migration is prevented by a calpain-dependent accumulation of MARCKS. *Biol. Cell.* 95 : 615 – 623.
- [Ditjennak]. Directorate General of Livestock. 2009. Livestock Statistics. Jakarta : Ministry of Agriculture Republic of Indonesia.
- Goll DE, Thompson VF, Taylor RG, Christiansen JA. 1992. Role of the calpain system in muscle growth. *Exp. Cell. Res.* 74 : 225 – 237.
- Koochmaraie, M., G. Whipple, D.H. Kretchman, J.D. Crouse and H.J. Mersmann. 1991. Postmortem proteolysis in longissimus muscle from beef, lamb and pork carcasses. *J Anim. Sci.* 69:617–624

- Schenkel FS *et al.* 2006. Association of a single nucleotide polymorphism in the calpastatin gene with carcass and meat quality traits of beef cattle. *J. Anim. Sci.* 84: 291 – 299.
- Sumantri C, Diyono R, Farajallah A, Inounu I. 2008. Polymorphism of calpastatin gene and its effect on body weight of local sheeps. *JITV.* 13 : 117 – 126.
- Van Eenennaam AL *et al.* 2007. Validation of commercial DNA tests for quantitative beef quality traits. *J. Anim. Sci.* 89 : 891 – 900.
- Yamin M *et al.* 2009. Increasing local sheep growth performance through rapid selection at fattening farm. *Sustainable Animal Production for Food Security and Safety. Proceeding the 1st International Seminar on Animal Industry (ISAI).* IPB ICC, 23 – 24 Nov 2009. Bogor : IPB Faculty of Animal Science. pp 57 – 60.
- Zhou H, Byun SO, Frampton CM, Bickerstaffe R, Hickford JGH. 2008. Lack association between CAST SNPs and meat tenderness in sheep. *Anim. Genet.* 39 : 328 – 332.