

METHODOLOGICAL APPROACHES IN RUMINANT METABOLIC RESEARCH¹

PENDEKATAN METODOLOGI DALAM PENELITIAN METABOLISME PADA RUMINANSIA

Djokowoerjo Sastradipradja²

ABSTRACT

Climatic ~~factors~~ of high temperatures and humidities of humid tropical countries, alongside with the **occurrence** of diseases and parasites have created a complex situation in terms of the effect on animal production. Studies on quantitative relationships for better understanding of digestion, metabolism and related areas are lacking. There are however a number of appropriate methodologies available for conducting **onfarm** metabolic research aiming at giving better understanding of animal production and health in the humid tropics. The paper discusses methodologies and experimental approaches developed in the authors's laboratory on growing, ~~draft~~, pregnant and lactating **ruminants**. Several examples of the results are presented.

ABSTRAK

Faktor-faktor suhu tinggi dan kelembaban lingkungan tropika lembab, digabungkan dengan tingginya kejadian **penyakit** dan **parasit** menciptakan **situasi** yang **kompleks** yang berpengaruh negatif terhadap upaya produksi ternak. Penelitian-penelitian mengenai **aspek kuantitatif fungsi pencernaan**, **metabolisme hewan** dan **bidang-bidang terkait** boleh dikatakan **sangat minim**. Meskipun demikian, **sebenarnya cukup tersedia** metodologi yang **sesuai** untuk **melaksanakan "onfarm metabolic research"** yang bertujuan memperoleh **pemahaman** yang **mendalam** mengenai produksi dan kesehatan **ternak** di daerah tropika lembab. **Makalah ini membahas beberapa metodologi dan pendekatan percobaan yang dikembangkan di laboratorium penulis pada hewan ruminansia yang sedang tumbuh, bunting, laktasi dan ternak kerja. Beberapa contoh hasil penelitian disajikan untuk ilustrasi.**

¹ Center for Metabolic Research, presented at a one-day seminar "Improvement of Animal Performance through Onfarm Metabolic Research" organized by the Faculty of Veterinary Medicine Bogor Agricultural University (IPB) at JL. Taman Kencana 3 Bogor 16151, 30 April 1997, INDONESIA

² Department of Physiology and Pharmacology, Faculty of Veterinary Medicine, Institut Pertanian Bogor (Bogor Agricultural University), Jalan Taman Kencana 3, Bogor 16151, INDONESIA

INTRODUCTION

Food is necessary to built up tissue and to act as a source of energy. Food has to be digested into absorbable units in the gastrointestinal tract. Metabolism is the process refered to all the chemical and energy transformations that occur in the body.

The humid tropical countries have special and often unique problems associated with animal production. Both imported and indigenous animals used to build the livestock industry shows variation in adaptability to the existing **agro-ecological** setting of the region. The countries have dietary materials available, particularly for ruminants, which in **many** instances have not been evaluated for their ability to support the desired production. Humid tropical countries have adverse climatic factors of high temperatures and humidities and, in addition, there are a wide variety of diseases and parasites which reduce production. Direct effects of these factors **and** interactions between the factors have created a complex situation in terms of the effect on production.

As qualitative knowledge increased, detailed consideration is given to develop quantitative relationships to increase further understanding and integrate various aspects, **e.g.** to bring together quantitative approaches concerned with elucidating mechanisms, used in the **study** of digestion (monogastric and ruminant), metabolism and related areas. In his address to the 11th Symposium on Energy Metabolism of **Farm** Animals on "Past Achievements and Future Perspective in Energy Metabolism", the late Sir Kenneth L. **Blaxter** (1989) made the following **remarks**: *"Current world literature did not contain much information which relates to climatic and seasonal or exogenous hormonal effects on metabolism, pregnancy in mammals, egg secretion in birds and effects of muscular work have not been considered. Nor has the comparative aspect of metabolism been considered On the practical side we need these are items of commerce. We also need, more accurate and rapid methods for estimating the energy values of feed & sice; to take into account breed and strain differences in animal requirements. Better estimates of the composition of body gains (retentions) are required. New demands will arise as a result of modern techniques of genetic manipulation of livestock and the use of exogenous hormones derived through, biotechnology, to increase reproductive performance and the rate and composition of growth. On the fundamental side, we must explore aspects of adaptation of metabolism to undernutrition, season and climate at the cell, organ and whole animal levels and unravel the complexity of the endocrine and neural control mechanisms involved focussing on*

the roles of the sympathetic nervous system and growth factors. New techniques must be explored since they may well enable studies to be made in natural environments rather than the laboratory. Biochemical methods of assessing the energy status of animals demands attention". These hold true for the humid tropical situation as well.

Brody (1945) stated that growth is the basis of and closely related to many animal productive processes, which includes egg, **milk**, fat and other production. He defined growth as the constructive and assimilatory synthesis of one substance at the expense of another, **i.e.** nutrient which undergoes dissimilation. Growth, and hence animal production, is biological synthesis, production of new biochemical units and it is the aspect of development concerned with the increase in living substance or protoplasm and includes one or all of the following processes, **e.g.** cell multiplication, cell enlargement, and incorporation of material taken from the environment. From the practical point of view of quantitative measurement of growth of the organism as a whole, nonprotoplasmic inclusions in the body must be considered as parts of the growth process.

The aforesaid processes are more dramatic during periods following starvation and injury. Seasonal variations in animal feed supplies **often** cause interrupted patterns of growth in young animals due to feed restriction resulting from seasonal variations or other causes. A subsequent restoration of feed supplies causes the animals to exhibit enhanced gain in weight called compensatory growth. The underlying physiological processes of growth are complex involving feed utilization, visceral tissue mass and **carcase** protein and fat deposition, variation in energy requirements according to a particular condition and also in other nutrients, to meet maintenance and body performance, partition of nutrients, endocrine control, etc. So **defined**, growth (which includes production) is inseparable from metabolism.

Our studies on the responses of ruminant (and other livestock) animals to different rations or related treatments rely basically on traditional balance methods alongwith proximate analysis of foodstuffs and waste. To **measure** the transfer of substances within the intact animal and their partition to organs and tissues, tracer isotope methodologies are used in combination with experimental approaches such as the application of the Fick principle and dilution techniques. The data range over a wide field area and include important considerations as the energy cost of maintaining animals, of transforming feed into body tissue and the desired animal products. Data on body composition enable assessment of growth responses to changes in nutrition or in the

environment. The studies are reviewed in this presentation highlighting on experience in the use of methodologies appropriate to the existing conditions.

The methodologies and experimental approaches presented are based on the work on growing animals. As the discussions proceed to touch on other **animal productive** states, the appropriate methodologies will be inserted.

GROWING RUMINANTS

Animal Balance Trial

The balance trial provides information related to the requirements of nutrients for growth, maintenance and production and the **availability** of nutrients from **feedstuffs**. It is used to show changes in percentage absorption of nutrients as a **function** of the level of nutrients in the diet. For energy balance, the energy categories of feed, **i.e.** gross energy (GE), faecal energy (**FE**), digestible energy (DE), urinary energy (UE), Energy loss with digestive gas production (**En-CH₄**), metabolizable energy (ME), energy expenditure (EE) = health production (HP) and retained energy (RE), should be calculated. Energies in feed, **faeces** and urine are determined by bomb calorimetry or calculated from organic **ingredients** multiplied by their respective caloric equivalents, **e.g.** UE may be found from g **urinary-N.d⁻¹** x 34.0 kJ, but energy in gases and **CH₄** is rather difficult to measure. It could be reasonably well approximated from gas test results or simply taken as 8-10 % of GE, or using **Blaxter's equation** **En.CH₄** = 4.28 + 0.059 D **Kcal/100Kcal** GE (D = digestibility of energy, %). For the energy balance, data on HP and RE are needed. When physical exercise is absent, it is safe to use the relationship ME = HP + RE. Since ME is found from GE **subtracted** by all energy **found** in waste, either HP or RE should be determined while the other could then be calculated by difference **from** the ME. If HP should be **measured**, a suitable technique of animal calorimetry is required. On the other hand if RE is chosen to be determined, the slaughter technique or a suitable **in vivo** body composition technique should be used. Alongside energy balance, protein (or N) balance could be simultaneously measured.

Whole Animal Calorimetry and Body Composition

All major components of the body are in a continuous **state of flux**: intakes, excretions, degradations and resynthesis. All **endproducts** of digestion not only act as substrates for the accretion of body tissue (anabolism), but also **serve** as **precursors** for the production of the high-energy compounds needed (catabolism) for maintenance and to provide the energy for the synthesis of the macromolecules of the body. In addition, there **is** an extensive **interconversion** of metabolites and the rate of productive processes in **the** body is governed by the complex interplay between the intermediary metabolism of the **various** nutrients **and** the **rate** of synthesis and degradation of tissue lipid, protein, nucleic acid and carbohydrate **stores**.

1, Measurement of Whole Body HP by Carbondioxide Entry Rate' Technique

Due to unavailability of respiration chamber **calorimetry** for large animals, but on the other hand we have in our **laboratory** at our disposal radioisotope detection equipment, Carbondioxide Entry Rate Technique (CERT) using tracer ^{14}C -bicarbonate is the method of choice to measure whole body energy expenditure of small ruminants (Sastradipradja, 1992). The method involves primed continuous infusion of the label solution into a blood vein and after reaching steady state condition of isotope concentration in body fluid bicarbonate, serial blood samples are withdrawn. The CO_2 production rate (rCO_2) is calculated by dividing the rate of label infused by the plateau specific activity of blood bicarbonate. At an accepted RQ value, the rCO_2 value is converted into its energy equivalent which is HP. Double polyethylene catheters are implanted in the jugular veins for minimal disturbance to the animal, easy delivery of isotope and convenient serial blood sampling. It is important to measure the specific activity of CO_2 over a sufficiently long period of time (**e.g.** 12h or longer) to ensure a more representative mean value of specific activity. For such long trials, instead of blood, samples of any body fluid **can** be taken (**e.g.** saliva instead of blood). To get faster attainment of stable specific activity we apply the primed-continuous infusion technique and our experience shows that the ratio between primer dose and infusion rate per minute is 80 to 1 giving satisfactory results. Primer dose is 1 ml of $\text{NaH}^{14}\text{CO}_3 = 40 \mu\text{Ci}$ delivered within 1 min. followed by continuous infusion at $0.5 \mu\text{Ci}/\text{min}$. The CO_2 production **can** be estimated **from** the plateau specific activity according to the equation (Corbett *et al.*, 1971):

$$\text{CO}_2 = \frac{\text{rate of tracer bicarbonate infused}}{\text{plateau specific activity of CO}_2}$$

The advantage of CERT is that the **animal** is **free** to move **around without restraint**. The use of CERT also enables **measurement** of **gluconeogenesis** involving $^{14}\text{CO}_2$ fixation. Another advantage of **CERT** is **the** ability of **measuring** glucose kinetics during the **same** trial by administering ^3H -glucose at **the same time** with the bicarbonate label.

2. *Measurement of HP as the Difference Between ME and RE*

In our work with **swamp** buffaloes during the last four years, we have calculated HP as the difference between ME and RE included physical work (**Mahardika et al.**, 1997) with satisfactory results (see exercise metabolism below).

3. *Heart Rate as A Predictor of HP*

We are still developing this technique on small **ruminants** and cattle following the success of **longterm heart** rate measurement in swamp buffaloes using the Polar Sport Tester (Finland).

4. *Estimation of Body Composition in vivo*

4.a *Urea Space Technique* (Rule et al., 1986)

Urea is used as a marker for body water. A measured amount of urea in saline is injected into the jugular vein within one minute. Exactly 12 minutes after the injection, a sample of blood is withdrawn from the jugular vein and the concentration of urea determined. Urea space is found by dividing the dose (mg) of urea **infused** by the increment of blood urea concentration following the **infusion** from the **preinfusion** value times body weight times 10. The empty body water (EBW, %) = $59.1 + 0.22 \text{ US (\%)} \cdot 0.04 \text{ BW}$, while empty body fat (%) = $19.5 \cdot 0.31 \text{ US (\%)} + 0.05 \text{ BW}$. The choice of this technique is necessary considering the hazards involved in the use of radioisotope labelled water on large ruminants **and** the costly slaughter technique with ruminant livestock animals. We have introduced this technique in our work with goats and sheep. Validation of the *in vivo* body composition method using US was separately done on four goats by doing slaughter technique (ST) analysis in addition to the *in vivo* calculations on the same animals in question (**Arta Putra et al.**, 1997). EBW estimated according to Rule et al. (1986) resulted in an underestimation of 4.2% from the EBW value by ST. Therefore, the use of US in goats required a correction factor of 1.044. For use with small ruminants, **Panaretto's** equation for body protein (**Panaretto**, 1963) and for body fat (Panaretto and Till, 1963) are

based on EBW. The use of our **corrected** values in aforesaid **equations** resulted in **good** agreement with our ST results. The **results** are **presented** in **Table 1**.

Table 1. **Body Composition (% BW) According to Slaughter Technique Analysis and in vivo Equations Estimated from Four Female Growing Goats**

	Body water		Body protein		Body fat	
	ST	Eq. 1	ST	Eq. 2	ST	Eq.3
	----- %BW-----		----- %BW-----		----- %BW-----	
Mean \pm SD	60.9 \pm 5.94	60.9 \pm 0.06	20.2 \pm 0.39	20.2 \pm 0.02	16.2 \pm 5.74	17.6 \pm 0.08

ST = slaughter technique analysis

Eq. 1 = EBW according to Rule *et al.* (1986) US equation multiplied by 1.044,

Eq. 2 = Panaretto (1963) body protein equation using EBW value from Eq. 1,

Eq. 3 = Panaretto and Till (1963) body fat equation using EBW value from Eq. 1.

Validation results with the Javanese thin tail (JTT) sheep (Saka and Sastradipradja, unpublished data) revealed that only the equation which relate EWB with US was significantly different **from** the regression equation according to Rule *et al.* (1986), as follows

$$\text{EBW (\%)} = 64.1 - 21.1\text{US\%} - 0.378\text{BW}$$

$$(P < 0.05; S_{y.x} = 2.185; R^2 = 0.425)$$

Consequently, only this equation can be used for sheep with correction factors respectively for the intercept, US and BW to be 1.0846, -95.9091 and 9.45. From our studies with female swamp buffaloes, the equation of Rule *et al.* (1986) if applied for working buffaloes should be modified to:

$$\text{Fat} = 13.69 - 0.21\text{US} + 0.03\text{BW}; \quad \text{Protein} = 19.7 + 0.08\text{US} + 0.11\text{BW}.$$

4.b Body Density Method (Kleiber, 1961)

Application of this technique in small ruminants and cattle is **being** conducted **after** it has been **successfully** practiced in the swamp buffalo. Contrary to the situation in swamp buffaloes who love wading in water, other livestock animals will resist to immersion in water. Therefore, a solid frame should be constructed where the animal can be immobilized and the whole device including the animal can be subsequently lowered in the water tank leaving the head above water level.

Supporting Measurements on Metabolism and Digestion

Glucose is an **important nutrient** for all tissues, especially it is indispensable as energy source for nerve cells and for foetus in pregnant animals, for the synthesis of lactose, fat and providing carbon skeleton for **others**. An important component of **growth** is fat synthesis. Glucose is required during growth, in particular for the supply of the NADPH via the pentosephosphate pathway for **fat** synthesis. A lower amount is required for **protein** synthesis. The major portion of glucose available to the **ruminant** is supplied by gluconeogenesis. Additional demands for **glucogenic precursors occur during** lactation, namely for the lactose secreted in **the** milk and for the reducing power needed **to** provide energy for milk synthesis. Impaired gluconeogenesis is believed to contribute to various metabolic disorders frequently seen in ruminants, such as acetonaemia in dairy cows and pregnancy toxaemia or twin lamb disease in sheep. In ruminant studies, glucose metabolism warrants attention.

Successful operation of whole animal metabolism requires low levels to function properly. Many metabolic peculiarities of ruminants stem from the proper functioning of the splanchnic region. It is understandable that special attention should be given to metabolism in this region.

A great deal of research has been carried out on the digestive system of the ruminant giving **understanding** of the characteristic metabolism which cope with the peculiar products of microbial digestion. We attempted to measure products of ruminal digestion and integrate it for our overall understanding of animal performance.

1. Gluconeogenesis Involving CO₂ Fixation

Tracer labeling of the bicarbonate pool (CERT) may enable one to estimate gluconeogenesis (GNG) **from** carbon transfer involving **CO₂** fixation into the glucose pool. The rationale involves using the transfer quotient (TQ) between **CO₂** and glucose to assess the extent of **CO₂** fixation in GNG from precursors like propionate and lactate. The TQ in question is calculated as the ratio between the plateau specific activities (per at. C) of **¹⁴C** in glucose (product) and in bicarbonate (precursor). The rate of GNG involving **CO₂** fixation is found by multiplying the value of the glucose **flux** by the TQ times 6. Measurement of the glucose **flux** and pool size is done simultaneously with the CERT infusion in the same animal by pulse injection with tritiated glucose (Sastradipradja, 1992).

2. Metabolism of the **Portal Drained Viscera**

Many metabolic peculiarities of ruminants stem **from** the viscera, that is a group of organs whose blood supply **drains** into the portal vein. Quantification of nutrients in the **splanchnic** region is useful to **understand the** metabolism in this region. We have experience to **study** this **aspect** in small ruminants by the **arteriovenous** difference technique (Katz and **Bergman**, 1969) involving surgical procedures to obtain **samples from** the **mesenteric, portal and** hepatic veins and of any artery (Astuti, 1995; Sastradipradja *et al.*, 1997). Portal blood flow were measured on unaesthetized animals (**i.v.** injection of **xylazine** 0.05 **ml/kg** BW, followed by **i.v.** injection of ketamine 0.11 **ml/kg** BW). Lidocaine (1 **ml/** animal) was applied locally on the site of incision. The primer dose of **PAH-³H** was 5.75 **µCi** delivered in one minute via the jugular vein followed by continuous infusion at 0.6 **µCi/min** into the right mesenteric vein. After around two hours of **infusion**, blood samples were collected for analysis of blood gasses (**CO₂, O₂, VFA**) and other relevant constituents. Metabolism of the region and absorption rates of nutrients can be measured. The technique is invasive however, which need appropriate skills to perform. So far we haven't been successful in placing permanent catheters in aforesaid blood vessels. Feasible techniques to measure ruminal fermentation production rates would be an alternative approach to measure the nutrients' supply to the animal.

3. In vitro **Techniques for Ruminal Fermentation**

The techniques closely simulate **in vivo** conditions especially for **rumen** fermentation studies: rate of formation of endproducts (VFA), rate of release of **NH₃**, production rate of microbial protein and gas production.

3.a' **Microbial Protein Synthesis** (Suwandyastuti *et al.*, 1985)

A shortterm incubation of **rumen** contents at 39 °C in an artificial **rumen**: test **tubes**, glass syringes or **flasks**. The anaerobiosis is maintained by introducing **N₂** gas replacing the air in the artificial **rumen** above the incubation mixture. **³³P** is used as tracer for measuring microbial protein synthesis (MPS).

3.b **Incubation of Rumen Contents in Glass Syringes (gas test)** (**Menke** and **Steingass**, 1988)

The amount of gas which is released when feedstuffs are incubated **in vitro** with **rumen** fluid measures **CO₂** and **CH₄** production. It is closely related to digestibility and **consequently** to the energetic feed value of feedstuffs for ruminants. The incubation **mixture** consists of feed

sample, main and **trace** element solution, buffer solution, **rezazurin solution** (indicator), **reduction** solution and rumen fluid. Hundred ml syringes are used **preheated** at 39 °C. The syringes **plus** contents **free** of air bubbles are incubated in a (39 °C) preheated incubator. The **volume** of **gas** in the piston is read off eight hours after the **start** of the incubation (**V₈**). Incubation is **continued** and the final reading is done **after** 24 hours (**V₂₄**). Regression equations will be derived relating ME with gas production, feed nutrient components and **digestibilities**.

3.c VFA Production Rates: **Zerotime** in vitro *Method* (Whitelaw *et al.*, 1970)

A sample of **rumen** contents is taken and subsamples incubated *in vitro* under anaerobic **conditions**. The rate of production of individual and total VFA is calculated **from** the increments in acid concentration obtained by incubating the subsamples for different periods and extrapolating back to zero time incubation to give the rate of VFA production per unit volume at the time the sample was removed. Equations for performing the calculations are given by **Whitelaw et al.** (1970). The **rumen** volume should be known in order to calculate total ruminal production.

3.d *Use of Nylon Bags Incubated in The Rumen* (IAEA, 1985)

The digestibility test requires animals fitted with permanent **rumen** fistulae. Nylon bags should have a pore size 20 - 40 µm, dimensions 15 x 8 cm, sample size 3-5 g of air-dry feed, ground through screen 2-5 mm, incubation times up to 24h for protein concentrates, up to 72h for roughage feeds. The technique can describe both the rate and the extent of degradation, affected by the **rumen** environment such as ammonia level, pH, **type** of feeds, trace minerals, etc.

4. In vivo *Estimation of Digestion*

Evaluation on the intake **and** digestion characteristics of feedstuffs, needs quantitative data to describe the movement of **digesta** along the gastrointestinal tract.

4.a *Estimation of Rumen Volume* (IAEA, 1985)

The volume of **rumen** liquid is **estimated from** the dilution of ⁵¹Cr-EDTA tracer introduced **intraruminally** by way of a syringe and needle in a period between **rumen** contractions. After 1 hour and then every hour up to eight hours, **sample/withdraw** representative **rumen** contents by **way** of a stomach tube. The animal should receive a (nearly) continuous feeding regime. Plot