

ISSN 1411-982X
12/Akred-LIPI/P2MBI/9/2006

Indonesian Journal of
**AGRICULTURAL
SCIENCE**

Indonesian Agency for Agricultural Research and Development



Accredited A by the Indonesian
Institute of Sciences
No. 1417/D/2006

INDONESIAN JOURNAL OF AGRICULTURAL SCIENCE

Vol. 9 No. 2, October 2008

Indonesian Journal of Agricultural Science is previously published as Indonesian Journal of Crop Science (1985 - 1999). This journal is published in one volume of two issues per year by the Indonesian Agency for Agricultural Research and Development, Indonesia. It is available on-line at: <http://pustaka-deptan.go.id>.

The journal publishes primary research articles from any source if they make a significant original contribution to the experimental or theoretical understanding of some aspect of agricultural science in Indonesia. The definition of agricultural science is kept as wide as possible to allow the broadest coverage in the journal.

*Director of Indonesian Center for Agricultural Library and Technology
Dissemination*

Mei Rochjat D.

Editorial Board

Chairman

Supriadi *Plant Pathology*
Indonesian Medicinal and Aromatic Crops
Research Institute

Members

Budiman Hutabarat *Socioeconomics*
Indonesian Center for Agricultural Socio Economic
and Policy Studies

Fahmuddin Agus *Soil Science*
Indonesian Soil Research Institute

Raffi Paramawati *Food Science and Technology*
Indonesian Center for Agricultural Engineering
Research and Development

M. Winarno *Horticultural Plant Breeding*
Indonesian Center for Horticultural Research
and Development

Sutoro *Agronomy*
Indonesian Center for Agricultural Biotechnology and
Genetic Resources Research and Development

I Made Tasma *Plant Breeding and Molecular Biology*
Indonesian Center for Agricultural Biotechnology and
Genetic Resources Research and Development

Managing Editor

Endang Setyorini Indonesian Center for Agricultural Library and Technology
Dissemination

Intan Yudia Nirmala Indonesian Center for Agricultural Library and Technology
Dissemination

Ujang Sahali Indonesian Center for Agricultural Library and Technology
Dissemination

All inquiries and manuscripts should be sent to the:

The editorial office

Indonesian Center for Agricultural Library and Technology Dissemination

Jl. Ir. H. Juanda 20

Bogor 16122, Indonesia

Telephone : Bogor (0251) 8321746

Fax. : 62-251-8326561

E-mail : pustaka@pustaka-deptan.go.id.

Website : <http://pustaka-deptan.go.id>.

THE SEGREGATION PATTERN OF INSECT RESISTANCE GENES IN THE PROGENIES AND CROSSES OF TRANSGENIC ROJOLELE RICE

Satoto^a, Yuli Sulistyowati^b, Alex Hartana^c, and Inez H. Slamet-Loedin^b

^aIndonesian Center for Rice Research, Jalan Raya 9, Sukamandi, Subang 41256, Indonesia

^bResearch Center for Biotechnology, Indonesian Institute of Science, Jalan Raya Bogor km 46, Cibinong 16911, Indonesia

^cBogor Agricultural University, Kampus IPB Darmaga, Bogor 16680, Indonesia

ABSTRACT

Successful application of genetic transformation technique, especially in developing rice variety resistant to brown plant hopper and stem borer, will depend on transgene being expressed and the gene inherited in a stable and predictable manner. This study aimed to analyse transgene segregation pattern of the progenies and the crosses of transgenic rice cv. Rojolele harboring *cry1Ab* and *gna* genes. The third generation (T_2) of five transgenic Rojolele events containing *gna* and/or *cry1Ab* were evaluated for two generations to identify the homozygous lines and to study their inheritance. The homozygous lines were selected based on the result of PCR technique. The segregation patterns of *gna* and *cry1Ab* were studied in eight F_2 populations derived from Rojolele x transgenic Rojolele homozygous for *cry1Ab* and or *gna* and their reciprocal crosses. Data resulted from PCR of F_2 population were analysed using a Chi Square test. The study obtained six homozygous lines for *gna*, namely A22-1-32, A22-1-37, C72-1-9, F11-1-48, K21-1-39, K21-1-48, and two homozygous lines for *cry1Ab*, namely K21-1-39 and K21-1-48. Both *cry1Ab* and *gna* transgenes had been inherited through selfing and crossing with their wild type as indicated from the F_1 containing *gna* and *cry1Ab* as many as 48.4% and 47.4%, respectively. In six of the eight crosses, *gna* was inherited in a 3:1 ratio consistent with Mendelian inheritance of a single dominant locus, while in the remaining two crosses, *gna* was segregated in a 1:1 ratio. The presence of *cry1Ab* in F_2 populations also showed a 3:1 segregation ratio in all crosses. In the F_2 population derived from F_1 plant containing *cry1Ab* and *gna*, both transgenes segregated in a 9:3:3:1 dihybrid segregation ratio. This study will add to the diversity of genetic sources for insect resistance and allow further use of these transgenic lines for pyramiding resistance to brown plant hopper and stem borer or separately in rice breeding programs whenever the efficacy tests and biosafety requirements have been completed.

[**Keywords:** *Oryza sativa*, transgenic plant, genetic resistance, segregation, genetic inheritance]

INTRODUCTION

Brown plant hopper and stem borer are the major insects causing rice production loss in Indonesia. The progress of classical rice breeding for brown plant hopper and stem borer resistance is hampered

due to the quick formation of the new biotype of brown plant hopper causing break of resistance even in the elite breeding lines, while for stem borer the major problem is the lack of resistant genes available in the respective species and their wild relatives (Rao and Padhi 1988). The International Rice Research Institute (IRRI) has screened more than 1500 entries for yellow rice stem borer and 6000 entries for stripe rice stem borer resistance (Jackson 1995), resulted some lines which have low to moderate resistance..

Plant genetic engineering is one of the complementary tools to the conventional breeding for the development of new varieties. This technique has some advantages such as the wider germplasm options, reduction of the number of backcrossing needed to eliminate one of the parental genetic backgrounds and the precision of transferring only the target gene without any other unknown genes introgressed (Conner 1997). A number of transgenic food crops have been released globally using this techniques such as maize, soybean, and rice (www.agbios.com). In this reported work, the parental material harbors *gna* gene encoding for snowdrop (*Galanthus nivalis*) agglutinin and synthetic *cry1Ab* gene encoding crystal protein from *Bacillus thuringiensis* (Sardana *et al.* 1996). Genetic transformation using *gna* gene in tobacco for resistance to aphid was first reported by Hilder *et al.* (1995), followed by the work on rice targeted to brown plant hopper (Rao *et al.* 1998; Sudhakar *et al.* 1998), while *cry* gene has been extensively used in maize for resistance to stem borer (Armstrong *et al.* 1995; Jansens *et al.* 1997) and also in rice (Ghareyazie *et al.* 1997; Breitler *et al.* 2000). Other biotic stress resistance genes introduced to rice were among others the *wasabi defensin* gene for resistance to blast diseases (Kanzaki *et al.* 2002), *potato proteinase inhibitor II* gene (Duan *et al.* 1996), and *CpTi* gene (Xu *et al.* 1996).

Plants have mechanisms for defending themselves from microorganisms and insects. Plant produces

lectins as one of their defense strategies against insects (Carlini *et al.* 2002). GNA that binds specifically to a sugar called mannose is rather toxic to certain kinds of insect pests of important crops, including rice (Fitches *et al.* 2001). Puzstai studies showed that purified GNA was not toxic to rats (Puzstai *et al.* 1990). The gene coding for GNA was an attractive choice for increasing insect resistance of crops and at that time the gene was introduced into a number of crops, including potatoes and rice and proved to increase their resistance to some important insect pests (Rao *et al.* 1998; Foissac *et al.* 2000).

Bacillus thuringiensis (Bt) crystal proteins are protoxins that emerge when exposed to alkaline medium. The protoxins are proteolytically cleaved into smaller active form (molecular weight 60-70 kDa) derived from the N-terminal half of the protein. Although the mode of action is largely unknown, it is predicted to bind to receptor proteins of the insect gut causing pore formation (Hoffman *et al.* 1988). Bt insecticide is commonly used in organic farming.

Transformation of Indonesian cultivar Rojolele rice by particle bombardment (Slamet-Loedin *et al.* 1998) and *Agrobacterium* has been developed in Indonesia using a number of genes (Slamet-Loedin *et al.* 1997; Mulyaningsih *et al.* 2004; Rahmawati and Slamet-Loedin 2006). In this reported work, the inheritance study was carried out on five different independent transgenic lines harboring target genes the *cry1Ab* and snowdrop lectin genes obtained through co-transformation by particle bombardment techniques (Slamet-Loedin *et al.* 1998). For further use in the breeding programs, it is important to ensure that the genes are dominant genes and segregate following Mendelian segregation. The study aimed to analyse the segregation pattern of *cry1Ab* and *gna* genes in F₂ populations derived from F₁ progenies of crosses between a transgenic Rojolele rice containing *cry1Ab* and/or *gna* genes, and their wild types as well as their reciprocal crosses.

MATERIALS AND METHODS

This study was carried out from June 2000 to July 2002 at the laboratories and the transgenic biosafety containment glass house of the Research Center for Biotechnology, Indonesian Institute of Science, Bogor, Indonesia.

Plant Materials

All breeding lines in this reported study were the progenies of Rojolele, a local long duration variety

from javanica subgroup (150-160 days) that has premium grain quality, few number of tillers, good grain filling, low productivity, and sensitive to brown plant hopper and stem borer. Rojolele was chosen because it has good cell regeneration ability and high economic value, but it is susceptible to brown plant hopper and stem borer.

The plant materials consisted of progenies of the third generations (T₂) of five transgenic events of Rojolele resulted from the earlier experiments (Slamet-Loedin *et al.* 1998), namely line A22-1, C72-1, E65-1, F11-1, and K21-1. The lines were obtained through co-transformation of three different plasmids. The first plasmid was p-Ubi-Cry IAB containing *cry1Ab* gene coding for δ endotoxin from *B. thuringiensis*, for resistance to stem borer. The second plasmid was p-Ubi-Gna containing *gna* gene coding for lectin from snowdrop for resistance to brown plant hopper and both genes were controlled by ubiquitin promoter. The third plasmid was pWRG 1515, containing *hpt* gene coding for *hygromycine phosphotransferase* and *gus A* gene coding β -glucuronidase controlled by CaMV35S promoter. *Hpt* gene was used to recognize the transformant among the bombarded cells in the early transformation process, while *gus A* gene was used to detect whether targeted gene has already been inserted into plant genome. All lines at the stage of the primary transgenic (T₀) have been analysed by molecular and biochemical techniques to consist of the two genes of interest (*gna* and *cry1Ab*) or individually apart from the marker genes (*hpt* and *gus-A*). The material used in this study were the segregated T₂ populations. The lines were selected based on the presence of the genes on T₁ generation. Transgenic lines A22-1 and C72-1 harbor only *gna* (lectin from snowdrop coding for resistance to brown plant hopper). Line E65-1 harbors only *cry1Ab* gene from *B. thuringiensis* (coding for δ endotoxin responsible for resistance to stem borer), whereas F11-1 and K21-1 contain both *gna* and *cry1Ab* genes.

PCR Analysis

The presence of inserted genes was analysed using PCR techniques. Specific primer sets were used to detect the presence of the codon optimized synthetic *cry1Ab* gene (Sardana *et al.* 1996) and the *gna* gene (Hilder *et al.* 1995). Primer sets used were as follows: forward (5'-CATTGTGTCTCTCTCCCC-3') and reverse (5'-CCGTTAGAGAAGTTGAAAGG-3') for *cry1Ab* and forward (5'-ATGGCTAAGGCAAGTC TCCTC-3') and reverse (5'-TCATTACTTGCCGTCAC AA G-3') for *gna*. As an internal control to ensure the plant

DNA was present in the reaction in most PCR reactions, a multiplex PCR was carried out with *gos-5* internal rice gene (Mayer *et al.* 1992) using the following primer set: forward (5'-CGACCTCGAGGACATCGGC AACAC-3') and reverse (5'-GCCGAGCAGCAGGAACTTGAGCAG G-3'). Expected band gene fragment sizes for *cry1Ab*, *gna* and internal control *gos-5* are 1012, 400 and 231 bp, respectively.

Seeds from the progenies of the five transgenic lines were germinated for 2 weeks. Five centimeters of the first leaf were harvested and stored in the liquid nitrogen. Genomic DNA isolation was carried out following the method of Zheng *et al.* (1995).

Individual genomic DNA samples were subjected for PCR. The PCR reaction contained 14 μ l mix buffer (1x PCR buffer, dNTP 0,05 mM, primer forward 2,50 ng μ l⁻¹ and primer reverse 2,50 ng μ l⁻¹ and Taq polymerase 0,05 μ l μ l⁻¹) were put into PCR tube then added 1 μ l DNA sample. This mixture was put into thermocycler *Gene Amp* (PCR system 2400, Perkin Elmer) and amplified for 40 cycles. PCR conditions for *gna* gene were 1 cycle 95°C 1 minute, 40 cycles with temperature of 95°C, 65°C, and 72°C, 1 minute each, 1 cycle 72°C 2 minutes and storage under 4°C. For *cry* gene, the PCR conditions were 1 cycle 95°C 1 minute, 40 cycles with temperature of 95°C, 55°C, and 72°C, 1 minute each, 1 cycle 72°C 2 minutes and storage under 4°C. Amplified DNA separation was done by electrophoresis in the 1.2% gel agarose and the result was documented using Bio Rad Gel Doc 1000.

Determination of Homozygosity of the Transgenic Lines

T₂ generation lines were selected from the T₁ parental which positively contained both *gna* and/or *cry1Ab* genes and having more than 30 seeds each. From the T₂ generation of rice lines A22-1, C72-1, E65-1, F11-1, and K21-1, eight progeny lines were selected. Fifty seeds of each selected progeny lines were grown and analysed for their inheritance. Selection of homozygous lines was done based on the gene occurrence identified from PCR result. Eleven individual plants were then selected to fulfill testing requirement number for each gene study. Eleven is the minimum number of plants to show at least one unexpected phenotype appeared under 95% probability for character controlled by a recessive allele (Sedcole 1977). However, to reduce the number of PCR reactions needed and cost, instead of carrying out directly 11 reactions for each line, initially the test was carried out on five individual plant samples from each line. Since if from five samples (five PCR reactions) there

was already sample showing no amplicon (negative for the presence of the gene), the parental must be heterozygous, therefore addition of six other sample assay would not be needed. When the five initial samples gave positive results for the presence of the expected band, six additional samples were added to obtain 11 sample number, the minimum set of number to ensure the homozygosity at the level of confidence 95%.

The Segregation Pattern of *cry1Ab* and *gna* Genes in the Transgenic Rice

The segregation pattern of both *cry1Ab* and *gna* genes in transgenic rice was studied based on the PCR positive result on F₂ population derived from both Rojolele wild type x transgenic Rojolele and their reciprocal crosses. The crossing was made between transgenic Rojolele and its wild type to ensure the success rate, since inter-variety group crossing such as to the Ciherang will have to overcome biological incompatibility if exists. In the first season, crossing between Rojolele and transgenic Rojolele homozygous for *gna* and *cry1Ab* and their respective reciprocal crosses were made. In the second season, F₁ seeds from the crossing were planted. The presence of both *cry1Ab* and *gna* genes in each individual F₁ plant was identified using PCR. The F₁ and its reciprocal crosses positively containing *cry1Ab* and *gna* genes were then selfed to produce F₂ seeds. In the third season, the F₂ seeds resulted from the F₁ and its reciprocal cross showing positive *cry1Ab* and or *gna* were planted and analysed their gene inheritance. The segregation pattern of both genes was analysed using a Chi Square test.

RESULTS AND DISCUSSION

Determination of Homozygosity of the Transgenic Lines Based on the Molecular Analysis

Plant materials used in this experiment were five transgenic progeny lines resulted from genome modification of Rojolele variety. The lines was obtained from co-transformation of *gna* and *cry1Ab* genes located at different plasmids, therefore some lines have two genes and some lines only have one of the two genes (Slamet-Loedin *et al.* 1998). In this study further analysis was carried out to identify the homozygosity status of the parents for further reciprocal analysis. In the case of transgenic application, the inserted gene will be integrated in one of

the chromosome pairs. Therefore if the gene is a dominant gene and inserted in one locus, it will follow a 3:1 Mendelian segregation pattern for a single dominant gene in the subsequent generation since rice is primarily self-pollinated plant.

As previously described in the protocol, PCR analysis using specific primer for the *gna* gene was carried out on the two steps to reduce the cost. At the step 1 at screening, five PCR reactions of each germinated seed from the same line were carried out. When all five samples showed the expected bands, then additional six samples are added to obtain 11 positive progenies. The PCR analysis results of the two-step analyses are summarized in Table 1 and 2. Based on the presence of the PCR amplicon using primers specific for *gna* on the first five samples, in total there were 10 transgenic lines harbour *gna* gene (A22-1-27, A22-1-31, A22-1-32, A22-1-37, C72-1-9, F11-1-31, F11-1-48, K21-1-46, K21-1-39, and K21-1-48), thus we continued with additional samples to reach 11 samples. The result showed that six lines (A22-1-32,

Table 1. Transgenic lines of rice harboring *gna* gene based on PCR amplicon presence.

| Transgenic event | Target genes presence in T ₁ | Lines showing <i>gna</i> amplicon on first five samples | Lines showing <i>gna</i> amplicon on 11 samples |
|------------------|---|---|---|
| A22-1 | <i>gna</i> | A22-1-27 A22-1-31 A22-1-32 A22-1-37 | A22-1-32 A22-1-37 |
| C72-1 | <i>gna</i> | C72-1-9 | C72-1-9 |
| E65-1 | <i>cry1Ab</i> | - | - |
| F11-1 | <i>cry1Ab</i> and <i>gna</i> | F11-1-31 F11-1-48 | F11-1-48 |
| K21-1 | <i>cry1Ab</i> and <i>gna</i> | K21-1-46 K21-1-39 K21-1-48 | K21-1-39 K21-1-48 |

Table 2. Transgenic lines of rice harboring *cry1Ab* gene based on PCR amplicon presence.

| Transgenic lines | Target genes presence in T ₁ | Lines showing <i>cry1Ab</i> on first five samples | Lines showing <i>cry1Ab</i> amplicon on 11 samples |
|------------------|---|---|--|
| E65-1 | <i>cry1Ab</i> | - | - |
| F11-1 | <i>cry1Ab</i> and <i>gna</i> | - | - |
| K21-1 | <i>cry1Ab</i> and <i>gna</i> | K21-1-39 K21-1-44 K21-1-46 K21-1-48 | K21-1-39 K21-1-48 |

A22-1-37, C72-1-9, F11-1-48, K21-1-39, and K21-1-48) derived from five transgenic events were homozygous for *gna* within 95% confidence interval (Table 1). As mentioned above, Sedcole (1977) explained statistically that 11 is the minimum number of plants must be tested to show at least one unexpected phenotype appeared under 95% probability for a character controlled by a recessive allele. If from the 11 individual plants there is one phenotype from unexpected recessive allele, the plant population must have originated from heterozygous parent plant. On the contrary, if all of the 11 individual plants showed phenotype from expected dominant allele, it means that the population was originated from homozygous parent plant for the respective gene. PCR analysis results of K21-1-48 and A22-1-32 lines are presented in Figure 1 and 2, respectively.

Step 1 PCR analysis for *cry1Ab* gene on five DNA samples of four lines (K21-1-39, K21-1-44, K21-1-46, and K21-1-48) harbored the *cry1Ab* gene (Table 2), thus six additional samples were analysed. Figure 3 showed the gel electrophoresis of K21-1-48 showing amplicon band of *cry1Ab*. Further analysis using 11 DNA samples showed only two lines namely K21-39 and K21-48 that possibly homozygous for *cry1Ab* gene within confidence interval of 95% (Fig. 4).




Fig. 1. PCR result of five samples for *gna* gene in the second generation of transgenic rice of K21-1-48 line, 1 = λ , 2 = -, 3 = fragment *cry*, 4 = fragment *gna*, 5 = control (+), 6 = control (-), 7 = water, 8-22 = five DNA samples of transgenic line K21-1-48.




Fig. 2. PCR result of transgenic rice of A22-1-32 line homozygous for *gna*; 1 = λ *Hind*III, 2 = fragment *gna*, 3 = control (+), 4 = control (-), 5 = water, 6-16 = DNA samples of transgenic rice A22-1-32.




Fig. 3. PCR result of five samples for *cry1Ab* gene in the transgenic rice of T2 generation, 1 = λ *Hind*III, 2 = water, 3 = fragment *cry*, 4 = control (+), 5 = control (-), 6-35 = DNA samples of transgenic lines (27-31 = five DNA samples for K21-1-48). Positive control plant is the plant DNA sample of earlier generation shown to have *cry1Ab* gene by southern and PCR analysis. Negative control was DNA sample of the wild type nontransgenic Rojolele.




Fig. 4. PCR result of homozygous transgenic rice of K21-1-48 line containing *cry1Ab* gene; 1 = λ *Hind*III, 2 = fragment *cry*, 3 = control (+), 4 = control (-), 5 = water, 6-16 = DNA samples of the homozygous transgenic line K21-1-48.

The Segregation Pattern of *Cry1Ab* and *gna* Genes in the Transgenic Rice

All lines identified to be homozygous for the transgenes were crossed with the Rojolele wild type; reciprocal crosses were also carried out. It was observed that eventhough the genetic background of parental plants was the same, some of transgenic lines flowered earlier than their wild type, and therefore, the number of plants suitable for crossing was limited. In general, there was no difficulty to cross Rojolele wild type to the homozygous transgenic lines.

Rojolele wild type x transgenic Rojolele crosses produced F_1 seeds with seed set varied from 16% (Rojolele x K21-1-39) to 63% (C72-1-9 x Rojolele) (Table 3). Rojolele line K21-1-39 had the lowest percentage of seed set because it produced less pollen.

PCR of the F_1 plants showed that 31 out of 64 plants (48%) contained *gna* gene, while for *cry1Ab*, 9 out of 19 plants (47%) contained the gene (Table 4). PCR analyses of F_1 plants of transgenic lines homozygous for *gna* and *cry1Ab* with Rojolele were presented Figure 5 and 6, respectively.

Some of the F_1 plants that have neither *gna* nor *cry1Ab* genes may be because the crossing was

failed to take place, therefore the parent (Rojolele) was self-pollinated. Another possibility was that the parent of the expected homozygous plant may actually be a heterozygous plant due to the stringency of our identification test that was at 95% level. Similar

Table 3. Number of F₁ seeds resulted from crosses between Rojolele wild type and homozygous transgenic rice lines.

| Crosses | Number of crosses | Number of F ₁ seeds resulted from each cross | Seed set (%) |
|---------------------|-------------------|---|--------------|
| Rojolele x A22-1-32 | 13 | 476 | 61 |
| Rojolele x A22-1-37 | 12 | 443 | 62 |
| Rojolele x C72-1-9 | 22 | 755 | 57 |
| Rojolele x K21-1-39 | 5 | 47 | 16 |
| Rojolele x K21-1-48 | 17 | 545 | 53 |
| A22-1-32 x Rojolele | 19 | 677 | 59 |
| A22-1-37 x Rojolele | 14 | 323 | 39 |
| C72-1-9 x Rojolele | 18 | 678 | 63 |
| K21-1-39 x Rojolele | 18 | 232 | 22 |
| K21-1-48 x Rojolele | 16 | 452 | 47 |

result was reported by Datta *et al.* (2002) in the crosses between two homozygous transgenic lines, TT-103 (homozygous for *Xa21* gene) and TT-9 (homozygous for *RC7* and *Bt* genes) resulted some F₁ that have no *RC7* gene. Another cross (TT-9 X TT-103) also produced some F₁ plants having no *Xa21* gene. The inheritance of transgenes through various crossing experiments performed in the present study indicates that the transgenes have already integrated into the plant genome and will be useful in rice breeding as the lines can be used as donor parent to improve the resistance of rice varieties to stem borer.

Results of the PCR analyses of samples taken from the F₂ population derived from the F₁ plants showed that there were three genotypes observed, those are genotypes containing *gna* or *cry1Ab*, or both genes (*gna* and *cry1Ab*), and those neither have any of the gene. This result indicates that both *gna* and *cry1Ab* were integrated independently in the rice genome. The recent report of Pusztai indicated the possibility that lectin from snowdrop gene may have not passed

Table 4. Number of F₁ rice plants positively harbor *gna* or *cry1Ab* gene.

| Crosses | Number of plants used in PCR analysis for <i>cry1Ab</i> gene | Number of plants positively harbor <i>cry1Ab</i> gene | Number of plants used in PCR analysis for <i>gna</i> gene | Number of plants positively harbor <i>gna</i> gene |
|---------------------|--|---|---|--|
| Rojolele x A22-1-32 | | | 14 | 3 |
| A22-1-32 x Rojolele | | | 4 | 4 |
| Rojolele x A22-1-37 | | | 7 | 3 |
| A22-1-37 x Rojolele | | | 3 | 3 |
| Rojolele x C72-1-9 | | | 10 | 7 |
| C72-1-9 x Rojolele | | | 12 | 4 |
| Rojolele x K21-1-48 | 9 | 5 | 7 | 4 |
| K21-1-48 x Rojolele | 10 | 4 | 7 | 3 |
| Total | 19 | 9 | 64 | 31 |




Fig. 5. PCR for *gna* gene in F₁ rice plants; 1 = *gna* gene fragment, 2 = control (+), 3 = control (-), 4 = water, 5,6,7 = samples of C72-1-9 x Rojolele, 8,9,10 = samples of A22-1-37 x Rojolele, 11,12,13,14 = samples of A22-1-32 x Rojolele, 15,16,17 = samples of Rojolele x A22-1-37, and 18,19,20 = samples of Rojolele x A22-1-32.

the food safety test for release yet, therefore, in our subsequent works the *gna* gene study has been discontinued to avoid possible food safety issues. This results negate the hypothesis suggested that transgenes tend to cluster in one locus even they are located in different plasmids (Kohli *et al.* 1999). This result also suggests that co-transformation is a possible approach when a gene is intended to be out-segregated in the final product, such as the selectable antibiotic marker gene required for transformation initially but preferred not to be presence in the final product.

The amplification result of F_2 population for segregation pattern of either *gna* or *cry1Ab* gene, and

segregation of both genes was presented in Table 5, 6, and 7. In general F_1 plants performed better than their parents particularly for plant height and percentage of unfilled grains. The height of the F_1 plants was normally higher than that of both parents suggesting that the *cry1Ab* and *gna* transgenes have been inherited through selfing and crossing. When crossed with the wild type, 48.4% and 47.4% of the F_1 population contained *gna* and *cry1Ab*, respectively. In six of the eight crosses, *gna* was inherited in a 3:1 segregation ratio consistent with Mendelian inheritance of a single dominant locus. In the remaining two crosses, Rojolele x C72-1-9 and C72-1-9 x Rojolele, *gna* was segregated in 1:1 ratio. Similar result was




Fig. 6. PCR for *cry1Ab* gene in the F_1 rice plants. λ = lambda DNA cut by *EcoRI*, 1 = *cry1Ab* gene fragment, 2 = control (-), 3 = control (+), 4 = water, 5-9 = Rojolele x K21-1-48, 11-15 = K21-1-48 x Rojolele.

Table 5. Segregation of *gna* gene in F_2 rice populations derived from F_1 and their respective reciprocal crosses.

| Crosses | n | Expected ratio ¹⁾ | | Observed ratio | | X^2 | df = 1; α = 0.05; X^2 = 3.84 |
|---------------------|----|------------------------------|-------|----------------|-----|-------|---|
| | | (+) | (-) | (+) | (-) | | |
| Rojolele x A22-1-32 | 27 | 20.25 | 6.75 | 20 | 7 | 0.012 | Ratio 3:1 |
| A22-1-32 x Rojolele | 27 | 20.25 | 6.75 | 16 | 11 | 3.560 | Ratio 3:1 |
| Rojolele x A22-1-37 | 26 | 20.50 | 6.50 | 19 | 7 | 0.150 | Ratio 3:1 |
| A22-1-37 x Rojolele | 30 | 22.50 | 7.50 | 26 | 4 | 2.170 | Ratio 3:1 |
| Rojolele x C72-1-9 | 58 | 29.00 | 29.00 | 27 | 31 | 0.280 | Ratio 1:1 |
| C72-1-9 x Rojolele | 51 | 25.50 | 25.50 | 26 | 23 | 0.250 | Ratio 1:1 |
| Rojolele x K21-1-48 | 59 | 44.25 | 14.75 | 48 | 11 | 1.270 | Ratio 3:1 |
| K21-1-48 x Rojolele | 60 | 45.00 | 15.00 | 48 | 12 | 0.800 | Ratio 3:1 |

¹⁾(+) = present and (-) = absent of *gna* gene in the F_2 , progeny of the respective crosses.

Table 6. Segregation of *cry1Ab* gene in F_2 rice population derived from F_1 and its reciprocal cross.

| Crosses | n | Expected ratio ¹⁾ | | Observed ratio | | X^2 | df = 1; α = 0.05; X^2 = 3.84 |
|---------------------|----|------------------------------|-------|----------------|-----|-------|---|
| | | (+) | (-) | (+) | (-) | | |
| Rojolele x K21-1-48 | 59 | 44.25 | 14.75 | 49 | 10 | 2.04 | Ratio 3:1 |
| K21-1-48 x Rojolele | 60 | 45.00 | 15.00 | 47 | 13 | 0.6 | Ratio 3:1 |

¹⁾(+) = present and (-) = absent of *cry1Ab* gene in the F_2 , progeny of the respective crosses.

Table 7. Segregation of *cry1Ab* and *gna* genes in F₂ rice populations derived from F₁ plant and its reciprocal cross.

| Crosses | n | Expected ratio | | | | Observed ratio | | | | X ² | df = 3; α = 0.05; X ² = 7.82 |
|---------------------|----|----------------|--------------|--------------|------------|----------------|--------------|--------------|------------|----------------|---|
| | | <i>cry/gna</i> | <i>cry/-</i> | <i>-/gna</i> | <i>-/-</i> | <i>cry/gna</i> | <i>cry/-</i> | <i>-/gna</i> | <i>-/-</i> | | |
| Rojolele x K21-1-48 | 29 | 16.4 | 5.4 | 5.4 | 1.8 | 19 | 4 | 5 | 1 | 1.16 | Ratio 9:3:3:1 |
| K21-1-48 x Rojolele | 30 | 16.9 | 5.6 | 5.6 | 1.9 | 23 | 2 | 4 | 1 | 5.39 | Ratio 9:3:3:1 |

reported by Christou *et al.* (1996) for *bar* gene which means that this aberrant segregation can be explained by the passage of the transgene exclusively through one gamete.

The presence of *cry1Ab* in F₂ population also showed a 3:1 segregation ratio in all crosses. In the F₂ population derived from F₁ plants containing *cry1Ab* and *gna*, both transgenes segregated in a 9:3:3:1 dihybrid segregation ratio.

CONCLUSION

Both *cry1Ab* and *gna* transgenes had been inherited through selfing and crossing with their wild type (48.4% of F₁ containing *gna* and 47.4% of F₁ containing *cry1Ab*). In six of the eight crosses, *gna* was inherited in a 3:1 ratio consistent with Mendelian inheritance of a single dominant locus. In the remaining two crosses, Rojolele x C72-1-9 and C72-1-9 x Rojolele, the *gna* gene was segregated in 1:1 ratio. Progeny of Rojolele wild type x Rojolele transgenic and its respective reciprocal crosses showed the same segregation pattern indicating that both *gna* and *cry1Ab* genes have been integrated in the rice genome.

Both *gna* and *cry1Ab* genes, which originally derived from different plasmids, were independently inherited and segregated in a 9:3:3:1 ratio indicating that both genes were located in two nonhomologous chromosomes; thus negated the other reports suggesting transgenes introduced by particle bombardment tend to integrate in the same loci even they were located at different plasmids originally.

ACKNOWLEDGMENT

The authors wish to thank the technical assistance given by Agus Rachmat, Anky Zannati, and Yeni Andriani. This work was supported by the funding from the Research Center for Biotechnology, Indonesian Institute of Science (DIP 2000-2002).

REFERENCES

- Amstrong, C.L., G.B. Parker, J.C. Pershing, S.M. Brown, P.R. Sanders, D.R. Duncan, T. Stone, D.A. Dean, D.L. DeBoer, J. Hart, A.R. Howe, F.M. Morrish, M.E. Pajean, W.L. Petersen, B.J. Reich, R. Rodrigues, C.G. Santino, S.J. Sato, W. Schuler, R.S. Sims, S. Stehling, L.J. Tarochione, and M.E. Fromm. 1995. Field evaluation of European corn borer control in progenies of 173 transgenic corn events expressing an insecticidal protein from *Bacillus thuringiensis*. *Crop Sci.* 35: 550-557.
- Breitler, J.C., V. Marfa, M. Royer, D. Meynard, J.M. Vassal, B. Vercambre, R. Frutos, J. Messenger, R. Gabarra, and E. Guiderdoni. 2000. Expression of a *Bacillus thuringiensis cry1B* synthetic gene protects mediterranean rice against the stripe stem borer. *Plant Cell Rep.* 19:1195-1202.
- Carlini, C.R. and M.F. Grossi-de-Sá. 2002. Plant toxic proteins with insecticidal properties. A review on their potentialities as bioinsecticides. *Toxicon* 40(11):1515-1539.
- Christou, P., P. Vain, A. Kohli, M. Leech, J. Oard, and S. Linscombe. 1996. Introduction of multiple genes into elite rice varieties - Evaluation of transgene stability, gene expression, and field performance of herbicide-resistant transgenic plant. In G.S. Khush (Ed.). *Rice Genetic III. Proc. the Third International Rice Genetics Symposium*, International Rice Research Institute, Manila, Philippines. 1011 pp.
- Conner, A.J. 1997. *Genetically Engineered Crops. Environmental and Food Safety Issues*. The Royal Society of New Zealand. Miscellaneous Series 39.
- Datta, K., N. Baisakh, T.K. Maung, J. Tu, and S.K. Datta. 2002. Pyramiding transgene for multiple resistance in rice against bacterial blight, yellow stem borer, and sheath blight. *Theor. Appl. Genet.* 106: 1-8.
- Duan, X., X. Li, Q. Xue, M. Abo-El-Saad, D. Xu, and R. Wu. 1996. Transgenic rice plants harboring an introduced *potato proteinase inhibitor II* gene are insect resistant. *Nature Biotechnol.* 14: 494-498.
- Fitches, E., S.D. Woodhouse, J.P. Edwards, and J.A. Gatehouse. 2001. *In vitro* and *in vivo* binding of snowdrop (*Galanthus nivalis* agglutinin; GNA) and jackbean (*Canavalia ensiformis*; Con A) lectins within tomato moth (*Lacanobia oleracea*) larvae; mechanisms of insecticidal action. *J. Insect Physiol.* 47(7): 777-787.
- Foissac, X., Ng.T. Loc, P. Christou, A.M.R. Gatehouse, and J.A. Gatehouse. 2000. Resistance to green leafhopper (*Nephotettix virescens*) and brown planthopper (*Nilaparvata lugens*) in transgenic rice expressing snowdrop lectin (*Galanthus nivalis* agglutinin; GNA). *J. Insect Physiol.* 46(4): 73-583.

- Ghareyazic, B., F. Alinia, C.A. Menguito, L.G. Rubia, J.M. de Palma, E.A. Liwanag, M.B. Cohen, G.S. Khush, and J. Bennet. 1997. Enhanced resistance to two stem borers in an aromatic rice containing a synthetic *cry1Ab* gene. *Mol. Breed.* 3: 401-414.
- Hilder, V.A., K.S. Powell, A.M.R. Gatehouse, J.A. Gatehouse, L.N. Gatehouse, J. Shi, W.D.O. Hamilton, A. Merryweather, C.A. Newell, J.C. Timans, W.J. Peumans, E. van Damme, and D. Boulter. 1995. Expression of snowdrop lectin in transgenic tobacco plants results in added protection against aphids. *Transgenic Res.* 4: 18-25.
- Hoffman, C., P. Luthy, R. Hutter, and V. Pliska. 1988. Binding of the delta endotoxin from *Bacillus thuringiensis* to brush border membrane vesicle of the cabbage butterfly (*Pieris brassicae*). *Eu. J. Biochem.* 173: 85-91.
- Jackson, M.T. 1995. Protecting the heritage of rice biodiversity. *Geo-Journal* 35: 267-274.
- Jansens, S., A. van Vliet, C. Dickburt, L. Buysse, C. Piens, B. Saey, A.D. Wulf, V. Gossele, A. Paez, E.M. Gobel, and M. Peferoen. 1997. Transgenic corn expressing a *cry9C* insecticidal protein from *Bacillus thuringiensis* protected from European corn borer damage. *Crop Sci.* 37: 1616-1624.
- Kanzaki, H., S. Nirasawa, H. Saiton, M. Ito, M. Nishihara, R. Terauchi, and I. Nakamura. 2002. Over-expression of the *wasabi defensin* gene confers enhanced resistance to blast fungus (*Magnaporthe grisea*) in transgenic rice. *Theor. Appl. Genet.* 105: 809-814.
- Kohli A, X. Fu, R.M. Twyman, and P. Christou. 1999. Transgene integration, organization and expression. *Rice Genetics Newsl.* 16: 135-138.
- Mayer, A., G.B. Zondag, and L.A.M. Hensgens. 1992. A simple screening method for transgenic rice tissue based on PCR. *Rice Genetic Newsl.* 8: 161-165.
- Mulyaningsih, E.S., S. Indrayani, and I.H. Slamet-Loedin. 2004. Analisis molekuler dan uji ketahanan padi transgenik yang mengandung gen khitinase generasi ketiga (T2) terhadap *Rhizoctonia solani* Kuhn dan *Phyricularia oryzae* cav. *Majalah Ilmiah Biologi Biosfera A Sci. J.* 22(3): 142-151.
- Pusztai, A., S.W.B. Ewen, G. Grant, W.J. Peumans, E.J.M. Van Damme, L. Rubio, and S. Bardocz. 1990. Relationship between survival and binding of plant lectins during small intestinal passage and their effectiveness as growth factors. *Digestion* 46: 308-316.
- Rahmawati and I.H. Slamet-Loedin. 2006. Introduksi gen *cry1B-cryIA* ke dalam genom padi (*Oryza sativa*) cv. Rojolele menggunakan transformasi *Agrobacterium*. *Hayati* 13(1): 19-25.
- Rao, P.S.P. and G. Padhi. 1988. Improved sources of plant resistance to yellow stem borer (YSB) *Scirpophaga incertulas* Walker in rice. *Intl. Rice Res. Newsl.* 13: 5.
- Rao, K.V., K.S. Rathore, T.K. Hodges, X. Fu, E. Stoger, D. Sudhakar, S. Williams, P. Christou, M. Barathi, D.P. Brown, K.S. Powell, J. Spence, A.M.R. Gatehouse, and J.A. Gatehouse. 1998. Expression of snowdrop lectin (GNA) in transgenic rice plants confers resistance to rice brown planthopper. *Plant J.* 15(4): 469-477.
- Sedcole, J.R. 1977. Number of plants necessary to recover a trait. *Crop Sci.* 17: 667-668.
- Sardana, R., S. Dukhandjiev, M. Giband, X. Cheng, K. Cowan, C. Sauder, and I. Altosaar. 1996. Construction and rapid testing of synthetic and modified toxin gene sequences *cryIA* (b&c) by expression in maize endosperm culture. *Plant Cell Rep.* 15: 677-681. *Proc. National Academic of Science USA* 95: 2767-2772.
- Slamet-Loedin, I.H., A. Estiati, W. Rahayu, S. Hutajulu, and S. Nugroho. 1997. Plant regeneration toward transformation of several javanica rice cultivars. *Intl. Rice Res. Notes* 22(1): 16.
- Slamet-Loedin, I.H., A.S. Rahayu, P. Deswina, S. Purwantomo, E.S. Mulyaningsih, D. Sudhakar, J.A. Gatehouse, I. Altosaar, and P. Christou. 1998. Production of fertile transgenics aromatic Indonesian javanica rice co-expressing the snowdrop lectin and *cry1Ab* anti-insect proteins. *Proc. 4th Asia-Pacific Conference on Agriculture Biotechnology*, 13-16 July 1998, Darwin, Australia.
- Sudhakar, D., X. Fu, E. Stoger, S. Williams, J. Spence, D.P. Brown, M. Bharathi, J.A. Gatehouse, and P. Christou. 1998. Genetic engineering of rice for resistance to sap sucking insect pests. *In* P.J. Larkin (Ed.). *Agricultural Biotechnology: Laboratory, Field, and Market. Proc. 4th Asia-Pacific Conference on Agricultural Biotechnology*, 13-16 July 1998, Darwin, Australia.
- Xu, D., Q. Xue, D. McElroy, Y. Mawal, V.A. Hilder, and R. Wu. 1996. Constitutive expression of a cowpea trypsin inhibitor gene, *CpTi*, in transgenic rice plants confers resistance to two major rice insect pests. *Mol. Breed.* 2: 167-173.
- Zheng, K., N. Huang, J. Bennet, and G.S. Khush. 1995. PCR-based marker-assisted selection in rice breeding. *Intl. Rice Res. Inst. Paper Series* 12.

Indonesian Journal of Agricultural Science

Vol. 9 No. 2, 2008

Contents

| | |
|---|----|
| The segregation pattern of insect resistance genes in the progenies and crosses of transgenic Rojolele rice | 35 |
| Satoto, Yuli Sulistyowati, Alex Hartana, and Inez H. Slamet-Loedin | |
| Properties and management implications of soils formed from volcanic materials in Lembang area, West Java | 44 |
| Edi Yatno and Siti Zaayah | |
| Identification and mapping of a gene for rice slender kernel using <i>Oryza glumaepatula</i> introgression lines | 55 |
| Sobrizal and Atsushi Yoshimura | |
| Role of polyamines in inhibition of ethylene biosynthesis and their effects on rice anther culture development | 60 |
| Iswari S. Dewi and Bambang S. Purwoko | |
| Mycotoxin contamination on corn used by feed mills in Indonesia | 68 |
| Budi Tangendjaja, Sri Rachmawati, and Elizabeth Wina | |
| Index | 77 |
| List of referee 2008 | 78 |
