

PROSIDING

SEMINAR NASIONAL HARI NYAMUK 2009

"Partisipasi Masyarakat dalam Program Pengendalian
Nyamuk Terpadu"

Dalam Rangka Hari Nyamuk Nasional 2009



IICC

(IPB International Convention Center) - Botani Square Bogor

Senin, 10 Agustus 2009

Kerjasama

Fakultas Kedokteran Hewan, Institut Pertanian Bogor (FKH- IPB)

Asosiasi Pengendalian Nyamuk Indonesia (APNI)

Unitas Kesehatan Kota (UKK) Bogor

Kwartir Cabang Gerakan Pramuka Kota Bogor

PROSIDING

PROSIDING

SEMINAR NASIONAL

HARI NYAMUK 2009

“Partisipasi Masyarakat dalam Program Pengendalian Nyamuk Terpadu”

Dalam Rangka Hari Nyamuk Nasional 2009

**IPB International Convention Center - Botani Square Bogor
Senin, 10 Agustus 2009**

EDITOR :

**Upik Kesumawati Hadi
Sri Utami Handajani
Risa Tiuria
Susi Soviana**

ISBN 978-602-95733-0-5

PENYELENGGARA



Sekretariat :

**Bagian Parasitologi dan Entomologi Kesehatan, FKH – IPB
Jl. Agatis Kampus IPB Darmaga Bogor, Telp./Fax. 0251 8421784
Email : ukphp_ipb@yahoo.co.id, sugik38@gmail.com**

PROSIDING SEMINAR NASIONAL HARI NYAMUK 2009

“Partisipasi Masyarakat dalam Program Pengendalian Nyamuk Terpadu”

Dalam Rangka Hari Nyamuk Nasional 2009

**ISBN 978-602-95733-0-5
@ 2009**

**Hak Cipta dilindungi Undang-undang
Diterbitkan oleh
Bagian Parasitologi dan Entomologi Kesehatan
Fakultas Kedokteran Hewan
Institut Pertanian Bogor
Jl. Agatis Kampus IPB Darmaga Bogor
Telp/Fax. 0251 8421784
Email : ukphp_ipb@yahoo.co.id**

Desain cover dan layout : Sugiarto

DAFTAR ISI

KATA PENGANTAR	i
DAFTAR ISI	ii
SAMBUTAN KETUA PANITIA	iv
SAMBUTAN REKTOR IPB.....	vi
KEYNOTE SPEAKER	
Program Pengendalian Hama Terpadu.....	1
PLENNARY SESION	13
Konsep Pengendalian Hama Terpadu.....	13
Penelitian Resistensi Vektor di Indonesia (Biomolekuler)	23
Pemberdayaan Masyarakat untuk Mengendalikan Vektor (<i>COMBI Approach</i>)	34
Pengalaman Penerapan Peraturan Daerah (<i>Law Enforcement</i>) dalam Pengendalian DBD di DKI Jakarta.....	44
New Technology for Mosquitoes Control.....	54
MAKALAH BEBAS	
Antimalarial Compounds from <i>Erythrina variegata</i> (Leguminosae)	58
Analisis Faktor Faktor Densitas Larva <i>Aedes aegypti</i> dan Endemisitas Penyakit Demam Berdarah Dengue di Kabupaten Bulukumba, Sulawesi Selatan	67
Pengaruh Komposisi Media Fermentasi <i>Metarhizium</i> sp. terhadap Mortalitas Larva <i>Aedes aegypti</i>	75
Analisis Porsi Substansi Vektor Malaria dalam Petencanaan Pembangunan Kesehatan Tahun 2007-2010 di Kabupaten Kupang Provinsi Nusa Tenggara Timur.....	85
Perilaku Bertelur Nyamuk <i>Aedes aegypti</i> pada Air Sumur Gali dan Air Comberan.....	92

"Partisipasi Masyarakat dalam Program Pengendalian Nyamuk Terpadu"

ii

Penyelenggara



Sebaran Tambak Benur dan Risikonya sebagai Habitat Larva <i>Anopheles</i>	99	✓
Dayaetas Telur <i>Aedes aegypti</i> pada Air Tercemar	107	
Populasi Nyamuk Tersangka Vektor Filariasis di Daerah Endemis Desa Jambu Ilir Kabupaten Ogan Komering Ilir Sumatera Selatan	120	✓
Kandungan Kimia Bawang Putih sebagai Repelen Nyamuk.....	129	
Pengendalian Penyakit DBD dengan Teknik Serangga Mandul (TSM) ..	137	
Habitat Jentik <i>Aedes aegypti</i> (Diptera: Culicidae) pada Air Terpolusi di Laboratorium	143	✓
Sebaran jentik Nyamuk <i>Aedes aegypti</i> (Diptera: Culicidae) di Desa Cikarawang, Kabupaten Bogor.....	154	✓
Pemetaan Tempat Perindukan Larva Nyamuk <i>Anopheles</i> sp. di Kayangan-Lombok Utara, Nusa Tenggara Barat.....	160	✓

LAMPIRAN

Daftar Panitia Penyelenggara.....	167
Daftar Peserta Seminar.....	169



Study of the effect of ultrasonic device against the dengue mosquito, *Aedes aegypti* (Diptera: Culicidae)

Upik Kesumawati Hadi, FX. Koesharto, SH Sigit & Sugiarto

Div. Parasitology and Medical Entomology
Fac. Vet. Med. Bogor Agricultural University, Indonesia
Jl Agatis, Darmaga Campus, Bogor 16680
Telp/Fax 0251 8421784, email upik_hadi@hotmail.com

ABSTRACT - The study of the effect of ultrasonic device against the dengue mosquito, *Aedes aegypti* adults was done in Peet Grady chamber. Three replications of tests were performed based on 24 hours expositions. 50 female mosquitoes on each replication were introduced by ultrasonic device (type ICE) in Peet Grady Chamber, and the other 50 females were put in other Peet Grady Chamber without the device. Observation and counts on knock down mosquitos (mortality rate) were done every one hour to six hours, then 24 hour after exposure. The result showed that During the first hour observation, 2.67% of mosquitoes were knockdown (died) by ultrasonic device (ICE type). In the long run (one to six hours) observation, 3.33 to 10% of the mosquitoes were died, however, after 24 hours exposure 74% of the mosquitoes died. On the control groups, all of the mosquitoes (100%) were still alive on the six hour observations, and after 24 hour observations only 7.33% mosquitoes died. Based on the observation, the ultrasonic device (ICE type) affected the mosquitoes after 24 hours observations by giving them knock down effect and finally died (74%).

KEY WORD: *Aedes aegypti*, knockdown effect, mosquito, ultrasonic.

INTRODUCTION

The recent increase in dengue epidemics all around the world and the real risk of resurgence of urban yellow fever has directed public concern on how to avoid *Aedes aegypti* (L.) and *Aedes albopictus* (Skuse) bites. The avoidance of mosquito bites consequently turns to the obvious- personal protection and how to obtain the perfect repellent or other devices without insecticides. Sonic, electronic and ultrasonic devices have all been advertised as repelling mosquitoes. Some devices are available in a



bracelet, watch or necklace form for personal protection while others plug into room outlets and claim to eliminate mosquitoes from a larger area (Barrido *et al.* 1993). Ultrasound is a form of cyclic sound pressure with a frequency greater than the upper limit of human hearing, this limit being approximately 20 kilohertz (20,000 hertz). The upper frequency limit in humans (approximately 20 kHz) is caused by the middle ear, which acts as a low-pass filter. If ultrasound is fed directly into the skull bone and reaches the cochlea without passing through the middle ear, much higher frequencies can be heard. This effect (sometimes called ultrasonic hearing) was first discovered by divers exposed to high frequencies (ca. 50 kHz) sonar signal. Carefully-designed scientific studies have been performed and confirmed what they call the hypersonic effect - that even without consciously hearing it, high-frequency sound can have a measurable effect on the mind. It is a standard fact in psychoacoustics that children can hear some high-pitched sounds that older adults cannot hear, because in humans the upper limit pitch of hearing tends to become lower with age (Takeda *et al.* 1992).

However, in Indonesia there is no information about the effect of the ultrasonic devices against the mosquitoes. Here, the research was aimed to study the effect of ultrasonic devices against the dengue mosquito, *Aedes aegypti*.

MATERIAL AND METHODS

Product tested was ultrasonic device (ICE type), 20kHz ~ 100kHz, 12 volt, 12 minutes interval produce by LG. The test was carried out in Entomology Laboratory, Division of Parasitology and Medical Entomology, Department of Animal Diseases and Veterinary Public Health, Faculty of Veterinary Medicine, Bogor Agricultural University, Bogor Indonesia. The test was done from February to April 2009. The mosquito used in this study was *Aedes aegypti*, strain Liverpool, female, 4-5 days old, colonized in Entomology laboratory.

The test were done in 190x190x190 cm³ Peet Grady (PG) chambers. Four to five days old *Aedes aegypti* were use for the test. Ultrasonic device (ICE type) was put inside right at the center of the PG and used, waited for one minute. Fifty sucrose-fed females were released into PG chamber, observed for several minutes to make sure that no mosquito fell down to the floor. Observation and counts on knock down mosquitos were done every hour up to six hours, then 24 hours after exposure.

RESULTS AND DISCUSSION

The result of the research was performed in Table 1. The result showed that during the first hour observation, 2.67% of mosquitoes were knockdown (died) by ultrasonic device (ICE type). In the long run (one to six hours) observation, 3.33 to 10% of the mosquitoes were died, however, after 24 hours exposure 74% of the mosquitoes died. On the control groups, all of the mosquitoes (100%) were still alive on the six hour observations, and after 24 hour observations only 7.33% mosquitoes died.

"Partisipasi Masyarakat dalam Program Pengendalian Nyamuk Terpadu"

Penyelenggara



Based on the observation, the ultrasonic device (ICE type) affected the mosquitoes after 24 hours observations by giving them knock down effect and finally died (74%). According to the theory is that rodents and insects can hear sounds that are above the range of human senses (upper than 20 kHz). By emitting a loud *ultrasound*, it's assumed that the mosquitoes will received the effect of ultrasonic wave through their antennae as receptors, and their will be repelled, driven insane or discomfort and finally dead.

Table 1. Mortality rate of mosquitoes after exposed by ultrasonic device (ICE type, 20~100 kHz, 12 volt, 12 minutes interval) in Peet Grady Chamber

Time (hours)	Initial number of mosquitoes	Number of mosquitoes that died	Number of mosquitoes that survived	Mortality rate (%)	Survival rate (%)	Knock down effect (%)	Final result (%)	Final result (%)
1	4.00	0.00	4.00	2.67	0.00	0.00	0.00	0.00
2	6.00	0.00	4.00	3.33	0.00	0.00	0.00	0.00
3	12.00	0.00	4.00	5.33	0.00	0.00	0.00	0.00
4	12.00	0.00	4.00	5.33	0.00	0.00	0.00	0.00
5	14.00	0.00	6.00	6.67	0.00	0.00	0.00	0.00
6	20.00	2.00	8.00	10.00	0.00	0.00	0.00	0.00
24	82.00	56.00	84.00	74.00	6.00	8.00	8.00	7.33

CONCLUSION

Ultrasonic device (ICE type, 20 kHz ~ 100 kHz, 12 volt, 12 minute interval) affected the mosquitoes (*Aedes aegypti*) after 24 hours observations by giving them (74%) knock down effect and finally died.

REFERENCES

- Barrido R., Brown J., Novak R. and Berenbaum B. 1993. A test of the efficacy of ultrasonic mosquito repellents. *The Vector Control Bulletin of the North Central States*. 2:65-69.
- Takeda S *et al.* (1992) Age variation in the upper limit of hearing. *European Journal of Applied Physiology* 65(5), 403-408.



Diskusi :

1. Penanya pertama

- a. Diketahui kontrol hanya 6 jam masa pengujiannya sedangkan yang menggunakan ultrasound hingga 24 jam, mengapa kontrolnya juga tidak diuji sama dengan perlakuan yang menggunakan ultrasound yaitu 24 jam?
- b. Pada kesimpulan tidak mencantumkan efektifitas?

Jawab :

- a. Kelompok kontrol dan perlakuan dengan ultrasound diamati dalam waktu yang sama yaitu 24 jam, sedangkan untuk efektifitas dari kelompok dengan perlakuan adalah 74% sedangkan untuk kontrol hanya 7.33% dalam waktu 24 jam.
- b. Penelitian entomologi berkaitan dengan masalah efikasi alat, sedangkan untuk keamanan digunakan penelitian terhadap monyet yang memiliki fungsi fisiologis hampir menyerupai manusia. Pengaruh gelombang ultrasound dalam jangka panjang dengan FK UNPAD



