

7. DAFTAR PUSTAKA

- Agisimanto D, Supriyanto A. 2007. Keragaman genetik pamelo Indonesia berdasarkan Primer Random Amplified Polymorphic DNA. *J Hort* 17:1-7.
- Altat N, Khan AR. 2007. The seedless trait in kinnow fruit. *Pak J Bot* 39: 2003-2008.
- Arias AB, Ramo'n-Laca L. 2005. Pharmacological properties of citrus and their ancient and medieval uses in the Mediterranean Region. *J Ethnopharmacology* 97: 89-95.
- Ara N, Bashar MK, Kalim Udin MD, Khalequzzaman KM. 2008. Evaluation of pummelo, *Citrus grandis* L. cultivars in northern area of Bangladesh. *J Agric Res* 46: 65-75.
- [BAFPS] Bureau of Agriculture and Fisheries Products Standard. 2004. Philippines National Standard BAFPS BAFPS 11:2004 for Fresh Fruit-Pummelo-Specification. Bureau of Products Standard, Department of Agriculture. Manila.
- Baker, EA, Procopious J, Hunt GM. 1975. The cuticles of *Citrus* species: Composition of leaf and fruit waxes. *J. Sci. Food Agric.* 26, 1093-1101.
- [BPS] Biro Pusat Statistik. 2012. Data Ekspor-Impor. <http://www.bps.go.id/>. [30 Juli 2012].
- [BSN] Badan Standardisasi Nasional. 2009. Standar Nasional Indonesia. SNI 7419:2009. Pamelos. Jakarta: Badan Standardisasi Nasional.
- Chacoff NP, Aizen MA. 2007. Pollination requirements of pigmented grapefruit (*Citrus paradisi* Macf.) from Northwestern Argentina. *Crop Sci* 47:1143-1150.
- Chacoff NP, Souto CP, Aizen MA, Premoli A. 2009. Is there genetic variation in seedless Argentinean grapefruit? Implications for crop production and conservation. *J Basic Appl Gen* 20: 27-35.
- Chaiwong S and Theppakorn T. 2010. Bioactive compounds and antioxidant capacity of pink pummelo (*Citrus grandis* (L.) Osbeck) cv. "Thong dee" in Thailand. *J. ISSAAS Vol. 16, No. 2:10-16*.
- Chavez DJ, Chapparo JX. 2011. Identification of markers linked to seedlessness in *Citrus kinokuni* hort. ex Tanaka and its progeny using bulked segregant analysis. *Hort Sci* 46: 693-697. Abstr.

- Craig EJ. 2002. Phytochemical guardians of our health, J. Amer Dietetic Association, 97:199-204.
- Cvetković BR, Jokanović MR. 2009. Effect of preservation method and storage condition on ascorbic acid loss in beverages. APTEFF, 40:1-7.
- Davies FS, Albrigo LG. 1994. Citrus. UK :CAB International. 254 p.
- de Lima JC, Crippa A, Garrido LR, Camargo UA, de Oliveira PRD. 2006. Molecular analysis of the seedlessness character in grape using RAPD markers. Crop Breeding and Appl Biotechnol 6:209-214.
- [Ditjenhort] Direktorat Jenderal Bina Produksi Hortikultura. 2012. Daerah sentra jeruk. <http://hortikultura.deptan.go.id> [23 Februari 2013].
- Freeman, B. 1978. Cuticular waxes of developing leaves and fruits of citrus and blueberry: Ultra structure and chemistry. University of Florida, Gainesville.
- Gattuso G, Barreca D, Gargiulli C, Leuzzi U and Caristi C. 2007. Review: Flavonoid composition of *Citrus* juices. Molecules 12: 1641-1673.
- Hasegawa S, Lamikl ZJ. 1994. Inhibition of Chemically Induced Carcinogenesis by Citrus Limited. Di dalam Food Phytochemicals for Cancer Prevention I. Fruits and Vegetables M. J. Huang, T. Osawa, C. T. Ho, R. T. Rosen (Eds.) American Chemical Society Washington DC PP. Hal. 209-219.
- Hasegawa S, Berhow MA, Fong CH. 1996. Analysis of Bitter Principles in *Citrus*. Di dalam: Linskens HF, Jackson JF, editor. *Modern Methods of Plant Analysis*, Volume ke-18 *Fruit Analysis*. Berlin Heidelberg: Springer-Verlag. Hlm 59-80.
- Hirai M, Kajiura I. 1987. Genetic analysis of leaf isozymes in citrus. Jpn J Breed 37:377-388.
- Hlušičková JB, Blažek J. 2006. Seed count, fruit quality and storage properties in four apple cultivars. J. Fruit Ornamental Plant Res 14: 151-160.
- Iqbal MM, Altaf N, Murwat EK, Hafiz IA, Bhatti IA. 2001. A marker for seedlessness in Kinnow mandarin. Pak J Biol Sci 4:69.
- [Kementan] Kementerian Pertanian. 2013. Produksi Nasional Hortikultura Tahun 2000-2011. <http://aplikasi.deptan.go.id> [23 Februari 2013].
- Ketsa, S. 1988. Effect of seed number on fruit characteristics of tangerine. Katsersat J. (Nat. Sci.). 22:225-227.
- Ladaniya MS. 2008. Citrus Fruit. Biology, Technology and Evaluation. San Diego: Academic Press. 593 hal.

- Susanto, S. 2002. Pengaruh pelilinan dan pembungkusan dengan plastik berlubang terhadap daya simpan buah jeruk besar 'Nambangan'. *Agrotropika* 7 (1): 27-33.
- Susanto S, Nakajima Y, Hasegawa K. 1993. Effect of Growth Hormone Application on the Development of Parthenocarpic of Pummelo Trees Grown in Plastic House. *Jpn J. Trop. Agr.* 37(4): 284-289.
- Tsai HL, Chang SKC, Chang SJ. 2007. Antioxidant content and free radical scavenging ability of fresh red pummelo (*Citrus grandis* (L.) Osbeck) juice and freeze-dried products. *J Agric Food Chem* 55 (8): 2867-2872. Abstr.
- [UNECE] United Nations Economic Commission for Europe. 2009. UNECE Standard Ffv-14. Concerning the Marketing and Commercial Quality Control of Citrus Fruit. Edisi 2009. United Nations. New York and Geneva.
- Uzun. A, Gulsen O, Yesiloglu T, Aka-Kacar Y, Tucz O. 2010. Distinguishing grapefruit and pummelo accessions using ISSR markers. *Czech J Genet Plant Breed* 46 (4):170-177.
- Van DT, Hai VM, Stackelberg MV, Jacobsen HJ, Dang BQ, Hung LV. 2004. Preliminary results of research for identification the linking between isozymes and seedless characterization of Grape-fruit (Pummelo) in Vietnam *Sci Technol Agric Rural Dev* 8:1082-1084.
- Vardi A, Levin I, Carmi N. 2008. Induction of seedlessness in citrus: from classical techniques to emerging biotechnological approaches. *J Am Soc Hort Sci.* 133: 117-126.
- Varoquaux F, Blanvillain R, Delseny M, Gallois P. 2000. Less is better: new approaches for seedless fruit production. *Tibtech* 18:233-242.
- Yen CR. 1984. Seeded and seedless fruits growth of "Sah Keng" litchi. *J Agric Res China* 33:257-264.
- Yu J, Wang L, Walzem RL, Miller EG, Pink LM, Patil BS. 2005. Antioxidant activity of citrus limonoids, flavonoids and coumarins. *J. Agric. Food Chem.* 53: 2009-2014.
- Zechmeister L, LeRosen AL, Went FW, Pauling L. 1941. Prolycopene, a naturally occurring sterioisomer of lycopene. *Proc Natl Acad Sci USA* 21: 468-474.
- Zhichang Z, Guibing H, Yangruo O, Yunchun L, Yang Y, Yeyuan Y. 2010. The earlier identification of the seedless characteristic of the wampee [*Clausena lansium* (Lour.) Skeels] hybrid by a random amplified polymorphic DNA (RAPD) Marker. *African J Biotechnol* 9:8578-8583.