

Reprinted from the

Far East Journal of
Mathematical Sciences (FJMS)
Volume 95, Number 1, 2014, pp 69-80

SHARPER ANALYSIS OF UPPER BOUND FOR THE
ITERATION COMPLEXITY OF AN INTERIOR-POINT
METHOD USING PRIMAL-DUAL FULL-NEWTON

STEP ALGORITHM

by

Bib Paruhum Silalahi



Pushpa Publishing House
Vijaya Niwas, 198 Mumfordganj
Allahabad 211002, INDIA
<http://pphmj.com/journals/fjms.htm>
fjms@pphmj.com & arun@pphmj.com

Information for Authors

Aims and Scope: The *Far East Journal of Mathematical Sciences* (FJMS) is devoted to publishing original research papers and critical survey articles in the field of *Pure and Applied Mathematics, Computer Applications and Statistics*. The FJMS is a fortnightly journal published in twelve volumes annually and each volume comprises of two issues.

Abstracting, Indexing and Reviews: Global Impact Factor : 0.692, Scopus, Mathematical Reviews, MathSciNet, ProQuest, IndexCopernicus, EBSCOhost, Zentralblatt MATH, Ulrich's web, Indian Science Abstracts, SCIRUS, OCLC, Google Scholar, Excellence in Research for Australia (ERA), AcademicKeys.

Submission of Manuscripts: Authors may submit their papers for consideration in the Far East Journal of Mathematical Sciences (FJMS) by the following modes:

1. **Online submission:** Please visit journal's homepage at <http://www.pphmj.com/journals/fjms.htm>
2. **Electronically:** At the e-mail address: fjms@pphmj.com or kkazad@pphmj.com
3. **Hard copies:** Papers in duplicate with a letter of submission at the address of the publisher.

The paper must be typed only on one side in double spacing with a generous margin all round. An effort is made to publish a paper duly recommended by a referee within a period of three months. One set of galley proofs of a paper will be sent to the author submitting the paper, unless requested otherwise, without the original manuscript, for corrections.

Abstract and References: Authors are requested to provide an abstract of not more than 250 words and latest Mathematics Subject Classification. Statements of Lemmas, Propositions and Theorems should be set in *italics* and references should be arranged in alphabetical order by the surname of the first author.

Page Charges and Reprints: Authors are requested to arrange page charges of their papers @ USD 40.00 per page for USA and Canada, and EUR 30.00 per page for rest of the world from their institutions/research grants, if any. However, for authors in India this charge is Rs. 800.00 per page. No extra charges for colour figures. Twenty-five reprints in print version and a copy in soft version are provided to the corresponding author ex-gratis. Additional sets of reprints may be ordered at the time of proof correction.

Copyright: It is assumed that the submitted manuscript has not been published and will not be simultaneously submitted or published elsewhere. By submitting a manuscript, the authors agree that the copyright for their articles is transferred to the Pushpa Publishing House, Allahabad, India, if and when, the paper is accepted for publication. The publisher cannot take the responsibility of any loss of manuscript. Therefore, authors are requested to maintain a copy at their end.

Subscription Information for 2014

Institutional Price for all countries except India

Electronic Subscription	€ 720.00	US\$ 995.00
Print Subscription includes Online Access	€ 1195.00	US\$ 1695.00

For Institutions: On seeking a license for volume(s) of the *Far East Journal of Mathematical Sciences* (FJMS), the facility to download and print the articles will be available through the institutional 9 digits IP address to be provided by the appropriate authority. The facility to download will continue till the end of the next calendar year from the last issue of the volume subscribed. For having continued facility to keep the download of the same subscribed volume for another two calendar years may be had on a considerable discounted rate.

Price in Indian Rs. (For Indian Institutions in India only)

Print Subscription Only	Rs. 15000.00
-------------------------	--------------

The subscription year runs from January 1, 2014 through December 31, 2014.

Information: The journals published by the "Pushpa Publishing House" are solely distributed by the "Vijaya Books and Journals Distributors".

Contact Person: Subscription Manager, Vijaya Books and Journals Distributors, Vijaya Niwas, 198 Mumfordganj, Allahabad 211002, India; sub@pphmj.com; arun@pphmj.com



Far East Journal of Mathematical Sciences (FJMS)
© 2014 Pushpa Publishing House, Allahabad, India
Available online at <http://pphmj.com/journals/fjms.htm>
Volume 95, Number 1, 2014, Pages 69-80

SHARPER ANALYSIS OF UPPER BOUND FOR THE ITERATION COMPLEXITY OF AN INTERIOR-POINT METHOD USING PRIMAL-DUAL FULL-NEWTON STEP ALGORITHM

Bib Paruhum Silalahi

Department of Mathematics
Faculty of Mathematics and Natural Sciences
Bogor Agricultural University
Jl. Meranti, Kampus IPB Darmaga
Bogor, 16680, Indonesia
e-mail: bibparuhum1@yahoo.com

Abstract

The use of interior-point methods to solve linear optimization problems has become a great attention to the researchers. The most important thing is that the interior-point methods have the best complexity compared to other methods and also efficient in practice. The worst upper bound for the iteration complexity of this method is polynomial. Roos, Terlaky and Vial presented an interior-point method using primal-dual full-Newton step algorithm that requires the best known upper bound for the iteration complexity of an interior-point method. In this paper, we present their method with a slightly better iteration bound.

1. Introduction

Recently, the use of interior-point methods (IPMs) for solving linear Received: July 8, 2014; Revised: August 20, 2014; Accepted: September 3, 2014
2010 Mathematics Subject Classification: 90C05, 90C51.

Keywords and phrases: interior-point method, upper bound, primal-dual full-Newton step.

optimization (LO) problems has become a major concern of the optimization researchers. This happens mainly due to the fact that interior-point methods have polynomial complexity, which is the best compared to other methods and these methods are also efficient in practice.

Interior-point method first appeared in 1984, when Karmarkar [4] proposed a polynomial-time method for LO problems. In the worst-case, for a problem with n inequalities and L bits of input data, Karmarkar's algorithm requires $O(n^{3.5}L)$ arithmetic operations on numbers with $O(L)$ bits.

In [8], Renegar improved the number of iterations to $O(\sqrt{n}L)$ iterations. Other variants of IPMs, called *potential reduction methods*, require also only $O(\sqrt{n}L)$ iterations. This was shown by Ye [13], Freund [1], Todd and Ye [12] and Kojima et al. [5].

Sonnevend [11] and Meggido [6] introduced a class of IPMs which uses the so-called central path as a guide line to the set of optimal solutions. These methods are called path-following methods. A variant of path-following methods was presented by Gonzaga [3], Monteiro and Adler [7] and Roos and Vial [10]. Their methods require $O(\sqrt{n}L)$ iterations, which is the best known upper bound for the iteration complexity of an IPM. Roos et al. in their book [9] obtained the same upper bound by using an algorithm which is a so-called primal-dual full-Newton step algorithm. Their upper bound for the number of iterations is

$$\left\lceil \sqrt{2n} \ln \frac{n\mu^0}{\varepsilon} \right\rceil, \quad (1.1)$$

where ε is the absolute accuracy of the objective function and $\mu^0 > 0$ denotes the initial value of the so-called barrier parameter.

In this paper, by careful analysis, we reduce the upper bound by a factor $\sqrt{2}$. The iteration upper bound that we obtained is

$$\left\lceil \sqrt{n} \log \frac{n\mu^0}{\varepsilon} \right\rceil.$$

2. Primal-dual IPM with Full-Newton Steps

The standard form of an LO problem is as follows:

$$\min\{c^T x : Ax = b, x \geq 0\}, \quad (P)$$

where $c, x \in \mathbb{R}^n$, $b \in \mathbb{R}^m$ and $A \in \mathbb{R}^{m \times n}$. Any LO problem can be transformed into standard form, by introducing additional variables [2]. The problem (P) is often called the *primal* problem. Associated with any LO problem is another LO problem called the *dual* problem, which consists of the same data (A, b, c) arranged in a different way. The dual of (P) is

$$\max\{b^T y : A^T y + s = c, s \geq 0\}, \quad (D)$$

where $s \in \mathbb{R}^n$ and $y \in \mathbb{R}^m$; (D) is called the *dual problem*.

The feasible regions of (P) and (D) are denoted by \mathcal{P} and \mathcal{D} , respectively. The (relative) interiors of \mathcal{P} and \mathcal{D} are denoted by \mathcal{P}° and \mathcal{D}° .

Finding an optimal solution of (P) and (D) is equivalent to solving the following system [9]:

$$\begin{aligned} Ax &= b, \quad x \geq 0, \\ A^T y + s &= c, \quad s \geq 0, \\ xs &= 0, \end{aligned} \quad (2.1)$$

where xs is the component-wise (or Hadamard) product of the vectors x and s and 0 denotes the zero vector. The first line in (2.1) is simply the feasibility constraint for the primal problem (P) and the second line represents the feasibility constraint for the dual problem (D) . The last equation is the so-called complementarity condition.

By using path-following IPM, the complementarity condition in (2.1) is replaced by $xs = \mu e$, where μ is any positive number and e is the all-one vector. This new constraint is referred to as the centering condition with

respect to μ . The resulting system is

$$\begin{aligned} Ax &= b, \quad x \geq 0, \\ A^T y + s &= c, \quad s \geq 0, \\ xs &= \mu e. \end{aligned} \tag{2.2}$$

If system (2.2) has a solution for some $\mu > 0$, then a solution exists for every $\mu > 0$ [9]. This happens if and only if interior point condition (IPC) is satisfied. The solutions are denoted as $x(\mu)$, $y(\mu)$ and $s(\mu)$. We call $x(\mu)$ the μ -center of (P) and $(y(\mu), s(\mu))$ the μ -center of (D) .

When μ runs through $(0, \infty)$, then $x(\mu)$ runs through a curve in \mathcal{P}^o which is called the *central path of* (P) . Similarly, the set $\{(y(\mu), s(\mu)) : \mu \in (0, \infty)\}$ is called the *central path of* (D) . If $\mu \rightarrow 0$, then $x(\mu)$, $y(\mu)$ and $s(\mu)$ converge to a solution of (2.1), which means that the central path converges to the set of optimal solutions of (P) and (D) . On the other hand, if $\mu \rightarrow \infty$, then $x(\mu)$ and $(y(\mu), s(\mu))$ converge to the so-called *analytic center* of (P) and (D) , respectively.

Next, it will describe how Newton's method can be used to obtain an approximate solution of system (2.2), for fixed μ . Given a primal-dual feasible pair $(x, (y, s))$, we want to define search direction Δx , Δy and Δs such that $(x + \Delta x, y + \Delta y, s + \Delta s)$ satisfy (2.2).

Since $Ax = b$ and $A^T y + s = c$, system (2.2) is equivalent to the following system:

$$\begin{aligned} A\Delta x &= 0, \\ A^T \Delta y + \Delta s &= 0, \\ s\Delta x + x\Delta s + \Delta x\Delta s &= \mu e - xs. \end{aligned}$$

The third equation is nonlinear, due to the quadratic term $\Delta x\Delta s$. By neglecting this quadratic term, according to Newton's method for solving nonlinear equations, we obtain the linear system

$$\begin{aligned} A\Delta x &= 0, \\ A^T \Delta y + \Delta s &= 0, \\ s\Delta x + x\Delta s &= \mu e - xs. \end{aligned} \tag{2.3}$$

The resulting directions of (2.3) are known as the primal-dual Newton directions. By taking a step along these directions, one finds new iterates $(x^+, (y^+, s^+))$ such that x^+ and s^+ are positive. The new iterates are given by

$$x^+ = x + \Delta x, \quad y^+ = y + \Delta y, \quad s^+ = s + \Delta s.$$

In the process of following the central path to the optimal solution, by using Newton steps, we generate a sequence of points within the neighborhood of central path. We need a quantity to measure the proximity of $(x, (y, s))$ to the μ -center.

Before defining this proximity measure, we reformulate the linear system (2.3), by scaling Δx , Δy and Δs to d_x , d_y and d_s as follows:

$$d_x = \frac{v\Delta x}{x}, \quad d_y = \frac{\Delta y}{\sqrt{\mu}}, \quad d_s = \frac{v\Delta s}{s},$$

where

$$v = \sqrt{\frac{xs}{\mu}}.$$

If we define $D = \text{diag}(\sqrt{x/s})$, then system (2.3) is equivalent to

$$\begin{aligned} ADd_x &= 0, \\ (AD)^T d_y + d_s &= 0, \\ d_x + d_s &= v^{-1} - v. \end{aligned} \tag{2.4}$$

The first two equations of system (2.4) show that the vectors d_x and d_s belong to the null space and the row space of the matrix AD , respectively. These two spaces are orthogonal, therefore, d_x and d_s are orthogonal. The orthogonality of d_x and d_s implies

$$\begin{aligned}\|d_x\|^2 + \|d_s\|^2 &= \|d_x + d_s\|^2 \\ &= \|v^{-1} - v\|^2.\end{aligned}$$

Note that d_x, d_s (and also d_y) are zero if and only if $v^{-1} - v = 0$, which happens only if $v = e$, and then x, y and s coincide with the respective μ -centers. Therefore, to measure the ‘distance’ of $(x, (y, s))$ to the μ -center, we use the quantity $\delta(x, s; \mu)$ defined by

$$\delta(x, s; \mu) := \delta(v) := \frac{1}{2} \|v^{-1} - v\|^2. \quad (2.5)$$

For any $\tau \geq 0$, the τ -neighborhood of the μ -center is given by the set

$$\{(x, y, s) : x \in \mathcal{P}, (y, s) \in \mathcal{D}, \delta(x, s; \mu) \leq \tau\}.$$

After a full-Newton step, the duality gap at the new iterates always assumes the same value as at the μ -centers, i.e., $(x^+)^T s^+ = n\mu$ (cf. [9, Theorem II.47]). An IPM with full-Newton steps can be described as in Figure 1.

Primal-dual IPM with full-Newton steps

Input:

an accuracy parameter $\varepsilon > 0$;

a proximity parameter τ , $0 \leq \tau < 1$;

strictly feasible (x^0, y^0, s^0) with $(x^0)^T s^0 = n\mu^0$ and $\delta(x^0, s^0; \mu^0) \leq \tau$;

a barrier update parameter θ , $0 < \theta < 1$.

begin

$$x := x^0; s := s^0; y = y^0; \mu = \mu^0;$$

while $n\mu \geq \varepsilon$ do

$$\mu := (1 - \theta)\mu;$$

$$x := x + \Delta x;$$

$$y := y + \Delta y;$$

$$s := s + \Delta s;$$

endwhile

end

Figure 1. Primal-dual IPM with full-Newton steps.

3. Analysis of the Primal-dual IPM with Full-Newton Steps

We first deal with the effect of a full-Newton step on the proximity measure. The next lemma implies that when $\delta(x, s; \mu)$ is small enough then the primal-dual Newton step is quadratically convergent, as stated in Corollary 3.1.

Lemma 3.1 (cf. [9, Theorem II.50]). *If $\delta := \delta(x, s; \mu) \leq 1$, then the primal-dual Newton step is feasible, i.e., x^+ and s^+ are nonnegative. Moreover, if $\delta < 1$, then x^+ and s^+ are positive and*

$$\delta(x^+, s^+; \mu) \leq \frac{\delta^2}{\sqrt{2(1 - \delta^2)}}.$$

Corollary 3.1. *If $\delta := \delta(x, s; \mu) \leq \frac{1}{\sqrt{2}}$, then $\delta(x^+, s^+; \mu) \leq \delta^2$.*

Initially, the duality gap is $n\mu^0$. In each iteration, it is reduced by the factor $1 - \theta$. Using this, one easily proves the following lemma.

Lemma 3.2 (cf. [9, Lemma II.17]). *After at most*

$$\left\lceil \frac{1}{\theta} \log \frac{n\mu^0}{\varepsilon} \right\rceil$$

iterations, the algorithm stops and we have $n\mu \leq \varepsilon$.

We have the following lemma that will be used in proving the next two theorems.

Lemma 3.3 (cf. [9, Lemma II.54]). *Let (x, s) be a positive primal-dual pair and $\mu > 0$ such that $x^T s = n\mu$. Moreover, let $\delta := \delta(x, s; \mu)$ and let $\mu^+ = (1 - \theta)\mu$. Then*

$$\delta(x, s; \mu^+)^2 = (1 - \theta)\delta^2 + \frac{\theta^2 n}{4(1 - \theta)}.$$

The next theorems present iteration bound of the primal-dual IPM with full-Newton steps.

Theorem 3.1. *If $\tau = 1/\sqrt{2}$ and $\theta = 1/\sqrt{n+1}$, then the algorithm requires at most*

$$\left\lceil \sqrt{n+1} \log \frac{n\mu^0}{\varepsilon} \right\rceil$$

iterations. The output is a primal-dual pair (x, s) such that $x^T s \leq \varepsilon$.

Proof. Let us take $\tau = 1/\sqrt{2}$. By using Corollary 3.1, since we have $\delta(x, s; \mu) \leq 1/\sqrt{2}$, after the primal-dual Newton step we have $\delta(x^+, s^+; \mu) \leq 1/2$. Then, after the update of the barrier parameter to $\mu^+ = (1 - \theta)\mu$ with $\theta = 1/\sqrt{n+1}$, by using Lemma 3.3, we get the following upper bound of $\delta(x^+, s^+; \mu^+)^2$:

$$\delta(x^+, s^+; \mu^+)^2 \leq \frac{1 - \theta}{4} + \frac{\theta^2 n}{4(1 - \theta)} = \frac{1}{2}.$$

The last equality follows by substituting $\theta = 1/\sqrt{n+1}$. Hence, we obtain $\delta(x^+, s^+; \mu^+) \leq 1/\sqrt{2} = \tau$. This means that the property

$$\delta(x, s; \mu) \leq \tau$$

is maintained after each iteration. Therefore, combining this with Lemma 3.2, we obtain the theorem. \square

Note that Theorem 3.1 holds for every $n \geq 1$. In practice, n is much larger. For such cases it is worth mentioning that slightly better iteration bounds can be obtained, as the following theorem.

Theorem 3.2. *If $\tau \in [0.6687, 0.6773]$ and $\theta = 1/\sqrt{n}$, then for $n \geq 47$, the algorithm requires at most*

$$\left\lceil \sqrt{n} \log \frac{n\mu^0}{\varepsilon} \right\rceil$$

iterations.

Proof. Let us take $\theta = 1/\sqrt{n}$. By using Lemma 3.1 and Lemma 3.3, we can verify that if

$$\frac{(1 - 1/\sqrt{n})\tau^4}{2(1 - \tau^2)} + \frac{1}{4(1 - 1/\sqrt{n})} \leq \tau^2, \quad (3.1)$$

then $\delta(x, s; \mu) \leq \tau$ is maintained. The region in the (τ, n) -space defined by (3.1) is depicted in Figure 2, where the smallest value of n is around 46.6274 at $\tau = 0.6731$. Therefore, $n = 47$ is the smallest integer value of n which satisfies (3.1). We can find out that for $n \geq 47$, the inequality (3.1) holds for $\tau \in [0.6687, 0.6773]$. Then we have the theorem. \square




Figure 2. The region defined by (3.1).

The value of n in Theorem 3.2 can be improved to $n \geq 6$, as stated in the following theorem.

Theorem 3.3. *If $\tau \in [0.7433, 0.8289]$ and $\theta = 1/\sqrt{n}$, then for $n \geq 6$, the algorithm requires at most*

$$\left\lceil \sqrt{n} \log \frac{n\mu^0}{\varepsilon} \right\rceil$$

iterations.

Proof. We use [9, Theorem II.52], a sharper quadratic convergence result of a primal-dual Newton step. This theorem states that if $\delta := \delta(x, s; \mu) < 1$, then

$$\delta(x^+, s^+; \mu) \leq \frac{\delta^2}{\sqrt{2(1 - \delta^4)}}.$$

Then, by using Lemma 3.3, we obtain that for $\theta = 1/\sqrt{n}$ the property $\delta(x, s; \mu) \leq \tau$ is maintained if

$$\frac{(1 - 1/\sqrt{n})\tau^4}{2(1 - \tau^2)} + \frac{1}{4(1 - 1/\sqrt{n})} \leq \tau^2. \quad (3.2)$$

Figure 3 depicts the region defined by (3.2). The smallest value of n is around 5.1971 at $\tau = 0.7968$, and we can verify that for the smallest integer value $n = 6$, the inequality (3.2) holds for $\tau \in [0.7433, 0.8289]$. Thus, we get the theorem. \square




Figure 3. The region defined by (3.2).

References

- [1] R. M. Freund, Polynomial-time algorithms for linear programming based only on primal scaling and projected gradients of a potential function, *Math. Program.* 51 (1991), 203-222.
- [2] D. Goldfarb and M. J. Todd, Linear programming, *Optimization*, G. L. Nemhauser, A. H. G. Rinnooy Kan and M. J. Todd, eds., *Handbooks in Operations Research and Management Science*, Vol. 1, pp. 141-170, North Holland, Amsterdam, The Netherlands, 1989.

- [3] C. C. Gonzaga, An algorithm for solving linear programming problems in $O(n^3L)$ operations, *Progress in Mathematical Programming: Interior Point and Related Methods*, N. Megiddo, ed., pp. 1-28, Springer-Verlag, New York, 1989.
- [4] N. Karmarkar, A new polynomial-time algorithm for linear programming, *Combinatorica* 4(4) (1984), 373-395.
- [5] M. Kojima, S. Mizuno and A. Yoshise, An $O(\sqrt{n}L)$ iteration potential reduction algorithm for linear complementarity problems, *Math. Program.*, Series A 50 (1991), 331-342.
- [6] N. Megiddo, Pathways to the optimal set in linear programming, *Progress in Mathematical Programming: Interior Point and Related Methods*, N. Megiddo, ed., pp. 131-158, Springer Verlag, New York, 1989. Identical version in: *Proceedings of the 6th Mathematical Programming Symposium of Japan*, Nagoya, Japan, 1986, pp. 1-35.
- [7] R. D. C. Monteiro and I. Adler, Interior-path following primal-dual algorithms. Part I: Linear programming, *Math. Program.* 44 (1989), 27-41.
- [8] J. Renegar, A polynomial-time algorithm, based on Newton's method, for linear programming, *Math. Program.* 40 (1988), 59-93.
- [9] C. Roos, T. Terlaky and J.-Ph. Vial, *Interior Point Methods for Linear Optimization*, Springer, New York, 2006. Second Edition of Theory and Algorithms for Linear Optimization, Wiley, Chichester, 1997.
- [10] C. Roos and J.-Ph. Vial, A polynomial method of approximate centers for the linear programming problem, *Math. Program.* 54 (1992), 295-306.
- [11] G. Sonnevend, An "analytic center" for polyhedrons and new classes of global algorithms for linear (smooth, convex) programming, A. Prékopa, J. Szelezsán and B. Strazicky, eds., *System Modelling and Optimization: Proceedings of the 12th IFIP-Conference held in Budapest, Hungary, September 1985*, Vol. 84 of *Lecture Notes in Control and Information Sciences*, pp. 866-876, Springer-Verlag, Berlin, West Germany, 1986.
- [12] M. J. Todd and Y. Ye, A centered projective algorithm for linear programming, *Math. Oper. Res.* 15 (1990), 508-529.
- [13] Y. Ye, A class of potential functions for linear programming, *Management Science Working Paper 88-13*, Dept. of Management Science, University of Iowa, Iowa City, IA 52242, USA, 1988.

Editorial Board

Gunjan Agrawal, India

Natig M. Atakishiyev, Mexico

Antonio Carbone, Italy

Hasan Coskun, USA

Zhenlu Cui, USA

Manav Das, USA

Shusheng Fu, China

Wei Dong Gao, China

Toshio Horiuchi, Japan

Lisa M. James, USA

Koji Kikuchi, Japan

Victor N. Krivtsov, Russian Federation

Alison Marr, USA

Jong Seo Park, South Korea

Alexandre J. Santana, Brazil

A. L. Smirnov, Russian Federation

Chun-Lei Tang, China

Carl A. Toews, USA

Vladimir Tulovsky, USA

Qing-Wen Wang, China

Peter Wong, USA

Pu Zhang, China

George S. Androulakis, Greece

Carlo Bardaro, Italy

Yong Gao Chen, China

Claudio Cuevas, Brazil

Maslina Darus, Malaysia

Massimiliano Ferrara, Italy

Salvatore Ganci, Italy

Demetris P. K. Ghikas, Greece

Jay M. Jahangiri, USA

Young Bae Jun, South Korea

Hideo Kojima, Japan

Yangming Li, China

Manouchehr Misaghian, USA

Cheon Seoung Ryoo, South Korea

K. P. Shum, China

Ashish K. Srivastava, USA

E. Thandapani, India

B. C. Tripathy, India

Mitsuru Uchiyama, Japan

G. Brock Williams, USA

Chaohui Zhang, USA

Kewen Zhao, China

Principal Editor
Azad, K. K. (India)

OUR PUBLICATIONS

1. Advances and Applications in Discrete Mathematics (GIF : 0.378)
2. Advances and Applications in Fluid Mechanics (GIF : 0.689 ; SCOPUS)
3. Advances and Applications in Statistics (GIF : 0.812)
4. Advances in Computer Science and Engineering (GIF : 0.705)
5. Advances in Differential Equations and Control Processes (GIF : 0.607)
6. Advances in Fuzzy Sets and Systems (GIF : 0.703)
7. Current Development in Oceanography (GIF : 0.587)
8. Far East Journal of Applied Mathematics (GIF : 0.727)
9. Far East Journal of Dynamical Systems (GIF : 0.685)
10. Far East Journal of Electronics and Communications (GIF : 0.701 ; SCOPUS)
11. Far East Journal of Mathematical Education (GIF : 0.604)
12. Far East Journal of Mathematical Sciences (FJMS) (GIF : 0.692 ; SCOPUS)
13. Far East Journal of Theoretical Statistics (GIF : 0.767)
14. JP Journal of Algebra, Number Theory & Applications (GIF : 0.681 ; SCOPUS)
15. JP Journal of Biostatistics (GIF : 0.581)
16. JP Journal of Fixed Point Theory and Applications (GIF : 0.624)
17. JP Journal of Geometry and Topology (GIF : 0.725 ; SCOPUS)
18. JP Journal of Heat and Mass Transfer (GIF : 0.604 ; SCOPUS)
19. International Journal of Functional Analysis, Operator Theory and Applications (GIF : 0.597)
20. International Journal of Materials Engineering and Technology (GIF : 0.801)
21. International Journal of Numerical Methods and Applications (GIF : 0.619)
22. Universal Journal of Mathematics and Mathematical Sciences (GIF : 0.564)
23. Surveys in Mathematics and Mathematical Sciences (GIF : 0.558)
24. International Journal of Nutrition and Dietetics

New Journals From 2014

25. Advanced Studies in Artificial Intelligence
26. Advances and Applications in Software Engineering
27. Journal of Advanced Studies in Telecommunications
28. Universal Journal of Applied Physics and Technology
29. Universal Journal of Economic and Business Management
30. Universal Journal of Electrical Engineering and Informatics
31. Universal Journal of Mechanical Sciences and Technology